



This is a digital copy of a book that was preserved for generations on library shelves before it was carefully scanned by Google as part of a project to make the world's books discoverable online.

It has survived long enough for the copyright to expire and the book to enter the public domain. A public domain book is one that was never subject to copyright or whose legal copyright term has expired. Whether a book is in the public domain may vary country to country. Public domain books are our gateways to the past, representing a wealth of history, culture and knowledge that's often difficult to discover.

Marks, notations and other marginalia present in the original volume will appear in this file - a reminder of this book's long journey from the publisher to a library and finally to you.

Usage guidelines

Google is proud to partner with libraries to digitize public domain materials and make them widely accessible. Public domain books belong to the public and we are merely their custodians. Nevertheless, this work is expensive, so in order to keep providing this resource, we have taken steps to prevent abuse by commercial parties, including placing technical restrictions on automated querying.

We also ask that you:

- + *Make non-commercial use of the files* We designed Google Book Search for use by individuals, and we request that you use these files for personal, non-commercial purposes.
- + *Refrain from automated querying* Do not send automated queries of any sort to Google's system: If you are conducting research on machine translation, optical character recognition or other areas where access to a large amount of text is helpful, please contact us. We encourage the use of public domain materials for these purposes and may be able to help.
- + *Maintain attribution* The Google "watermark" you see on each file is essential for informing people about this project and helping them find additional materials through Google Book Search. Please do not remove it.
- + *Keep it legal* Whatever your use, remember that you are responsible for ensuring that what you are doing is legal. Do not assume that just because we believe a book is in the public domain for users in the United States, that the work is also in the public domain for users in other countries. Whether a book is still in copyright varies from country to country, and we can't offer guidance on whether any specific use of any specific book is allowed. Please do not assume that a book's appearance in Google Book Search means it can be used in any manner anywhere in the world. Copyright infringement liability can be quite severe.

About Google Book Search

Google's mission is to organize the world's information and to make it universally accessible and useful. Google Book Search helps readers discover the world's books while helping authors and publishers reach new audiences. You can search through the full text of this book on the web at <http://books.google.com/>

the 1990s, the number of people in the world who are undernourished has increased from 600 million to 800 million.

There are a number of reasons for this increase. First, the world population has increased from 5 billion in 1987 to 6 billion in 1999, and is projected to reach 8 billion by 2025. Second, the number of people who are undernourished has increased from 1.2 billion in 1987 to 1.6 billion in 1999. Third, the number of people who are undernourished has increased from 1.2 billion in 1987 to 1.6 billion in 1999. Fourth, the number of people who are undernourished has increased from 1.2 billion in 1987 to 1.6 billion in 1999.

There are a number of reasons for this increase. First, the world population has increased from 5 billion in 1987 to 6 billion in 1999, and is projected to reach 8 billion by 2025.

Second, the number of people who are undernourished has increased from 1.2 billion in 1987 to 1.6 billion in 1999.

Third, the number of people who are undernourished has increased from 1.2 billion in 1987 to 1.6 billion in 1999.

Fourth, the number of people who are undernourished has increased from 1.2 billion in 1987 to 1.6 billion in 1999.

Fifth, the number of people who are undernourished has increased from 1.2 billion in 1987 to 1.6 billion in 1999.

Sixth, the number of people who are undernourished has increased from 1.2 billion in 1987 to 1.6 billion in 1999.

Seventh, the number of people who are undernourished has increased from 1.2 billion in 1987 to 1.6 billion in 1999.

Eighth, the number of people who are undernourished has increased from 1.2 billion in 1987 to 1.6 billion in 1999.

Ninth, the number of people who are undernourished has increased from 1.2 billion in 1987 to 1.6 billion in 1999.

Tenth, the number of people who are undernourished has increased from 1.2 billion in 1987 to 1.6 billion in 1999.

Eleventh, the number of people who are undernourished has increased from 1.2 billion in 1987 to 1.6 billion in 1999.

Twelfth, the number of people who are undernourished has increased from 1.2 billion in 1987 to 1.6 billion in 1999.

Thirteenth, the number of people who are undernourished has increased from 1.2 billion in 1987 to 1.6 billion in 1999.

Fourteenth, the number of people who are undernourished has increased from 1.2 billion in 1987 to 1.6 billion in 1999.

Fifteenth, the number of people who are undernourished has increased from 1.2 billion in 1987 to 1.6 billion in 1999.

Sixteenth, the number of people who are undernourished has increased from 1.2 billion in 1987 to 1.6 billion in 1999.

Seventeenth, the number of people who are undernourished has increased from 1.2 billion in 1987 to 1.6 billion in 1999.

Eighteenth, the number of people who are undernourished has increased from 1.2 billion in 1987 to 1.6 billion in 1999.

Nineteenth, the number of people who are undernourished has increased from 1.2 billion in 1987 to 1.6 billion in 1999.

Twentieth, the number of people who are undernourished has increased from 1.2 billion in 1987 to 1.6 billion in 1999.



THE STUDENT'S GUIDE
TO
CLINICAL MEDICINE
AND
CASE-TAKING

BY
FRANCIS WARNER, M.D. LOND., F.R.C.P.

ASSISTANT PHYSICIAN, LECTURER ON BOTANY, AND LATE MEDICAL REGISTRAR TO THE
LONDON HOSPITAL; LATE PHYSICIAN TO THE EAST LONDON HOSPITAL FOR CHILDREN
EXAMINER TO THE UNIVERSITY OF ABERDEEN

SECOND EDITION



LONDON
J. & A. CHURCHILL
11, NEW BURLINGTON STREET

1885

C

1512 . f . 6 .

INTRODUCTION TO SECOND EDITION.



FURTHER experience gained since the publication of the former edition, has shown that students commencing clinical work need to be taught to think, and reason for themselves, as well as to observe. It is hoped that the plan of this work tends to give such training if properly used.

The size of this work has not been increased, but new material has been added, and corrections have been made in accordance with the advances of clinical medicine, especially in the chapter on diseases of the nerve-system. A special scheme for taking notes of children has been added, and the index has been made more complete.

F. W.

24, HARLEY STREET, W.

December, 1884.



INTRODUCTION TO FIRST EDITION.

DURING the three years that I held the office of Medical Registrar to the London Hospital, I saw that the student, on commencing his duties as clinical clerk, required some guide as to the method of arranging the history of his case, and the facts observed. A card of "instructions for case-taking" was provided, almost similar to that given at page xi. It was further evident that with each case the student needed, when taking his cases, to be told what special points to note in the history, and what special points to look to under each of the heads of the "instructions." Further, zeal was much stimulated in the thoughtful student by telling him why these special points should be enquired for, and their presence or absence noted.

Such points, with regard to the more commonly recurring diseases, have been put together, and presented in the following pages.

The object has been to provide, in a small space, a guide for the student to use at the bedside, when wanting to know what to look for, and what to note. Pathology and treatment are not touched upon, and for this reason, independent of the general incompleteness of this little work, the student is recommended to read, in some text-book, all about his case in hand. Much attention has been given to the special conditions met with in disturbance and disease of the nervous system.

To encourage enquiry as to the origin of disease, the principal causes in each case are indicated under the heading "*causation*," which will usually be found on the left-hand page, sometimes on the right-hand ; thus the student may find his enquiries directed on reasonable grounds. As to the scheme of the work, as far as possible the facts indicated as specially to be observed are arranged under the ordinary heads of a case on the left-hand page, and on the corresponding right-hand page are given explanations, characters of the special disease, or points of interest in its natural history, etc. This plan could not in all cases be adhered to, and general convenience and the necessities of printing had then to take precedence of the original scheme.

Names printed in the text in **thick type** are heads of chapters contained in the work, which may be found on reference to the index. Thus, in taking a case of fever, look for "**Signs of Fever**," and if **Vomiting** or **Jaundice** be present look up these heads by means of the index.

F. W.

24, HARLEY STREET, W.
1881.

CONTENTS.

	PAGE
INSTRUCTIONS FOR CASE-TAKING	xi—xiii
ADDITIONAL INSTRUCTIONS FOR CHILDREN'S CASES	xiv, xv
TABLE SHOWING HEALTHY DEVELOPMENT OF AN INFANT	xvi

GENERAL DISEASES—CLASS I.

Common bad Hygienic Conditions—Signs of Fever—
Table of the Specific Fevers: Enteric Fever; Typhus;
Scarlet Fever; Measles; Variola; Varicella—Erysipelas
—Diphtheria—Pyæmia—Puerperal Fever—Ague—
Hooping-cough—Syphilis, inherited; Syphilis, acquired 1—17

GENERAL DISEASES—CLASS II.

Signs of Defective Development—Anæmia; Pernicious
Anæmia—Cancer—Rickets—Emaciation—Œdema or
Anasarca—Amyloid Degeneration—Scrofulosis: General
Miliary Tuberculosis—Table of Differences and Resem-
blances between Enteric Fever and Tuberculosis—
Diabetes Mellitus—Addison's Disease—Purpura—De-
velopmental Defects—Senile Degeneration 19—33

ARTHRITIC DISEASES.

Arthritis—Rheumatoid Arthritis—Table of Joints for
Description—Rheumatism; Gonorrhœal Rheumatism—
Gout 34—39

DISEASES OF THE NERVOUS SYSTEM.

General Conditions of the Nervous System—Intel- ligence—Speech—Sleep—Head-pain—Headache— Vomiting—Coma—Vertigo—Delirium—Typhoid State —Paralysis—Tendon Reflexes—Skin Reflexes—Table of Diagnosis of Functional from Organic Paralysis—Elec- tric Tests—Convulsion—Spasm—Laryngismus—Tremor —Athetosis—Motor Power—Sensation—Muscular Anæ- sthesia—Special Senses—Cranial Nerves—Bell's Paralysis of the Face—Brain Disease—Ophthalmoscopic Appear- ances—Pupils—Signs of Spinal Cord Disease—Minor Para- lyses: Progressive Muscular Atrophy; Pseudo-hyper- trophic Paralysis; Paralysis of Extensors of Forearm; Cross Paralysis; Bulbar Paralysis; Paralysis of Face— Neuralgia: Sciatica; Intercostal Neuralgia—Hemiplegia —Chorea—Table of Diagnosis of Chorea from Sclerosis— Hysteria; Epilepsy; Table of Diagnosis of Epilepsy from Hysteria—Cerebral Tumour—Cerebral Meningitis— Chronic Hydrocephalus—Table of Diagnosis of Hydro- cephalus from Rickets—Alcoholism—Acute Alcoholism— Delirium Tremens—Insanity—General Paralysis of the Insane—Paralysis Agitans—Sclerosis of the Cord—Te- tanus—Locomotor Ataxy—Infantile Paralysis—Graves' Disease—Plumbism—Diphtheritic Paralysis—Herpes Zoster	40—101
--	--------

DISEASES OF THE VASCULAR SYSTEM.

Physical Examination of the Heart: Pulse; Passive Congestion—Important Anastomoses—Table of Signs of Valvular Lesions, Mitral Regurgitation and Obstruc- tion; Aortic Regurgitation and Obstruction; Hyper- trophy and Dilatation of the Heart—Cardiac Displace- ments—Valvular Disease—Heart Disease—Table of Diagnosis of Functional from Organic Palpitation—Angina Pectoris—Pericarditis—Congenital Defects of the Heart —Tricuspid Regurgitation—Thoracic Aneurism—Disease of Vessels	102—123
---	---------

DISEASES OF THE RESPIRATORY SYSTEM.

PAGE

Clinical Regions of the Chest—Physical Examination of the Chest—Auscultation—Cough—Sputum—Hæmoptysis; Table of Diagnosis of Hæmoptysis from Hæmatemesis—Dyspnoea—Pulmonary Œdema—Contraction of Lung—Solidification of Lung—Table of Diagnosis of Pneumonia from Pleuritic Effusion—Pleurisy; Empyema; Hydrothorax—Phthisis—Pneumonia—Emphysema—Bronchitis—Asthma—Laryngeal Diseases. 124—149

DISEASES OF THE DIGESTIVE SYSTEM.

Signs of Digestive Functions: Appetite—Intestinal Worms—Examination of the Mouth and Throat: Tongue; Palate; Tonsils; Pharynx; Teeth; Gums—Diarrhoea—Vomiting—Acute Abdominal Pain—Dysphagia—Hæmatemesis—Melæna—Obstruction of the Bowels—Gastric Ulcer—Typhlitis—Abdominal Cancer: Cancer of Stomach; Cancer of Intestines—Ulceration of Bowels: Dysentery; Tubercular Ulceration—Abdominal Tumours: Ovarian; Kidney; Spleen; Abdominal Aneurism; Tumours arising from the Pelvis—Fæcal Accumulations—Phantom Tumour—Intussusception—Examination of Abdomen—Fluid in Peritoneum—Diagnosis of Ovarian Tumour from Ascites—Peritonitis. 150—173

DISEASES OF THE LIVER.

Jaundice; Table of Causation; Jaundice from Obstruction or Independent of Obstruction—Large Livers: Lardaceous, Fatty, Hydatid—Cancer of the Liver—Small Livers—Acute Yellow Atrophy—Cirrhosis of the Liver—Syphilitic Disease of the Liver—Gall-Stones; Biliary Colic—Hydatid of Liver 174—182

DISEASES OF THE URINARY SYSTEM.

PAGE

Bright's Disease — Uræmia — Albuminuria — Hæmaturia—Paroxysmal Hæmaturia—Acute Bright's Disease—Granular Contracted Kidneys—Fatty Kidneys—Amyloid and Large White Kidneys—Bright's Disease, Acute or Chronic—Diseases of the Bladder—Renal Calculus ; Renal Colic—Description of Urine ; Examination of the Deposit, Chemically and Microscopically—Normal Constituents of the Urine ; Abnormal Constituents ; Albumen, Sugar, Bile, Leucine, Tyrosine—Urinary Calculi—Murexide Test for Uric Acid	184—203
SIGNS OF PREGNANCY — COINCIDENT SIGNS AND SYMPTOMS	204, 205
INDEX	206—211

INSTRUCTIONS FOR CASE-TAKING.

- I. Enter name, age, occupation, address, date of admission to hospital, and date at which the notes were taken.
- II. State what the patient complains of, as far as possible using his own words. With **children** say what the friends complain of.
- III. Family History.—Number and condition of health of those living. Ages and diseases of those dead. Specially enquire as to points in the inheritance bearing on the case and its causation.

Personal History.—Habits, occupations, residences, previous illnesses and diseases. Indications of **scrofula**, **gout**, **rickets**, **syphilis**, etc. Give dates.

History of Present Illness.—Date and manner of commencement; date when last at work. Order of the occurrence of symptoms, with date. Indicate the day of illness on the temperature chart. In taking this history look up the causation and course of the disease as given in the text. Probable causes.

- IV. Present Condition.—General condition. **Intelligence**; mental state; **sleep**; complaints of pain, etc. **Nutrition**; emaciation; **anæmia**; **œdema**; complexion; any specially obvious abnormal condition or source of distress, etc. Position of patient in bed; orthopnœa; dorsal decubitus; etc. Pulse= ; Temperature= ; Respirations= ; Weight= .

Lymphatic Glands in neck, axilla, groins; size, hardness, mobility; tendency to suppuration.

Locomotor System.—State of bones, muscles, joints, scars, nodes. Skin, dry or moist; bed-sores.

V. Nervous System.—**General Condition.** **Intelligence;** sleep; speech. **Vertigo;** head-pain. **Delirium;** paralysis; convulsion; tremor; coma, etc.

Motor Power.—Ability to stand or work; movements of extremities; gait in walking; co-ordination of limbs. **Reflexes.**

Sensibility.—Tactile sensibility of skin; sensibility to heat and cold, also to pricking. Anæsthesia; hyperæsthesia; dysæsthesia. **Special senses.**

Cranial Nerves.—Movements of eyes, tongue, palate, face. State of pupils. Ophthalmoscopic examination.

VI. Vascular System.—**Pulse**, frequency and other characters; condition of the vessels, especially the arteries. Cyanosis. **Heart;** palpate, auscultate, percuss. Note precordial dulness if normal. **Palpitation**, pain or signs of heart disease.

VII. Respiratory System.—Dyspnoea, frequency and characters of the respiratory movements. Cough; **expectoration;** hæmoptysis.

Physical Examination.—Inspection; palpation; percussion; auscultation. Signs of bulging or contraction of chest or solidification of lungs, etc. Larynx.

VIII. Digestive System.—**Tongue;** teeth; throat. Appetite; thirst. **Vomiting;** hæmatemesis; melaena. State of bowels; tenesmus; griping; piles. Fulness or pain after food; flatulence; pyrosis; colic or other disturbance. Abdominal pain or tenderness.

Liver.—Size and general characters as determined by percussion and palpation: whether tender or not, **Jaundice**.

Spleen.—Size as determined by percussion and palpation.

Abdomen.—Physical examination. Whether tender, distended, retracted. **Ascites**. **Tumour**.

IX. Urinary System.—**Urine**, quantity, colour, reaction, Sp. gr. Albumen, bile, sugar. Deposit, its general, chemical, and microscopical characters. Frequency of micturition; if accompanied by pain; hæmaturia.

X. Generative System.—Menstruation: frequency; duration; quantity increased or otherwise; whether painful; other discharges. Conditions of uterus and pelvic organs.

XI. Treatment.—Prescriptions and diet, etc., should be entered in the notes, and all alterations noted, with the dates.

XII. Diagnosis.—Should enumerate the principal disease, secondary lesions, complications and specially important conditions, symptoms or points in the treatment.

ADDITIONAL INSTRUCTIONS FOR CHILDREN'S CASES.

II. State complaints made concerning the child, or obvious conditions of disease.

III. Family History.—Number and condition of health of those living. Ages and diseases of those dead. Specially enquire as to the inheritance bearing on the case. History of mother's health during the intra-uterine life of the child. Note any miscarriages, with dates.

Personal History.—Whether healthy at birth; how brought up; if suckled; if farinaceous food has been used. Previous illnesses, and diseases.

IV. Present Condition.—*General condition*: plumpness; skin elastic, clear or muddy looking, with aged appearance.

Condition of Development.—Anæmic; hæmorrhagic flea bites; sweating. Bones, feel them all while the child is stripped. Signs of **syphilis**, **ricketts**, etc. Note warmth of the limbs; whether the child sheds tears in crying. T. = ; P. = ; R. = ; W. = . Signs of **defective development**.

V. Nervous System.—General condition. Note the amount of movement of limbs, hands, and feet, or whether this is absent. Intelligence, as indicated by movements of face and eyes directed towards objects noticed. Sleep; making noises; consciousness; exhaustion; coma. Paralysis; examine each limb. Spasm; tremor; contraction.

Motor Power.—Reflex action on tickling hands, putting finger in mouth, etc. Playfulness; ability to laugh. Power over large joints, small joints, movements of fingers, etc.

Cranial Nerves.—Movements of eyes and face.

Head.—Its shape and circumference. Fontanelle is patent, prominent, or depressed. State of other sutures. Ophthalmoscope.

VI. Vascular System.—Pulse: frequency and character. Cyanosis. Heart: palpate, percuss, auscultate.

VII. Respiratory System.—Dyspnoea; frequency of respiratory movements; laryngeal stridor; spasm, or obstruction. Warmth or coldness of breath. Cough.

Physical Examination.—Inspection; signs of collapse at bases and clavicular regions. [To examine back, let child be held leaning over nurse's shoulder.] Palpation; rhonchi may sometimes be felt. Percussion. Auscultation. In children, and especially infants, the feeling of resistance of lung, or its elasticity beneath the finger struck, gives valuable information as to its consolidation or clearness.

VIII. Digestive System.—Tongue, lips, throat; state of dentition. Appetite and liking for food; how it is fed. Vomiting. State of bowels. Abdomen: whether full or empty; palpate; note size of liver and spleen. State of umbilicus. Pain after food; flatulence; abdominal tenderness; griping of bowels. Test milk used for cream and acidity.

In examining an infant, it is necessary to determine if it be well developed. See **Developmental Defects.**

The child should be weighed, and the circumference of the head, at its longest, should be measured.

xvi ADDITIONAL INSTRUCTIONS FOR CHILDREN'S CASES.

The following Table is for a healthy well-developed infant of good-class:—

Infant at birth weighs six to ten pounds ; head circumference, 11·15—12·5 inches.

Age in months.	Weight.	Head Circumference.	Points indicating Stage and Progress of Development.
	lbs.	ins.	
I.	7 to 10	14·5	Power to suck ; regular succession of feeding and sleeping ; hand reflex.
II.	11·0	15·25	Hair in eyelashes and eyebrows ; may be occasional strabismus.
III.	13·5	16·5	Capability of shedding tears ; no strabismus.
IV.	15·0	17·0	Constant movement while awake.
V.	15·5	17·0	Turning head to a light or sound.
VI.	16·0	17·25	Recognizing objects, as mother, nurse.
VII.	17·5	17·5	Holding object in hand, and carrying it to mouth.
VIII.	18·5	17·75	Various sounds made ; commencing dentition.
IX.	19·5	18·0	Some power to hold up head when lying down.
X.	19·6	18·25	Holding an object without dropping it.
XI.	19·7	18·4	Power to transfer object from one hand to the other.
XII.	20·0	18·5	Commencing to crawl or stand with assistance.

THE
STUDENT'S GUIDE TO CLINICAL MEDICINE.

GENERAL DISEASES—CLASS I.

Specific diseases caused by some poison received by the patient from without, and in many cases communicable from one patient to another.

COMMON BAD HYGIENIC CONDITIONS.

- Drawing drinking-water from a cistern over w.c.
- Overflow pipe from a cistern opening into a sewer or soil-pipe instead of into open air.
- Drain from a kitchen sink opening direct into a sewer or cess-pit instead of into open air over a drain.
- Want of proper traps cutting off house-sewer from street-sewer, and each soil-pipe from house-drain.
- Want of ventilation of house-drains and soil-pipes, with arrangements for access of air into them, and exit of sewer-gas from them, may cause foul smells.

FEVER, SIGNS OF.

General condition.—Temperature raised ; respirations and pulse frequent ; skin dry and hot, or sweating ; rigors ; fever pains ; aching in back and limbs ; prostration of muscular power ; face presents depressed look. P. = ; T. = ; R. = .

Mode of onset.—Sudden, with rigors, headache, pains in back and limbs ; gradual, with anorexia and thirst, loss of strength.

Digestion.—Anorexia ; thirst ; bowels confined or relaxed ; describe the motions passed. **Tongue** furred, dry, or moist ; papillæ may be enlarged. State of gums ; teeth. Throat ; tonsils. **Vomiting.** *Spleen* may be enlarged. *Liver*, see Jaundice.

Vascular system.—Pulse frequent, soft, may be dicrotous and intermittent. Heart's action quick ; note strength of impulse and first sound. Tendency to capillary congestion.

Respiratory system.—Respirations frequent ; tendency to congestion of the lungs. **Pulmonary Oedema ; Bronchitis ; Pneumonia ; Pleurisy.**

Nervous system.—Mental condition, see general condition of **Nervous System.** Sleep ; **Headache ; Delirium ; Typhoid State.**

Urine.—Scanty. Sp. gr. high. Commonly a deposit of **Lithates.** It may be jaundiced or albuminous. Urea may be in excess.

Look for rash and the characters of the **Specific Fevers** ; local and general complications.

FEVER, SIGNS OF.

General condition.—An exanthematous fever does not often recur in the same individual. The date and mode of onset are important, so also whether sudden or gradual, with or without rigors.

Digestion.—Sordes and accumulations of mucus may occur on lips and teeth. Note any inability to take food or to swallow. Jaundice is common in relapsing fever, and may be present with **Pyæmia**, **Typhus**, etc.

Vascular system, see **Pericarditis**.—Danger may arise from failure of heart's action, and weakness of the circulation. Note complexion of the lips and face, fulness and tension of pulse.

Respiratory system.—Note fulness or shallowness of respirations. Examine lungs frequently, even if there be no symptoms of their disturbance. Note cough or **Expectoration**.

Nervous system.—Mental condition disturbed; delirium not necessarily of bad prognosis. Hyperpyrexia and adynamia dangerous.

Urine.—**Albuminuria** may be temporary, or it may lead to chronic Bright's disease.

Look for causation; cold, contagion in case of specific fevers, bad **hygienic conditions**.

Causation.—Contagion; concurrent or previous illness in same house; smells from sewers; water supply; date of exposure to contagion; infection by clothes.

SPECIFIC ERUPTIVE FEVERS.**DAYS OF FEVER AND RASH.**

ENTERIC FEVER.—Onset gradual; temperature slowly rising, falling at end of 3rd or in the 4th week with exacerbations at night. Small oval hyperæmic spots on abdomen in successive crops in 2nd and early in 3rd weeks.

SIGNS AND SYMPTOMS.

Abdominal pain and tenderness; gurgling over cæcum. Bowels usually relaxed. **Spleen** large. Temperature may be excessive. Occasionally sudamina. **Bronchitis** common. Bowels may be constipated.

TYPHUS FEVER.—Onset severe, with rigors and pains in back and limbs. Temperature rises rapidly 4 to 5 days, falling about 14th day. Mulberry-coloured maculæ, at first slightly raised, then dull mottling, appear in 1st week, disappear end of 2nd week.

Headache and nervous symptoms prominent; delirium usual. Much tendency to heart failure and hypostatic congestions. **Bronchitis**. Bowels not usually relaxed.

SCARLET FEVER.—Onset rather sudden, with chilliness. Temperature rising rapidly. Rash 2nd day on neck, chest, and trunk, extending to the limbs; minute red points, quickly becoming a diffused erythema. Rash passes off about 7th day, leaving desquamation of skin. Temperature falls about same time.

Tongue thickly coated, with enlarged red papillæ protruding; tip quickly becoming red. Fauces inflamed; tonsillitis. Desquamation specially seen on hands and feet. Occasionally there is no rash. Delirium.

SPECIFIC ERUPTIVE FEVERS.**COMPLICATIONS.**

ENTERIC FEVER.—Signs of heart failure. **Delirium.** **Typhoid State.** Hypostatic congestion of lungs. **Albuminuria.** Hæmorrhage from bowels; perforation of intestine. Profuse sweating, see **Tuberculosis.** Phlebitis. Sequential abscesses. Tendency to relapses of fever and other symptoms.

TYPHUS FEVER.—Active **Delirium** passing into the **Typhoid State.** Hypostatic pneumonia. **Albuminuria.** Weak action of ventricles, and very soft pulse.

CAUSATION.

Impure water. Sewer gas. Probably not contagious, but by the evacuations. Note occupation; residence, and its hygienic condition; sources of milk supply.

Contagious from the sick to the healthy. Its spread is favoured by over-crowding, bad hygienic conditions, and starvation.

SCARLET FEVER.—Albuminuria and anæmia with anasarca from **Acute Bright's Disease.** Inflammation of the throat may be excessive, with ulceration. Arthritis. Scarlatinal rheumatism. Inflammation of the middle ear. Glandular abscess in neck. Hyperpyrexia. **Pleurisy** or empyema rather than pneumonia. **CONVULSIONS.** *Scarlet Fever.*—Rheumatic symptoms often commence at the beginning of the 2nd week with swelling in sheaths of tendons, and some redness, tenderness, and moisture of skin. Subsequently stiff neck not uncommon.

Highly infectious, especially through the dust of the skin. The type varies greatly in different epidemics; in some, greater tendency to complications or death.

SPECIFIC FEVERS.

DAYS OF FEVER AND RASH.

MEASLES.—Rash appears about 4th day ; begins on face, spreading to the trunk and limbs. Fine red points, becoming flat and forming crescentic patches. Temperature begins to fall two or three days after rash appears.

SIGNS AND SYMPTOMS.

Specially common in children. Onset with chills or rigors. Sleepiness. Catarrh ; conjunctivæ watery ; coryza. If rash is full, desquamation may follow.

VARIOLA.—Rash appears 3rd day, first on forehead as red papules, soon becoming vesicles, feeling hard as shot ; 5th day they become umbilicated and purulent ; 8th day pustules mature, then scab. Temperature rises rapidly ; falls as rash appears ; secondary fever with the suppuration.

Incubation after inoculation, 7 days ; after infection, 12 days. Onset with great pains in limbs and back. Rigors. **Vomiting.** The pustules may become confluent or remain distinct.

VARICELLA.—With onset, small red spots appear on trunk, face, scalp, becoming vesicles, but these are not cellular or umbilicated ; they crust. Temperature not high.

Very little constitutional disturbance.

SPECIFIC FEVERS.**COMPLICATIONS.**

MEASLES.—Mostly in respiratory system. **Bronchitis.** Acute broncho-pneumonia, which may become chronic. **Laryngitis** may be severe; it precedes the rash. There may be vomiting and diarrhoea. **Delirium.** Rarely cutaneous hæmorrhages. Occasionally **Albuminuria.**

CAUSATION.

Very infectious, especially during the eruptive stage.

VARIOLA.—Mucous surfaces frequently affected, especially conjunctivæ, throat, nose. **Bronchitis; Pneumonia; Pleurisy; diarrhoea; Albuminuria; abscesses.** Cutaneous hæmorrhages and bleeding from the mucous surfaces. **Typhoid State.**

Very infectious. Inoculable by pus of vesicles, also by scabs.

VARICELLA.—None are usual,

Infectious.

ERYSIPELAS.

Describe the part affected, whether much swollen, red, œdematous; whether the margins of the inflamed part are defined or diffused. Note any vesicles or bullæ. Look for enlarged lymphatic glands. Note signs and **Symptoms of Fever.**

Complications.—Cellulitis; abscess; gangrene. **Delirium**; the **Typhoid State**; **Albuminuria**; **Pneumonia**; **Pleurisy**; **Phlebitis**; **Pericarditis**; œdema of larynx; inflammation of fauces. **Diarrhœa**; relapses of the disease.

DIPHTHERIA.

General condition of the patient; ability to swallow, strength of voice, **Dyspnœa**, position of body. P.= ; T.= ; R.= . Examine **Mouth and Throat** for redness and swelling of the fauces, soft palate, uvula, pharynx; patches of membranous exudation, whitish or greyish, often multiple; membrane may be peeled off, leaving surface of mucous membrane raw, but not excavated. Examine lips, cheeks, gums, glands under the jaw. Note voice or cry. See signs of **Laryngeal Disease.**

Laryngeal symptoms.—Commencing with a short cough, and some difficulty of breathing; breathing noisy, stridulous, with a metallic-sounding cough; weak voice; struggling for breath in paroxysms; chest collapsing; pulse weak; face bluish; extremities cold; sweating. Note if tracheotomy be performed or not.

Complications.—**Pneumonia**; **Pleurisy**; **Albuminuria**; adynamia; false membrane on conjunctiva and skin; **Paralysis.**

ERYSIPELAS.

An acute febrile disease characterized by local diffused inflammation of skin and cellular tissue with bullæ and vesiculation. Idiopathic erysipelas usually attacks the face, commencing about the eye.

Causation.—Epidemic and endemic causes. Exposure to cold, and bad **hygienic conditions**; contagion. It may follow injury or operation. Those once attacked by the disease are liable to recurrence.

DIPHTHERIA.

A febrile contagious disease, characterized by the formation of membranous exudations on the fauces and respiratory mucous membrane, frequently obstructing the larynx, often attacking the mucous membrane of the nose and causing an acrid discharge. It is asthenic in its course, and attended by great debility, frequently proving fatal through **Laryngeal Obstruction** or by pneumonia. The period of incubation is various. It may commence with lassitude, febrile disturbance, sore-throat; or those preliminary symptoms may be absent, laryngeal stridor being the first symptom noticed. Sometimes swelling of the glands under the jaw first attracts attention. There may be membrane in the larynx, and none on the fauces. There is less pain on attempting to swallow than with quinsy. Fever not prominent; rarely runs high. When paralysis follows, it is usually after convalescence.

Causation.—Communicable from the diseased to the healthy by secretion of mouth, vomits, expired air. Bad water; sewer gas, and bad **hygienic conditions**. Most common in children. It may be epidemic or endemic.

PYÆMIA.

Examine the body all over for any wound, local inflammation, or suppuration. A very slight wound may produce the disease, *e.g.*, a thorn under the nail, etc. See **Signs of Fever** and **General Condition of the Nervous System**, prostration, **Coma**, **Typhoid State**.

Causation.—Suppuration connected with diseased bone; whitlow; **Phlebitis**; softening clots; ulceration from tertiary syphilis; **Periostitis**. Occasionally it is secondary to internal suppuration or ulceration, *e.g.*, enteric fever, gastric ulcer, abscess of kidney, etc.

Complications.—Occasionally a cutaneous erythema. **Jaundice**, without signs of obstruction; **Albuminuria**; hæmorrhages in skin or from mucous membranes. Low forms of inflammation; **Pericarditis**; **Pneumonia**; **Pleurisy**; empyema; **Peritonitis**.

PUERPERAL FEVER.

General condition.—General signs of **Fever**. Patient usually assumes the dorsal decubitus. There is much tendency to adynamia and the **Typhoid State**, with sweating and **Delirium**.

Abdomen.—Usually distended and tympanitic; bowels often costive. There may be local tenderness over the uterus or in either iliac fossa.

Genito-urinary system.—Note pain or difficulty on micturition or defæcation. Lochial discharge, its amount, if offensive; any clots or pieces of placenta or membranes passed. **Albuminuria**. **Breasts**, if milk is secreted; condition of the glands, tenderness.

Causation.—Epidemic at periods. Endemic in a house or hospital; due to inoculation by nurse or attendant, *e.g.*, from a case of erysipelas, or another puerperal case. Infection with an acute specific fever, *e.g.*, scarlet fever. Local septic poisoning from metritis, decomposing clots, or portions of placenta. Bad hygienic conditions.

PYÆMIA.

Usually commences by an insidious onset, or with chilliness or rigors, and fever with sweating and great prostration. It is characterized by the formation of multiple abscesses, and arthritis with a tendency to suppuration in or around the joints. The tendency is to death by exhaustion, the patient passing into the typhoid state, or by its complications. It may be mistaken for **Rheumatism** or **Enteric Fever**, or may be confounded with broncho-pneumonia, which often accompanies it. Any source of suppuration may lead to the disease, whether the pus be discharged, as from an open abscess, or retained in deep parts, as from periostitis and acute necrosis.

PUERPERAL FEVER.

An acute febrile disease, probably of septic origin, following shortly after confinement, and incommunicable to those not in the puerperal state. The onset occurs about 2nd or 3rd day after confinement, with chilliness or rigors, and the signs of **Fever**. Usually this is attended with scanty and offensive uterine discharges. The secretion of milk is often suspended, but what is formed does not hurt the infant.

Complications.—Pleurisy. **Empyema**. Pneumonia. Pericarditis. Albuminuria. Pelvic cellulitis or parametritis. Peritonitis. Mammary abscess. **Phlebitis**, or phlegmasia dolens. **Arthritis**.

Varieties of Fever.—It may be specially characterized by **Peritonitis**; by pelvic cellulitis; metritis, with much abdominal pain and tenderness. Simple fever, with alteration of secretions, but no other local manifestations; the fever tending to exhaustion, adynamia, and the **Typhoid State**.

AGUE.

Enquire as to the periodicity of the paroxysms. Describe an attack, giving, if possible, the duration of the stages. Note conditions of health in the inter-paroxysmal period. Paroxysms may occur daily—quotidian; with one-day interval—tertian; with two-days interval—quartan. Note general appearance and condition; whether **Anæmia** or cachexia. Examine optic discs: sometimes hæmorrhages are seen in the retina. Urine. T. = ; R. = ; P. = . Also note condition of **Spleen** and liver. General condition of the **Nervous System**.

Complications and Sequelæ.—Enlargement of spleen and occasionally liver; digestive organs disturbed. **Dysentery**; jaundice; **Anæmia**; melanæmia (pigment granules in blood); retinal hæmorrhages; cachexia; **Neuralgia**; brow-ague.

Causation.—Endemic, in low and ill-drained districts. Symptoms may follow in a few hours after imbibing the poison, or may be delayed.

HOOPING-COUGH.

General condition.—State of nutrition; look for signs of **Rickets**. P. = ; T. = ; R. = ; W. = . Enquire for signs of catarrh preceding the development of hooping; simple cough, with expectoration, running at nose, etc.

Respiration.—Physical examination of lungs; the chest, its shape and movements, signs of collapse. Cough; paroxysms, describe them, their frequency, duration, and mode of subsidence; note the amount of asphyxia and venous congestion.

Complications.—Pulmonary collapse; specially in cases of **Rickets**, which usually do badly. Bronchitis and bronchopneumonia; **Convulsions**; **Diarrhœa**. Epistaxis; blood often ejected from mouth.

AGUE.

Characterized by feverish paroxysms, recurring at regular intervals, the patient being well between the paroxysms.

- Paroxysm.*—1. *Cold stage*: Lassitude, headache, malaise, chilliness, shivering, passing on to rigors, the teeth chattering and limbs trembling; muscular pains; epigastric discomfort; goose-skin; face dusky, pinched, shrunken. Pulse small; respirations quick; temperature rising rapidly.
2. *Hot stage*: Rigors and chilliness disappear, succeeded by a comfortable warmth; face less shrunken. Patient then feels hot; flushes; there may be mental excitement. Skin dry and frequently hot; pulse full and strong; respirations more rapid. Headache. Temperature rises higher. Urine abundant.
3. *Sweating stage*: Feeling of heat diminishes; temperature falls. Skin becomes moist and sweating profuse. Pulse and respiration fall in frequency. Headache passes off. Patient feels easy and sleeps, awaking feeling well. Urine scanty, depositing lithates.

Temperature may rise without a developed paroxysm.

HOOPING-COUGH.

Characterized by paroxysms commencing with a series of expiratory coughs, followed by deep, full inspiration with loud laryngeal spasm. Frequently vomiting and expectoration with paroxysms. Child may be comparatively well in intervals. Asphyxia in paroxysms very great; this may lead to ecchymosis under conjunctiva. Sublingual ulcer often results from stretching the frænum over the lower incisors during paroxysms. Ejection of blood not a bad symptom. Look for any spasmodic signs, such as *Tetanus* or chronic spasmodic conditions of muscles of one extremity; this is almost entirely confined to *Rachitic* children. Enquire as to a source of contagion.

SYPHILIS—Inherited.

General condition.—Unhealthy aspect ; dull earthy complexion ; old and shrivelled appearance. Rash on skin ; erythematous patches with abrupt margins ; coppery tint. Squamous skin lesions about mouth, chin, limbs, soles of feet. Sometimes a scab or a pustular rash with bullæ ; there may be much desquamation. The skin about nates and mouth mostly affected. Nails may be unhealthy and chippy.

Mucous membranes.—Mucous tubercles or condylomata at anus and at angles of the mouth ; diffuse stomatitis ; inflammation of gums and tooth-sacs. Thrush. Discharge from nose, often excoriating the lip ; snuffles. **Laryngitis** ; voice or cry hoarse.

Viscera may be affected : spleen large ; liver.

Bones.—**Periostitis** may be very extensive, causing much deformity of limbs and thickening of the skull. Skull thick ; forehead prominent ; craniotabes. Swelling of ends of long bones just above epiphyses.

Nervous system.—Deafness (nerve disease) and amaurosis more common than with the acquired disease ; palsy of a single nerve less common ; occasionally epilepsy or imbecility.

SYPHILIS—Inherited.

This may lead to deposit of gummata.

Eyes.—May be early the seat of iritis, later of keratitis, which occurs towards adult life and is usually symmetrical. There may also be CHOROIDITIS.

Nose.—Mucous membrane swollen ; this leaves nose sunken and flattened. Occasionally, in severe cases, the skin disease is obvious at birth, but usually child appears perfectly healthy till about six weeks old ; the thrush and the rash, etc., then appear.

Scars left in adult.—Bridge of nose sunken in ; linear scars near angles of mouth and about anus. Interstitial keratitis ; iritis ; choroiditis. Prominent forehead.

Nerve deafness.—Often only slight and temporary ; in some absolute and permanent.

Teeth.—All the incisors may be dwarfed and malformed. The upper central incisors are most reliable, dwarfed, usually narrow and short, with atrophy of the middle lobe, leaving a single broad vertical groove.

SYPHILIS—Constitutional and Acquired.

Stages :—Incubation ; efflorescence ; decline ; relapse ; sequelæ.

General condition.—Tendency to emaciation ; debility ; vague pains. **Anæmia**. Look for scar of primary sore.

Digestion.—Mucous tubercles of lips ; sores, leaving scars, at angles of mouth. **Tongue, Soft Palate**, pharynx ; ulceration on tonsils. Superficial and symmetrical ulcers in first stage ; deep, destroying parts, when tertiary ; destruction of these parts. Ulceration and condylomata of anus. **Liver**, perihepatitis ; gummata.

Respiration.—**Laryngeal Disease** with ulceration and tendency to contraction. Lung disease of chronic character.

Nervous system.—**Disease of Brain or Spinal Cord**. Gummata, forming tumour in brain. Palsy of **Cranial Nerves**, especially nerve iii. and nerve vi. ; disease of auditory nerve. Iritis ; **CHOROIDITIS** ; retinitis. Meningitis ; predisposition to **Ataxy**,

Locomotor system.—Nodes and thickening of bones ; **Periostitis**. Skin : syphilides, psoriasis, serpiginous tubercular patches, ulcers with ragged edges, etc.

Lymphatic glands.—Generally enlarged in neck and groins, without tendency to suppuration.

Special phenomena.—Gummatous masses in viscera and skin, etc. **Condylomata** and mucous patches on mucous membranes, or ulceration with tendency to contracting scars. Disease of testes.

SYPHILIS*—Constitutional and Acquired.

These phenomena may be considered as occurring in the second and third stages.

Second stage.—Follows six weeks to two months after inoculation. Rash on skin, scattered coppery eruption; or it may be scaly, papular, pustular, rather on flexor than dorsal aspect. On mucous membranes symmetrical ulcers, tonsils especially, with abrupt edges; condylomata may form anywhere. Iritis usually symmetrical. Occasionally slight periostitis.

Third stage.—Tendency to unsymmetrical ulceration of skin and mucous membranes, with great tendency to relapse. Scars tend to contraction and pigmentation. Tendency to sloughing.

Bone disease.—Periostitis, nodes, chronic thickening, destruction of nasal and palatal bones.

Gummata may form in any viscus. In liver they may be felt during life; in brain may cause signs of TUMOUR; in skin may lead to extensive sloughing.

Arteries often diseased. This may lead to aortic Aneurism, minute arterial aneurisms in brain, and hæmorrhage, Thrombosis, and gangrene.

* Mr. Hutchinson's Article—Reynolds' "System of Medicine."

GENERAL DISEASES—CLASS II.

Diseases often inherited, frequently arising from some internal changes in the patient's tissues or organs, but often due to causes acting from without.

SIGNS OF DEFECTIVE DEVELOPMENT.

Search for accompanying congenital defects of development.

Defect of heart ; cleft palate ; deformity of hands or feet ; supernumerary fingers and toes ; epicanthic folds in excess ; unusual shapes of the ears, and asymmetry between the two ears ; coarse or ichthyotic skin ; hair on forehead. Abnormal conditions of head ; size and shape ; fontanelle and sutures, whether open. They are sometimes prematurely ossified, and the forehead prow-shaped. *Eye* : Coloboma ; congenital defect of sight. Undescended testicle ; hydrocele ; hernia. *Skin* : Coarse, ichthyotic ; deficient in elasticity ; increased areolar tissue ; extremities blue ; chilblains.

Description.—But little spontaneous movement ; dirty habits ; fits ; paralysis ; inability to hold head up, to talk or walk. In low-class cases repetitive movements are common, *e.g.*, continuous movement of one arm, purposeless and rhythmical ; absence of intelligence ; not attracted by light or sound.

Lungs.—Liability to bronchitis.

Nervous system.—Defective in intelligence ; convulsions ; defective motor power ; insufficient power of co-ordination.

Causation.—Syphilis ; drunkenness in parents ; relationship between the parents. Most common in first member of a family.

ANÆMIA.

Pallor of skin and mucous membranes, lips, and conjunctivæ.

When the fingers are held up to the light the redness of the borders is seen diminished. **Edema** of feet ; possibly puffiness of face.

Circulation.—Examine arteries and veins in the neck ; condition of heart. See condition of the blood and its microscopical characters. Look for **Diseases of Vessels**. Breathlessness.

Nervous system.—**Headache** ; **Neuralgia**, especially spinal ; intercostal neuralgia ; drowsiness ; mental weakness and irritability ; muscular weakness ; pains in back. Examine optic discs.

Menstruation.—Disordered ; usually lessened, or absent.

Look for **Pernicious Anæmia**, leucocythæmia, enlarged glands, cancer, hæmorrhages from mucous membranes or under skin ; heart disease ; chronic lung disease ; **Bright's Disease**. Examine urine. Examine liver and spleen.
T. = ; P. = ; R. = ; W. = .

PERNICIOUS (Progressive) ANÆMIA.

Look for general signs of anæmia, and the ordinary causes. See amount of redness of the fingers held before a strong light. Examine optic discs ; there may be retinal hæmorrhages. Note condition of the joints and general power of the patient ; also state of digestion.

CANCER.

General condition.—**Anæmia** ; **cachexia** ; **Emaciation** ; loss of muscular strength. Temperature not raised.

Disturbed function of parts affected.—Pressure signs from growth of mass, *e.g.*—(1) Glands in transverse fissure of liver obstructing the vena portæ and causing **Ascites**, or the duct, causing jaundice ; (2) Mediastinal tumour ; (3) Pressure on veins, *e.g.*, vena cava or iliac veins ; (4) Intracranial tumour ; (5) Annular stricture of intestine.

ANÆMIA.

irculation.—Over jugular vein, especially on the right side, a thrill may be felt with the fingers, particularly in children ; but this does not necessarily indicate anæmia. A continuous humming sound heard, **Bruit de Diable**, over jugular like wind among trees, varying with the pressure of the stethoscope. Systolic blowing over the carotid or subclavian artery on very slight pressure. Over the pulmonary (2nd left) costal cartilage a systolic bellows, the second sound being often sharp and accentuated. Heart's action quick ; easily excited to palpitation. **Pulse** soft and frequent.

ausation.—Hæmorrhage ; menorrhagia. Sequent to acute disease. Defective hygienic surroundings. Hot rooms. Want of good food regularly taken. Dyspepsia ; chronic gastric disease ; **Alcoholism** ; **Plumbism** ; mental exhaustion ; fright ; **Malaria** ; heart disease ; **Cancer** ; often due to over-long lactation ; general delicacy ; disturbed nights as well as lactation ; re-establishment of menstruation during lactation. Coincident disease ; Bright's disease sequent to pregnancy ; rapid development of phthisis, which was quiescent during pregnancy.

PERNICIOUS (Progressive) ANÆMIA.

rofound increasing anæmia, accompanied by increasing debility and prostration, tending to death in many cases. Hæmorrhages ; spongy gums ; epistaxis ; breathlessness ; palpitation on exertion. Fat of the body not absorbed ; the subconjunctival fat yellowish. Excretion of urea diminished. There may be irregular pyrexia.

CANCER.

ausation.—Hereditary ; declining period of life ; sequent to blows. Organs commonly affected—uterus, mammæ, liver, stomach, peritoneum, other abdominal sites, lungs.

secondary deposits.—In liver, from the rectum, sigmoid flexure, stomach, etc. In lymphatic glands next to the organ affected.

Complications.—Serous effusion ; adjacent inflammations ; thrombus of veins.

RICKETS.

Enquire as to conditions of feeding ; ability to stand or walk ; age at which walking commenced ; previous health, especially as to symptoms and complications of rickets. Examine bones, head, abdomen. **Anæmia.**

Bones.—Ribs beaded ; enlargement of ends of ribs at their junction with the cartilages. Sternum thrust forward by the falling in of the ribs at side of chest ; hypochondriac regions depressed. Spine may be bent backwards, but is capable of being straightened on suspending the body, lifting the child by the arms. Shaft of long bones often bent, especially in tibiæ if child has walked ; epiphysis enlarged, particularly in radius. Skull may remain patent long after the first year ; the head is large, wide, and flat on the vertex. See diagnosis from **Chronic Hydrocephalus**. Head may be small and not ill-shapen. **Craniotabes**, or soft spots with deficiency of bone, can be felt sometimes in the occipital bone.

EMACIATION.

History as to probable causation. Emaciation, whether gradual or sudden, or coincident with other signs of disease. Distribution of the emaciation, especially in children. The emaciation often affects the body and extremities more than the head and neck. Examine all the organs and urine. Note weight of patient, and record it once a week. Specially enquire as to history of phthisis. Look for **Anæmia**, and the signs of any disease supposed to have produced the emaciation. When a muscle is struck, *e.g.*, biceps, note its irritability, longitudinal contractions, and transverse knotting.

Nutrition indicated by the relation of age, height, weight, etc.
W. = . Spare, thin, emaciated, stout, fat, good muscular development, strong, weak.

Growth rapid, moderate, slow.

RICKETS.

Thickening and deformity of bones. The child may be fat or ill-nourished. Much tendency to sweating, especially about the head; throws off the clothes at night; head much rubbed on the pillow, so that hair is worn from occiput. Dentition late; the teeth often devoid of enamel—soon decaying. General tenderness, so that child cries on being moved. Late in walking. *Late signs*: Head large, flat, square; figure too small in the legs.

Complications.—Tendency to catarrh of intestines; diarrhoea; abdomen large and prominent; Spleen and liver large. **Bronchitis**; collapse of base of lungs. If **Whooping-Cough** supervenes, it runs an unfavourable course with bronchitis; **Convulsions** and **Laryngismus**.

Causation.—Ill-feeding during infancy, especially with farinaceous food; intestinal catarrh; bad hygienic conditions; premature birth.

Digestion.—Teeth late in appearing, and often deficient in enamel; abdomen large, prominent, tympanitic, partly owing to weak condition of its muscles.

Nervous system.—Liability to convulsion; laryngismus; tetany, or chronic contraction of muscles of extremities.

EMACIATION.

Causation.—Chronic lung disease; **Phthisis**; caseous bronchial glands. **Cancer**. Chronic stomach disease; **Diarrhoea**; **Vomiting**. Starvation and ill-feeding, especially in infants. Defective hygienic conditions. **Senile Degeneration**. Sequent to acute disease. **Fever**. **Diabetes**. General **Tuberculosis**. Disturbance of the general condition of the **Nervous System**.

In infants often called **Marasmus**. Look for signs of **Congenital Syphilis**; collapse of lungs; **Rickets**. See state of skin; fulness or retraction of abdomen. State of bowels; constipation or chronic diarrhoea. State at birth, if suckled; how fed, and nature of foods. Thrush.

ŒDEMA OR ANASARCA.

Anæmia. Signs of disease of heart or vessels. Look for **Cardiac Dilatation** or degeneration.

Lungs.—Especially emphysema or phthisis.

Urine.—See signs of **Bright's Disease**.

If anasarca be thought to be due to passive congestion, look for the signs of passive congestion, and note if the œdema lessen or increase with such other signs ; *e.g.*, note if œdema lessen with the disappearance of pulmonary œdema, etc. If anasarca be due to Bright's disease, note if it lessen with lessening **Albuminuria**, and increase of the quantity and sp. gr. of the urine.

AMYLOID DEGENERATION.

Pasty, anæmic appearance. **Anasarca.**

Liver.—Large, firm-edged, uniformly enlarged, smooth. Usually no jaundice or ascites.

Spleen.—Large, firm, smooth.

Kidneys.—Urine very albuminous. **Anasarca.**

Intestines.—Diarrhoea.

Causation.—**Syphilis.** Chronic suppuration. **Phthisis**, with suppuration of bronchi. Chronic disease of bone, see **Scrofula**.

ŒDEMA OR ANASARCA.

Causation—

Obstruction at heart.—**Passive (Cardiac) Congestion.** Cardiac valvular disease; failure of the ventricles; fatty heart; dilated right ventricle. Adherent pericardium.

Obstruction at lungs.—**Emphysema.** Chronic bronchitis. Conditions obstructing circulation in one lung, *e.g.*, chronic pleurisy, empyema, collapse of one lung.

Local pressure on veins.—Pressure on vena cava or iliac veins in abdomen from enlarged glands, **Cancer**, **Aneurism**, pregnancy, **Abdominal Tumour**, pelvic effusion; pressure from ascites. Pressure on subclavian vein from thoracic aneurism or mediastinal tumour.

Changes in blood or vessels.—**Bright's Disease.** **Anæmia**; extreme debility from chronic disease, *e.g.*, cancer, **Phthisis**, diarrhœa in children, **Phlebitis**, phlegmasia dolens, varicose veins.

SCROFULA.

General condition.—W. = ; T. = ; R. = ; P. = . Intelligence dull; phlegmatic temperament. Coarse, flabby, ungainly children, outlines of body ill-marked. Features plain, complexion pasty. Liable to ophthalmia; tinea tarsi: otorrhœa, eczema, lichen, lupus, chilblains, blue hands. Forehead and back hairy.

Respiratory system.—Liability to bronchitis and chronic pneumonia passing on to **Phthisis**.

Digestive system.—Abdomen full; liability to diarrhœa. **Teeth**, no distinctive condition, liable to decay, may be devoid of enamel. Upper lip thick. Gums unhealthy. Tonsils large.

Locomotor system.—Bones thick; joint disease frequent. Skin easily inflamed. Glands enlarged and suppurate, specially in neck.

GENERAL MILIARY TUBERCULOSIS.

General condition.—Look for signs of strumous disease in bones, joints, spine; enlarged glands; **Emaciation**; state of skin. P. = ; T. = ; R. = . Look for **Signs of Fever**.

Respiratory system.—Signs of **Consolidation of Lung** or **Phthisis**; enlarged bronchial glands; cough.

Digestive system.—**Vomiting**; state of bowels, see **Ulceration of Bowels**; ability to take food.

Nervous system.—Signs of **Meningitis**; signs of **Brain Disease**. **Ophthalmoscopic examination** may show optic neuritis or tubercles in choroid. Tubercle in choroid indicates tuberculosis, not meningitis; optic neuritis with tubercle indicates probable tubercular meningitis.

The onset may be insidious, with a previous period of emaciation and lassitude, and after a few days or weeks may be followed by the somewhat sudden onset of a special complication, as **Pneumonia**; **Meningitis**. The general symptoms are mostly prostration, **Emaciation**, sweating, cough, moderate fever—this may be absent. Some of the complications are usually present, and frequently there are the signs of old strumous disease. The disease tends to death by exhaustion or by complications. The presence of miliary tubercles in the lungs does not necessarily cause any abnormal physical signs.

ENTERIC FEVER resembles GENERAL MILIARY TUBERCULOSIS.

- | | |
|--|---|
| <p>Diarrhoea from typhoid ulceration of Peyer's patches.</p> <p>I. Evening exacerbations of fever, mostly in 3rd or 4th weeks.</p> <p>III. Profuse sweating, with great debility and prostration.</p> <p>IV. Bronchitis and pneumonia common complications.</p> <p>V. Emaciation from fever.</p> <p>VI. Mental dulness from fever.</p> | <p>Diarrhoea from tubercular disease of intestines.</p> <p>Remittent hectic fever common, with caseous lung or glands, etc.</p> <p>Profuse sweating a part of the natural course of the disease.</p> <p>Chronic pneumonia may set up general tuberculosis.</p> <p>Emaciation from tuberculosis.</p> <p>Commencing Meningitis.</p> |
|--|---|

DIAGNOSTIC DIFFERENCES.

- | | |
|--|---|
| <p>I. Characteristic rash on the abdomen, etc.</p> <p>II. Diarrhoea and abdominal symptoms prominent.</p> <p>III. Spleen often large.</p> <p>IV. Lung symptoms late in appearing.</p> <p>V. Delirium and exhaustion, proportioned to height and duration of fever.</p> <p>VI. Occurs in those previously healthy.</p> <p>VII. Profuse sweating less common.</p> <p>VIII. High fever.</p> <p>IX. History of individual and family healthy.</p> <p>X. Any age.</p> | <p>No exanthem. Skin may be erythematous ; or sudamina.</p> <p>Bowels usually constipated.</p> <p>Spleen usually normal size.</p> <p>Lung symptoms appear early.</p> <p>Definite signs of meningitis.</p> <p>Usually previous lung disease.</p> <p>Sweating usual.</p> <p>Fever not high.</p> <p>Individual or family scrofulous.</p> <p>Usually young.</p> |
|--|---|

DIABETES MELLITUS.

General condition.—**Emaciation**; weakness. Skin harsh and dry. Mental aberration; low spirits. See **Nervous System**.

Digestion.—Appetite greatly increased; intense thirst. Tongue frequently devoid of epithelium, raw and cracked. Bowels costive; sometimes diarrhoea.

Urine.—Quantity usually greatly increased; greenish colour; high sp. gr.; sugar abundant. Micturition frequent.

Causation.—Most common in males and middle-aged adults; frequent in phthisical families. Exposure to cold. **Alcoholism**; excessive use of sugar; violent emotional disturbance; organic **Brain Disease**; over mental work or anxiety. It may be associated with **Gout**.

ADDISON'S DISEASE.

General condition.—Debility, faintness, pigmentation. Anæmia. Frequently tubercular tendency. Shallow, feeble breathing; breathlessness, sighing, gasping, especially on any effort. T. = ; usually subnormal. If not complicated, no emaciation. Feeble heart action, faintness, pulse thready. Death by asthenia, sudden or preceded by incoherence, delirium, convulsions.

Digestion.—Nausea, retching, **Vomiting**, epigastric pain. Examine buccal mucous membrane and that of lips. Hiccough; anorexia.

Nervous system.—Its general condition. Pains and sleeplessness. Loss of nerve-muscular power; extreme depression.

DIABETES MELLITUS.

Characterized by excessive thirst, excessive hunger, emaciation.

Urine saccharine, dense, and greatly increased in quantity, as a constant occurrence. Saccharine urine may be temporary, as after a convulsion or administration of chloroform. Diabetes is the more permanent condition of glycosuria, with constitutional symptoms and a tendency to certain complications; it usually has a fatal tendency. Onset of symptoms may be insidious or sudden, with nervous disturbance. Sugar may be detected in sweat, tears, saliva.

Urine.—The quantity of sugar usually greatest after food. Glucose may temporarily disappear; so also, not uncommonly, shortly before death.

Complications. — **Broncho-pneumonia; Phthisis; Pleurisy.** Serous inflammation of low type. **Head-pain;** sudden **Coma;** cataract; **Albuminuria.** Skin disease, boils, carbuncle, psoriasis, diarrhoea.

ADDISON'S DISEASE.

Characterized by pigmentation of the skin; attacks of syncope and extreme debility; **Anæmia**, often without emaciation. **Vomiting**, nausea, or epigastric pain. Discoloration is a bronzing colour, specially marked in face, hands, neck, groins, axillæ, penis, scrotum; areolæ very dark; buccal mucous membrane stained; conjunctiva always free. Tendency to advance to death by asthenia. Sometimes termination is sudden.

PURPURA.

Skin.—Description: Hæmorrhages into skin may occur in scurvy, typhus, measles, variola, or from injury. Ecchymoses occur on forehead and under conjunctiva from asphyxia as in epilepsy, if stage of tonic spasm is prolonged, and after severe paroxysms of hooping-cough.

It comes out in successive crops in aggregations of spots. Note their size, situation; whether separate or confluent. They do not fade on pressure, do not enlarge; but others may occur near those first produced, and become confluent with them; they soon absorb, undergoing changes like a bruise. Some fade, while others appear. They occur mostly in dependent parts, and are apt to occur in the legs in cardiac or other obstruction.

The spots may be small, "petechiæ," or elongated patches, "vibices," or in irregular patches, "ecchymoses." The colour is violet, purple, or blackish. At first the margins are abrupt, but these soon fade. Rarely the cuticle is raised, forming "blebs." Development of spots favoured by standing.

Causation.—Hepatic disease; rheumatism; syphilis; heart disease; any debilitating conditions; too restricted a diet.

Not uncommon in old age, and accompanying insanity.

Look for hæmorrhages of gastro-intestinal canal. **Albuminuria** and signs of heart disease; albumen may be temporary. See **Heart** and state of **Vessels**. General condition of lassitude, with pain in limbs and joints.

DEVELOPMENTAL DEFECTS.

Head.—Ill-shapen ; narrow, prow-shaped forehead ; hyper-ossification in various parts. It may be too large or too small. Ill-shapen in the anterior or posterior segments. Forehead overhanging.

Face.—Hare-lip ; epicanthic folds in excess.

Mouth.—Cleft palate and uvula.

Eyes.—Coloboma of iris or choroid, *i.e.*, a deficiency or cleft ; mall-shaped eyeball.

Ears.—Asymmetry ; one or both may be more or less rudimentary ; helix partly unrolled, with rudiment of third lobe ; frequently are deformed with ichthyosis.

Skin.—Ichthyosis ; hair in excess on forehead, arms, back in children ; eyelashes too long ; fine or coarse.

Fingers.—Webbed ; supernumerary fingers or two thumbs, etc. ; inspect feet and toes.

Heart.—See **Congenital Defects.**

Special abnormalities.—Patency of abdominal rings ; non-descent of testicle ; long prepuce ; imperforate anus ; nævus.

SENILE DEGENERATION.

General condition. — Nutrition ; atrophy, or fatty growth. Goitre occasionally. W. = . Skin, hair, colour ; abundance. Involution of generative system.

Locomotor system. — Power to move about ; state of joints ; power to walk.

Nervous system. — Look for tremors and **Paralysis Agitans**. Sleep ; pains ; mental power ; memory ; neuralgia.

Eyes. — Arcus senilis ; cataract ; presbyopia ; hearing and special senses.

Vascular system. — State of **Vessels** ; heart-force. Look for varicose veins ; purpura.

Respiratory system. — **Emphysema** ; dyspnœa on exertion (cardiac).

Digestive system. — Teeth, gums, jaw ; feeble digestion ; flatulence, constipation.

Urinary. — Albumen.

SENILE DEGENERATION.

Special degenerative and pathological tendencies. — The degeneration may be simply atrophy, the skin becoming wrinkled, or there may be fatty growth generally under skin. Skin loses elasticity, becomes wrinkled and pigmented ; it loses transparency and brilliancy. Hair grows on chin in old women ; cancer.

Nervous system. — Brain may be perfectly sound with an atrophied body.

Prognosis as to life. — Soundness of the organs. The degree of senility not being greater than the age of patient indicates. Longevity is often inherited. Adaptation of patient's life to state of his body.

ARTHRITIC DISEASES.

ARTHRITIS.

Note pain, tenderness, swelling, heat, redness, effusion in joints, periarthritis. Deposits or enlargement of ends of bones or out-growth therefrom. Position of joints; mobility or ankylosis. P. = ; T. = ; R. = .

Look for signs of **Rheumatism** and its complications; **Gonorrhœal Rheumatism**; **Gout** and its history; **Rheumatoid Arthritis**, especially when the arthritis has a chronic course with much stiffness and but little fever.

Tabulate the joints affected, indicating the condition of each —“effusion,” “swollen and painful,” “tender and red,” etc.

RHEUMATOID ARTHRITIS.

Joints.—Arthritis may be acute or subacute. There may be effusion, or only stiffness and pulpy feeling on manipulation. The hand, when made into fist and squeezed, is tender if finger joints are affected. Enumerate joints affected; it may attack temporo-maxillary articulation, or stiffen cervical spine. Every joint in the body may be ankylosed. Dislocation of affected joints may occur.

Causation.—Debilitating causes, hæmorrhages, mental depression, starvation, dampness, and possibly heredity. It may occur at any age.

ARTHRITIS.**JOINTS.****RIGHT.***Shoulder.*—*Elbow.*—*Wrist.*—*Hand.*—Note separately the metacarpo-phalangeal joints and internodes.*Hip.*—*Ankle.*—*Foot.*—Specially note metacarpo-phalangeal joint of great toe.**LEFT.***Shoulder.*—*Elbow.*—*Wrist.*—*Hand.*—*Hip.*—*Ankle.*—*Foot.*—

Temporo-maxillary and vertebral joints.

RHEUMATOID ARTHRITIS.

Small joints commonly first affected, but large joints may be equally attacked. The attacks last longer, are less severe ; less pyrexia and constitutional disturbance than with **Gout** and acute **Rheumatism**; more thickening left, with deformity of joints. No deposits of urate of soda ; no sweating. More commonly commences in fingers than toes, and not with a sudden short attack of single joints.

Complications and accompaniments.—Any organic disease. **Anæmia ; Neuralgia.**

RHEUMATISM.

History of rheumatism ; heart disease ; chorea in family and in collateral relations. Previous attacks in patient.

Present condition.—General signs of **Fever**. P. = ; T. = ; R. = . Skin moist, sweating, sudamina. Note any erythema.

Joints.—Whether tender or painful on movement ; swollen with effusion, with or without cutaneous redness. Enumerate the joints affected, specially noting whether large or small joints are mostly affected.

Vascular system.—Development of cardiac bruits from valvular disease ; pericarditis, with or without effusion. Always map out area of cardiac dulness. Pulse, regularity, etc.

Respiratory system.—**Pleurisy**, single or double ; extensive effusion common. Pneumonia, usually at base ; it may occur without special acute symptoms.

Nervous system.—Rheumatism may alternate with chorea, one following the other, near or at distant intervals. Occasionally delirium. Sleep.

Urine.—Usually a deposit of pink lithates during fever. Rarely a trace of albumen.

Rheumatism in children.—Symptoms often less severe than in adults ; less pain, but little fever ; skin often dry ; great tendency to heart disease, even when able to walk about ; often thought to be “growing pains ;” duration of fever a very few days, or it may be absent.

Nodules under skin not uncommon, even without arthritis or pain ; most common on prominences of bones or tendons, about elbow, knee, ankle, spine. They are usually accompanied by progressive heart disease.

Minor symptoms.—Liability to swollen joints on over-exertion ; stiff neck ; effusion in sheaths of tendons.

Purpura.—Great liability of serous inflammations without arthritis. Erythema.

RHEUMATISM.

A febrile disease, characterized by pyrexia and arthritis with effusion, the inflammation changing from joint to joint and attended with great pain. Skin moist, often sweating; this may be excessive and produce **miliaria**. Great tendency to serous inflammations attended with great effusion, usually quickly absorbed and not leading to suppuration. Tendency of all these conditions to relapse after convalescence. Subacute attacks often succeed the acute.

In children pain and fever often slight, but still tendency to heart damage very great.

Nodules.—Small masses of fibrous growth from size of a pin's head to an almond, often felt better than seen; painless, usually movable.

Complications.—Inflammatory conditions; endocarditis; **Pericarditis**; **Pleurisy**; **Pneumonia**; **Bronchitis**. Relapses of fever and arthritis. Erythema. Hyperpyrexia; **Delirium**; **Chorea**. **Albuminuria** occasionally.

Tonsillitis is frequent at the onset with fever, or it may precede it by a week or two. At same time there may be stiff neck.

Causation.—Exposure to cold and wet. Inherited tendency. Tendency to recurrence, especially in early years.

Exciting causes.—Exposure, over fatigue. Note recent antecedents; scarlet fever, tonsillitis, pharyngitis.

GONORRHOEAL RHEUMATISM.

Joints.—Wrist and knee affected by preference. Pain and effusion ; much stiffness, often causing a considerable amount of ankylosis. No tendency to suppuration, but infiltration and thickening around joint.

Generative system.—Muco-purulent or gleet discharge from urethra.

GOUT.

Joints.—Enumerate joints affected. Note periarthritic inflammation and infiltration, deposit of concretion, or thickening of bones. Examine bursæ for tophi. Take the history of previous joint affections. See **Arthritis**.

Vascular system.—When gout has lasted many years the vascular system often degenerates ; **Heart** becomes dilated and hypertrophied, especially with **Granular Kidneys**. **Pulse**, force and tension.

Digestive system.—General signs. These functions are often disturbed. Teeth much ground. Liver disease common. Enquire for piles.

Urine.—Often albuminous with signs of chronic Bright's disease. Amount of **Uric Acid** deficient.

GONORRHOEAL RHEUMATISM.

eldom seen in females. The disease runs its course through weeks or months. After slightly affecting many joints it becomes confined to one or two. No great pyrexia ; but little tendency to inflammation of internal organs.

GOUT.

In acute attack usually commences in early morning in one great toe. Severe pain, followed by swelling around the joint ; local œdema ; skin red and glazed, exquisitely tender. Attacks tend to recur at shorter intervals. Tophi or concretions of urate of soda may form around joints, in bursæ, or in the external ears.

Causation.—Most common in males at middle life. Hereditary tendency marked. Habits of intemperance ; exposure to weather. **Plumbism.** Any depressing circumstances or injury may excite an attack.

Complications and accompaniments.—**Chronic Bright's Disease.** Heart changes and **Disease of Vessels.** Skin affections ; psoriasis, eczema. **Diabetes.** Liver disease. Thrombus in veins. Tophi may discharge, forming sinuses.

DISEASES OF THE NERVOUS SYSTEM.

NERVOUS SYSTEM.

General conditions.—INTELLIGENCE ; SPEECH ; SLEEP ; HEAD-PAIN ; VERTIGO ; COMA ; VOMITING ; PARALYSIS ; CONVULSION ; SPASM ; TREMOR ; RHYTHMICAL MUSCULAR MOVEMENTS ; DELIRIUM.

INTELLIGENCE.—Giving good clear answers to questions. Memory : Memory for past events, or those of recent occurrence ; power to perform easy calculations. The face may temporarily or permanently lose the expression of intelligence.

SPEECH.—Stammering. Slow, jerky. Using inarticulate sounds only. Mute. **Aphasia.**

SLEEP.—Easily falling asleep ; sleeping soundly and waking up refreshed in the morning. Wakeful ; disturbed by dreams ; remembering dreams. Insomnia, *i.e.*, loss of sleep. Raving at night. Somnambulism. Tooth grinding.

HEAD-PAIN.—1. Its situation, whether general or local.

2. Its characters—heavy, dull, aching, throbbing, shooting, darting, sense of fulness. Whether constant, intermittent, recurrent, or periodical. Its intensity and variability.

3. Effects of movement and change of position, of light and sounds, etc.

4. Its mode of onset. If previous attacks, note periodicity.

5. If accompanied by soreness or tenderness at particular spots, see **Neuralgia.**

6. Look to state of **Special Senses**, especially **Sight** ; Inquire for dysæsthesia of sight.

7. If accompanied by **Vomiting**, note its relation to pain.

8. Look for signs of **Brain Disease**, **Convulsions**, **Paralysis**, **Hysteria**, **Condition of Sensation**. Examine **Optic Discs**. Look for **Neuralgia**.

9. Examine urine for sugar and albumen.

10. Character of pulse ; temperature.

History of neurosis in individual or family. History of phthisis or strumous affections.

NERVOUS SYSTEM.

General conditions.—Note all departures from the physiological condition. The muscular power should be such as to enable ordinary work to be performed.

INTELLIGENCE may be naturally dull or mental power may be lost from disease, *e.g.*, **senile decay**, dementia, general paralysis, epilepsy. Mental delusions may arise in sane people. Intelligence is proportioned to age, education, and surroundings. Ask as to school-work in children.

SPEECH.—**APHASIA** = loss of faculty to speak words, though he can recognize them when written or spoken.

AMNESIA = loss of faculty for the memory of words, but can repeat them if suggested to him.

SLEEP.—Restless tendency to turn the body may prevent sleep even if drowsiness is present; frequent in **Alcoholism**. **Insomnia** may be caused by heart disease or over mental exertion. Muscular twitching and cramps not uncommon from fatigue. Pain may prevent sleep.

HEAD-PAIN.—The first thing to decide is whether the case be one of organic or functional disease; in the latter case, the attacks, when recurrent, are commonly spoken of as headaches.

HEADACHE may be pericranial, frontal, occipital, or diffused, or bilateral. Headaches may recur periodically; in women frequently at the menstrual period. After an attack there is a certain amount of immunity. Attacks may be excited by over-work, sleeplessness, want of food, errors of diet, constipation, etc. With the attacks disorders of sight are common: sparks, coloured stars, zig-zags with coloured bright margins; hemiopia (seeing only half of any object looked at). Other senses may be disordered. Vomiting frequently terminates the attack. Accompanying the attacks, or alternating with them, may be much mental depression, mental weakness, and perverted ideas of things. Such recurrent headaches are common during pregnancy. Such attacks, accompanied by vomiting and coloured vision, are often spoken of as "bilious attacks."

History.—Look for signs of **Meningitis** and **Brain Disease**.

VERTIGO.

- I. Feeling of giddiness experienced by the patient, objects appearing stationary.
- II. External objects appear to move, *e.g.*, up and down, horizontally, approaching and receding.

Vertigo may be increased or relieved by movement and position.

Test hearing and sight. Examine for diplopia. Look for signs of **Brain Disease**. **Anæmia**. Examine **Vascular System**. Urine.

COMA.

History ; onset ; previous signs of **Brain Disease** ; **Convulsions** ; **Vomiting**.

Causation.—1. Injury to head.

2. Examine urine generally, and for sugar and albumen ; also for alcohol and poisons.

3. Cerebral Hæmorrhage. See signs of **Bright's Disease**. **Vascular Degeneration**.

4. Coma sequent to **Convulsion**.

5. Coma may occur during fevers.

6. **Meningitis** and coarse brain disease.

7. Heart failure. Examine pulse and heart's sounds.

8. **Delirium** frequently ends in coma.

9. **Alcoholism** and poisons.

Circulation.—Note pulse, small and soft in syncope, often hard in uræmia. First sound in heart failure.

Look for signs of Brain Disease. State of **Intelligence** and **Sensation**. Test power to perform certain acts, *e.g.*, protrude tongue, swallow food, move fingers, etc. Condition of sphincters. Note condition of sleep. **Delirium**. Subsultus tendinum. Position of body, *e.g.*, dorsal decubitus. Character of respiration, whether stertorous.

Examine for signs of Brain Disease and Paralysis. Examine urine ; lungs ; heart, its strength and sounds ; pulse ; condition of arteries. Smell breath for alcohol. **Edema**. Temperature. Action of sphincters. Eyes : strabismus ; Pupils. **Ophthalmoscope**. **Reflexes**.

VERTIGO.

May occur during sleep or on waking. It is common at climacteric period with degeneration of **vessels**, **Emphysema**, **Bright's Disease**. Vertigo may be due to diplopia dependent upon weakness of an ocular muscle or to some error of refraction as hypermetropia. Ménière's disease of ear; **Alcoholism**; excessive smoking; mental or physical exhaustion; dyspepsia; **anæmia**; heart disease; exposure to the sun. It may accompany simple recurrent **Headaches**.

COMA.

History. Coma may result from old-standing brain disease. There may be history of chronic disease capable of producing coma.

Causation.—1. Injury may produce compression of the brain. Collapse; shock; syncope.

2. See **Uræmia**. In **Diabetes** glycosuria may disappear before coma sets in.

3. Extensive cerebral hæmorrhage may cause deep coma. Hæmorrhage into pons causes universal powerlessness and contracted pupils, resembling opium poisoning.

4. Any severe exhaustion may cause coma.

5. Exposure to great heat, as summer sun.

6. Almost any brain disease may end in coma.

7. **Arterial Disease** may lead to cerebral hæmorrhage; heart disease to **Embolism**.

8. This is a great danger in fevers.

Look for—Coma may be partial or complete, constant or remittent. Signs of motor power may be partially or wholly lost. It may be a sign of the **Typhoid state**, with delirium; then the pulse is usually very soft. Wandering at night in febrile diseases may pass on into **Delirium** and coma. The lungs are usually congested, with pulmonary œdema or hypostatic pneumonia.

Examine for **Alcoholism**; smell breath and test urine. The vomits or washings of the stomach may be smelt and tested for poisons—opium, alcohol, hydrocyanic acid. Urine may be obtained by the catheter. Stomach-pump may be used in poisoning cases. Avoid mistaking brain disease for poisoning.

VOMITING.

Describe vomits; containing undigested food, frothy like yeast; look for *sarcinæ*; watery; smell; containing blood or bile.

See **State of Tongue** and bowels; **Abdominal Pain**; signs of dyspepsia.

Look for reflex causes, *e.g.*, pregnancy, **Ovarian Tumour**, disease of liver, **Gall-stones**; **Renal Calculus**. Examine urine.

Causation.—Stomach disease or derangement. **Œsophageal obstruction**. **Obstruction of Bowels**. Poisons. **Alcoholism**. **Uræmia**. Hepatic disease. Pelvic disease. **Pregnancy**. Ovarian disease. **Addison's Disease**. **Brain Disease** or disturbance. **Migraine**.

DELIRIUM.

Its characters; if attended with illusions and purposeless muscular movements, *e.g.*, subsultus tendinum, picking of bed-clothes, etc. Test consciousness by speaking to patient and requiring an answer to a question, or that he shall protrude his tongue, etc.

Causation.—**Plumbism**, **Alcoholism**, and such causes as may produce **Coma**. **Belladonna**. **Camphor**.

TYPHOID STATE.

Asthenia or **adynamia**. Temperature not high. Tongue tends to dryness, with crusting and formation of sordes on teeth and gums; lips cracked and dry; deglutition difficult. Pulse very soft, compressible, dicrotous, irregular. Heart's action weak; first sound hardly heard. Tendency to pulmonary congestion, œdema, and hypostatic pneumonia.

Drowsiness; **Delirium**; **Coma**; subsultus tendinum; picking bed-clothes. Paralysis of sphincters or retention of urine. Dorsal decubitus complete.

VOMITING.

If of cerebral origin it is—1. Purposeless, not specially after taking food, and not relieving symptoms.

2. Tongue clean ; no special signs of **Digestive Disturbance**.
3. General absence of premonitory symptoms or nausea before vomiting ; contents of stomach ejected easily without retching or much effort.
4. Vomiting frequently arrested by the horizontal position, recurring on becoming erect.
5. Concomitant signs of disturbance of the **Nervous System** or signs of **Brain Disease**.

If vomiting appear to be of cerebral origin use **Ophthalmoscope**. Take temperature ; look for other signs of **Brain Disease**. Intermittent pulse is an early sign of **Meningitis** in children.

DELIRIUM.

May be active ; violent ; low muttering. It usually commences at night with talking and wandering of the mind. When moderate in degree temporary consciousness may be restored by speaking loudly and clearly. Is usual in the course of fevers. It may be due to simple exhaustion ; as from hæmorrhage after labour, etc.

TYPHOID STATE.

A prostrated condition, nervous symptoms, heart failure. An unfavourable termination of **Delirium**, **Coma**, delirium tremens, and acute febrile diseases. Note at each observation strength of heart sounds, force of pulse, and the manifestation of any further nervous symptoms. Dorsal decubitus is usually complete, *i.e.*, the patient lies flat in the trough of the bed ; muscular power is prostrated. If prolonged, bed-sore may form. **Albuminuria** and hypostatic pneumonia frequently coincident.

PARALYSIS.

See **Hemiplegia; Palsy of Cranial Nerves; Minor Paralysis.**
Test **Motor Power.**

See signs of **Brain Disease; signs of Disease of Spinal Cord.**
View the part paralysed, and examine as to **Motor Power.**
Note the parts paralysed and the muscles affected, stating whether the fine and general movements of the limb are wholly lost. Note state of nutrition of the part, contractions, rigidity, etc. Test reflex action by tickling, pricking, etc. See **Sensation. Electric Tests. Examine Optic Discs.** Look for signs of **Syphilis. Vascular Degeneration.**

ELECTRIC TESTS.

If one muscle contract to a lesser force of the current than another, it is said to be more irritable. To ascertain the irritability of a muscle reduce the strength of the current to the lowest point at which it will produce action. A full power of current simply shows the strength of the muscle. If in hemiplegia there be a well-marked difference in the reaction of the two sides the paralysis is not feigned. Diminished contractility may be due to disease of brain, cord, motor nerve, morbid condition of the muscle.

The faradaic current may be applied over the muscle to be tested, or the galvanic current to the nerve supplying it.

Loss of electric contractility is a sure sign of disease. Faradization is sometimes useful to prove the presence of muscle in a fat limb in which it is suspected that tissue is wasted.

Functional aphonia may be from hysteria or exhaustion. Sometimes the fauces, palate, and pharynx are anæsthetic. Laryngoscope shows healthy, motionless, white true cords, and a larynx otherwise healthy.

History —Often sets in suddenly. Liable to relapses.

Look for **Hysteria, Phthisis.** Signs of mediastinal pressure.

PARALYSIS.

Paralysis may depend upon disease of nerves or nerve-centres, or may be only **Functional Paralysis**. When a muscle is paralysed, it usually atrophies in a short time, and on regaining strength regains its nutrition. In **pseudo-hypertrophic Paralysis**, the flexor muscles of the lower extremities become weak, but greatly enlarged.

In paraplegia, see **Spinal Cord Disease**.

Paralysed muscles often become rigid, *e.g.*, hemiplegia, **Infantile Paralysis**. General muscular weakness, not dependent on simple debility and not secondary to disease of viscera, is seen in **General Paralysis** and **Diphtheritic Paralysis**. See **Minor Paralyses**.

FUNCTIONAL.

Age and sex.—Most frequent at onset of puberty and climacteric period; almost confined to females.

Hysteria.—Present more or less. No signs of organic disease.

Atrophy of palsied part.—Palsied part well nourished. No bed-sore.

Sensation.—May be lost, hyperæsthetic, or perverted.

Reflex action.—Not obliterated.

Electric tests.—Reaction readily obtained.

Palsy of Cranial Nerves.—Not seen.

Aphonia.—Common; may be the only palsy.

Part paralysed.—Frequent change. Often partial of one limb or part. Sphincters not paralysed. Urine often retained.

ORGANIC.

Most common in degeneration; sexes more equally affected.

No signs of **Hysteria** or **Epilepsy**. Disease of heart, kidneys, etc.

Atrophy follows paralysis. Sacral bed-sore frequent.

If lost temporarily, usually returns before motor power.

In very many cases lost.

Lost in disease of cord.

Common; specially of face and tongue.

Rare from organic nerve disease. See **Aphasia** in right hemiplegia.

No changes without fresh lesion. May be permanently rigid, with coldness and tendency to slight œdema.

TENDON REFLEXES.*

Name of Reflex.	Mode of Excitation.	Nature of Result.	Level of Cord upon which Reflex depends.
Knee-jerk . . .	By striking patellar tendon with edge of hand or with percussion hammer, whilst leg hangs loosely over fellow, or over forearm of operator; also by striking quadriceps tendon, above patella.	A single upward jerk of the leg and foot, slight or distinct. This reflex occurs in health.	2nd and 3rd lumbar nerves.
Ankle clonus . .	With knee extended or very slightly flexed, by pressing quickly and firmly against anterior part of foot (so as to stretch calf-muscles) and then keeping up the pressure. Ankle clonus is not obtainable in a state of health.	A series of clonic contractions at the ankle-joint, continuing as long as the pressure is maintained, and instantly ceasing when it is relaxed. If the condition is very highly marked it may spread to the whole limb, or even to that of the opposite side.	1st to 3rd sacral nerves (lower part of lumbar enlargement).

Name of Reflex.	Mode of Excitation.	Nature of Result.	Level of Cord upon which Reflex depends.
Plantar reflex . .	Tickling sole of foot.	Movements of toes ; of these and foot ; or of these and leg.	1st, 2nd, and 3rd sacral nerves (lower part of lumbarenlargement).
Gluteal reflex . .	Irritation of skin of buttock.	Contraction of glutæi.	4th and 5th lumbar nerves.
Cremasteric reflex .	Irritation of skin of upper and inner part of thigh.	Drawing up of testicle.	1st and 2nd lumbar nerves.
Abdominal reflex .	Irritation of skin of abdomen along edge of ribs, and above Poupart's ligament.		
Epigastric reflex .	Stroking side of chest over 6th and 5th intercostal spaces.	A dimpling of corresponding side of epigastric region (contraction of highest fibres of rectus abdominis).	4th to 6th or 7th dorsal nerves.
Scapular reflex . .	Irritation of skin in interscapular region.	Contraction of posterior axillary fold (teres), or of several of scapular muscles.	6th or 7th cervical to 2nd or 3rd dorsal nerves.

* From a Table published by Dr. Gowers.

CONVULSION.

Paroxysm.—Note the order, progress, and kind of spasm; whether mostly **Tonic** or **Clonic Spasm**. Commencement, whether general or local; commencing on one side, *e.g.*, one hand or finger. Note suddenness of onset, whether attended with asphyxia and marked cyanosis; its duration. Face pale or flushed; fulness of veins; whether distortion of face; head retracted. Eyes: their position; strabismus; state of **Pupils**. Condition of consciousness.

Premonitory symptoms.—**Aura Epileptica**; muscular twitches; dilatation of pupils.

Causation.—**Brain Disease**; **Rickets**; **Syphilis**; **Bright's Disease**; **Epilepsy**; **Hysteria**. Acute diseases—(1) Cerebral; (2) Febrile; (3) Exanthemata; (4) Pulmonary. Reflex exciting causes, *e.g.*, indigestion, worms, teething, ear disease. Examine heart, urine, temperature.

Sequelæ.—**Paralysis**; amaurosis; strabismus; defect of speech; mental disturbance; mania; drowsiness; sleep; **Coma**.

SPASM.

1. **Tonic Spasm** = continuous muscular contraction during a longer or shorter interval.
2. **Clonic Spasm** = alternate contraction and relaxation of muscles.

Facial Spasm is usually one-sided only. The successive clonic spasms are of equal extent and severity, so that successive grimaces resemble one another. In many cases it is chronic in duration and unaccompanied by other spasms. In these particulars it differs from **Chorea**.

Writers' Cramp.—On attempting to write, the muscles ordinarily used in the act are thrown into a state of tonic spasm; this subsides on discontinuing the act of writing. Other dissimilar acts may be performed without spasm.

CONVULSION.

aroxysm.—Usually commences with tonic spasm and pallor or cyanosis, followed by clonic spasms. One side or one limb may be primarily or chiefly affected; then, occasionally, the eyes and head turn to that side, and there may be a few one-sided jerks of the head. **Pupils** usually dilated.

remontitory symptoms.—In children, frequently, fist is clenched, with thumb turned in. **Laryngismus** may precede convulsion.

ausation.—In children convulsions are very easily produced by slight causes. Ill-feeding, teething, worms, and **Rickets** very common predisposing causes. Pyrexia may be due to an acute disease or to continued tonic spasm. Urine may be albuminous from **Bright's Disease**, or may contain albumen or sugar consequent upon the convulsion.

sequelæ.—Convulsions may be symptomatic of brain disease, which may subsequently advance.

SPASM.

Tonic Spasm is frequently attended by pain, and may be preceded by hyperæsthesia. It is seen in the first stage of an **Epileptic** convulsion; trismus, or lock-jaw; tetanus; spasmodic talipes; spasmodic torticollis.

Clonic Spasm may be increased by effort or mental excitement, and may subside during sleep and under chloroform, *e.g.*, epilepsy. It causes movement or displacement of the limb or part affected. It is seen in chorea and muscular tic.

Causation.—Look for signs of **Hysteria**. Reflex exciting causes, *e.g.*, pregnancy, intestinal worms, teething. Dyscrasie, **Uremia**, fevers, spinal irritation, and meningitis. **Hydrophobia**. **Hysteria**. **Brain Disease**.

LARYNGISMUS.

Look to **Nervous System**. **Convulsion**. General convulsions often follow. It may occur in hysterical women, but is most common in infants. Paroxysms may be brought on by excitement or fatigue. Look for **Rickets**, teething, constipation. There may be tonic contraction of muscles of limbs.

TREMOR.

- I. Tremor absent when at rest, but of various intensity when executing a more or less co-ordinated movement, *e.g.*, raising a glass of water, picking up a pin.
- II. Tremor continuous and permanent. Purposive movements exaggerate it, but it does not disappear on repose.

Note the sets of muscles affected ; whether head is moved ; whether muscles supplied by **Cranial Motor Nerves** are affected. Test **Patellar Tendon Reflex**. Take sample of patient's writing.

RHYTHMICAL MUSCULAR MOVEMENTS.

Athetosis = gliding movements, frequently repeated in the same order. Generally accompanies epilepsy, and usually *hemiplegic* in situation.

LARYNGISMUS.

Characterized by paroxysmal convulsion of the laryngeal muscles and noisy inspiration ; no specific catarrh or special lung trouble, as in **Whooping-cough**. Muscles of chest and abdomen may be involved. Most common in young boys, and on waking from sleep. It may become almost continuous crowing, the veins being distended and face distressed. Child rarely dies in an attack.

TREMOR.

In **Paralysis Agitans**, tremor continues when at rest.

In **Sclerosis**, tremor is increased by movement, ceasing during repose ; so also in mercurial tremor.

In paralysis agitans, the face, head, and cranial nerves usually escape.

Sleep arrests tremor temporarily. Tremor may be general, affecting the head, or not ; it may be localized to a limb. Tremor is a simple vibratory repetition of purposeless movements, not displacing a limb greatly. Fine movements are those through small arcs.

Muscular tremor is a characteristic symptom in paralysis agitans. Disseminated sclerosis ; **Alcoholism** ; mercurial tremor ; **General Paralysis of the Insane**.

RHYTHMICAL MUSCULAR MOVEMENTS.

Chorea may be a congenital or an acquired disease ; it may be hemiplegic or both-sided.

MOTOR POWER.

Ability to stand, walk, walk up stairs, work, etc. State some act the patient can or cannot perform ; how far he can walk. Power over large joints, small joints, finer movements of fingers, *e.g.*, writing.

Movements of upper and lower extremities.—Test power of simple movement, and power to overcome resistance. Test movements of larger joints and muscles ; and power over individual digits.

Movements of head and trunk.—Patient lying on his back, let him erect trunk without use of hands. Examine spine.

Respiratory movements.—Note respiratory rhythm ; movements, whether principally thoracic or diaphragmatic.

Co-ordination of the limbs.—Gait in walking ; walking well and firmly with head erect ; also walking straight with eyes shut ; walking stiff, one joint being kept immobile from pain ; hip movements much restrained in **Sciatica**. Circumducting one leg, swinging it round, not moving it forward as the other, seen in **Hemiplegia**. Staggering, moving trunk over place where the legs are. Lifting legs inordinately high, then bringing them suddenly down. Walk with eyes shut. Test for **Muscular Anæsthesia**.

MOTOR POWER

May be lessened from general weakness or be lost in one or two extremities only, or in a certain group of muscles. See **Paralysis**.

Movements of upper and lower extremities.—Palsy of upper extremity, if of cerebral origin, is usually accompanied by weakening of lower extremity. Let patient move limbs to order; lift weights; pick up a pin, etc.

Movements of head and trunk.—Motor power over spine may be lost from caries of spine. **Pseudo-hypertrophic Paralysis**. View spine; feel for curvatures.

Respiratory movements.—Cheyne's respiration = a series of respirations hurried and deep up to a certain point, then subsiding to a dead pause.

Co-ordination of the limbs.—If defective, examine joints. **Sciatica**. **Spasms**. **Tremors**. **Paraplegia**. **Chorea**. In **Paralysis Agitans** there is a tendency to propulsion or retropulsion. In **General Paralysis**, stumbling and staggering, or tottering. In **Ataxy**, muscular power in the legs is not lost; the patient may walk, feeling the ground with a stick. In **Hemiplegia** the patient in walking swings round the leg, and then keeping it stiff balances the trunk upon it.

SENSATION.

Objective sensibility (ascertained by examination).—Tactile sensibility of skin. Examine separately the flexor and extensor surfaces, face, trunk. Test the least distance at which two points can be distinguished in various regions. Sensibility to heat and cold. Apply to various parts two test tubes, one containing hot water, the other cold. Or apply a hot and cold sponge alternately.

Subjective sensibility (sensations experienced by patient).—Localized pain in the area of a certain cutaneous nerve, constant or periodical, suggests enquiry as to **Neuralgia**. Sensibility may be lessened, anæsthesia; exalted, hyperæsthesia. Sensation may be perverted, the patient experiencing altogether abnormal sensations, dysæsthesia, *e.g.*, numbness, "pins and needles," a sense of burning, heat and cold. If subjective sensations are complained of, examine for an objective cause, *e.g.*, local tenderness, local inflammation or disease, periostitis. Reflex causes, gastric, uterine, etc. See **Head-pain, Vertigo, Hysteria, Neuralgia, Muscular Anæsthesia**.

MUSCULAR ANÆSTHESIA.

Let patient carry his hand to his mouth, and repeat the act with his eyes shut; let him state the position of his limbs with his eyes shut; let him distinguish between different weights. In all such attempts he fails. Test **reflex action**, and **electric excitability** (usually diminished). Note what muscles are affected; state of muscular nutrition; presence or absence of pain. Test cutaneous sensibility.

SENSATION.

næsthesia, loss or diminution of sensibility ; *hyperæsthesia*, exaltation of sensibility. Both these conditions frequently met with in **Hysteria**.

Emianæsthesia is usually functional ; it may paralyse the special senses of side affected ; it is frequent in *hysteria*. **Analgesia** is the loss of sensibility to pricking, pinching, etc. It may be temporarily removed or transferred to the other side of the body.

Subjective sensibility may be *anæsthetic*, *hyperæsthetic*, or *dysæsthetic*, *i.e.*, sensibility may be lessened, exalted, or perverted. The brain centres of the organs of special sense may be altered in any of these ways ; so also the sense of touch. As sensations of physical life we may speak of "organic sensations," or those due to the changes occurring in the organs of digestion, circulation, respiration, etc. ; the "appetites," a group of uneasy feelings produced by the recurring wants or necessities of the physical system, as sleep, exercise, repose, thirst, hunger, etc. Special *dysæsthesiæ* are the epileptic *aura*, the lightning pains of *ataxy*, the sensation of girthing frequent in spinal cord disease.

MUSCULAR ANÆSTHESIA.*

'A loss of the feeling of muscular action, attended by irregularity, sluggishness, and diminished force of voluntary movement ; but unattended by any necessary loss of cutaneous sensibility or by distinct paralysis.'

A condition frequently seen in **Hysteria**. Usually there is no pain in the limbs, but pain is common in **Locomotor Ataxy**. It may be local. It often precedes *paraplegia*. Usually impaired or lost in **General Paralysis**. Some muscular *anæsthesia* may accompany attacks of *migraine*. See **Headache**.

* Dr. Reynolds' "System of Medicine."

SPECIAL SENSES.

Sight.—Test acuteness of vision with test-type. Examine for perception of colour. To completely examine the sense of sight, further test power of accommodation, refraction, action of ocular muscles separately and in the combined movements of the eyes. Examine the field of vision. See **Pupils**. **Ophthalmoscopic Appearances**.

Hearing.—Test hearing with a watch held at the greatest distance at which it can be heard from each ear. If watch cannot be heard thus, test auditory power of the nerve for sounds conducted through the skull, *i.e.*, place watch on forehead or between teeth. Look for otorrhœa; examine throat; use ear speculum.

Taste.—For acids, bitters, sapid substances; determine each separately at anterior and posterior portions on either side.

Smell.—For pungent substances, *e.g.*, ammonia; aromatic substances, *e.g.*, oil of cinnamon.

CRANIAL NERVES.

Observe movements of eyes, tongue, face, lips, palate, muscles of mastication and deglutition. **Pupils**.

Test **Special Senses**; sensibility of head and face.

Nerve I.—Olfactory, See **Smell**.

Nerve II.—Optic, see **Sight**, **Pupils**, **Ophthalmoscopic appearances**.

Nerve III.—(Palsy).—Ptosis or drooping of the upper eyelid; permanent external strabismus; dilated pupil; loss of accommodation for near objects.

NYSTAGMUS = purposeless vibratory movements of the eyes; usually the movements are in the horizontal plane.

Nerve IV.—Superior oblique muscle. Palsy produces no appreciable deviation of the axis of the eye, but diplopia results and the diagnosis generally depends upon the relative position of the two images.

Nerve V.—Motor to temporals, masseters, and pterygoid muscles. Examine condition of its separate branches. See **Neuralgia**, **Trigeminal**. Examine power of **Taste**. Look for tooth grinding.

SPECIAL SENSES.

Sight.—Defects may occur from errors of accommodation, myopia, hypermetropia, or astigmatism, from changes in the optic nerve or other parts. Illusions may represent an aura preceding an epileptic fit; common in delirium and insanity, not uncommon with recurrent **Headache**.

Hearing.—Deafness may result from obstruction of the Eustachian tube from pharyngeal catarrh, or tonsil disease; wax in ear; disease of tympanum. The nerve may be paralysed from disease, *e.g.*, **Syphilis**; rarely from cerebral tumour. Tinnitus common with and without ear disease.

Taste.—Taste may be lost on one side only. It is impaired in some cases of palsy of Nerve VII.

Smell.—Test either nostril separately.

CRANIAL NERVES

are some sensory, others nerves of special sense, while others are purely motor. The condition of the parts that they supply, as found on examination, often throws much light on the condition of the brain. Paralysis of an ocular muscle or the tongue would indicate intra-cranial disease.

Nerve III.—Paralysis often partial, *e.g.*, ptosis only. Accommodative power alone may be lost, *e.g.*, in **Diphtheritic Paralysis**. This nerve is frequently paralysed from **Syphilis**.

NYSTAGMUS.—A chronic condition, usually congenital, and dependent upon deeply-seated brain lesion.

Nerve V.—Sensory branches give sensibility to the lateral and anterior parts of the head and the eyeball, and common sensibility with taste to the anterior two-thirds of the tongue. It is the afferent nerve in reflex winking on touching the eyeball; if palsied, the eyeball becomes insensitive and the cornea ulcerates and sloughs. See **Neuralgia, Trigeminal**.

CRANIAL NERVES.

Nerve VI.—External rectus of the eye.

Nerve VII.—Examine movements of face in natural expression, in forced voluntary movements, *e.g.*, to grin and show teeth, to frown, to elevate the forehead, to whistle. See respiratory movements of *alæ nasi*. Facial movements, are they symmetrical; compare the two sides of the face. See position of the angles of the mouth, and slope and curve of the upper and lower lips. The depth of the nasolabial groove. *Orbicularis oris*, its power of holding air in the mouth with the cheeks blown out. *Orbicularis oculi*, its action in closing the eyelids, in producing similar folds of the eyelids on the two sides; a similar width of palpebral fissure on the two sides; a firm application of the lower eyelid to the globe, with the punctum applied to the conjunctiva. Note action of *Occipito-frontalis* and *Corrugator*. Test reflex actions of the eyes. Note pronunciation. Examine with care the movement of the soft palate and tongue. Test **Hearing, Sight, Smell, Taste**. Look for dryness of mouth from want of saliva.

Nerve VIII.—**Pneumogastric; Glosso-pharyngeal; Spinal Accessory; Pneumogastric.** Not purely a cerebral nerve; partly spinal, and receiving branches from the sympathetic.

Motor branches.—To larynx, pharynx, œsophagus. Pharyngeal, concerned in reflex act of deglutition.

Superior laryngeal.—Mostly sensory, but motor to arytenoid and crico-thyroid. Its stimulation inhibits inspiration, *e.g.*, when opening of larynx is irritated.

CRANIAL NERVES.

Nerve VI.—It is opposed by Nerve III.

Nerve VII.—Motor to muscles of face, these muscles being used in expression, respiration, eating; certain reflex actions, *e.g.*, eyelids, mouth.

Intra-cranial branches.—*Great petrosal* through Michel's ganglion to levator palati and azygos uvulæ. *Small petrosal* through otic ganglion to tensor palati and tensor tympani and parotid gland. *Tympanic branches* to stapedius and laxator tympani. *Chorda tympani* to submaxillary gland and lingualis.

Bell's Paralysis of the Face differs from the facial paralysis produced by brain disease in being more complete and general in distribution; in the latter the muscles about the angles of the mouth are mostly affected as seen in grinning. Bell's paralysis affects all the muscles on the side of the face; the eyelids, however, retain a little power. The creases of the face are obliterated, as seen on the forehead and in the naso-labial groove; the eye remains more or less permanently open, and the tears overflow. The patient cannot distend the mouth with air, and food accumulates in the cheeks.

Causes.—Cold, disease of ear, syphilitic disease of temporal bone, pressure of glands on facial nerve.

Nerve VIII.—

Vagus nerve.—Is concerned in certain reflex actions, *e.g.*, deglutition, reflex movements of glottis.

Laryngeal branches.—Palsied in **Diphtheritic Palsy**, in **Bulbar Paralysis**, and much dulled in the **Typhoid State**. Concerned in reflex throat cough.

Superior laryngeal.—Afferent nerve in reflex movements, closing larynx in deglutition or when irritated.

CRANIAL NERVES

Nerve VIII.—Continued.

Recurrent laryngeal.—Chiefly motor ; supplies all the muscles of the larynx except the crico-thyroid.

Cardiac branches.—Inhibitory ; pulse may be irregular from brain disease, and small from mental depression.

Pulmonary branches.—Afferent fibres convey the feeling of the necessity to breathe. Motor fibres supply the bronchi.

Gastric branches.—Regulate the peristaltic movements, and the secretion of gastric juice.

Abdominal branches.—Supply liver and are connected with the renal plexus.

Glosso-pharyngeal nerve.—Gives common and gustatory sensibility to the tongue, supplying circumvallate papillæ at back of tongue.

Spinal accessory nerve.—A motor nerve closely associated with the pneumogastric and giving it motor fibres, some of which go to larynx.

Nerve IX.—Principally motor to the tongue and depressors of the larynx and lower jaw.

CRANIAL NERVES.

Nerve VIII.—Continued.

Recurrent laryngeal.—Left winds round arch of aorta, right round innominate artery. When paralysed glottis is passively narrowed on inspiration, and passively dilated on expiration. It may be paralysed by thoracic **Aneurism** or mediastinal tumour, and thus lead to palsy of corresponding vocal cord.

Pulmonary branches.—Concerned in spasmodic **Asthma**, **Whooping-cough**, **Laryngismus Stridulus**. When paralysed leads to congestion of the lungs, *e.g.*, in **Typhoid State**.

Vagustic branches.—Afferent in cerebral **Vomiting**. Dyspepsia may result from brain disturbance.

Abdominal branches.—Mental shock may excite **Diabetes**. Anxiety causes flow of pale urine of low sp. gr.

Glossopharyngeal nerve.—It is concerned in reflex deglutition

Vagus accessory nerve.—Motor to sterno-mastoid and trapezius; fibres pass to the larynx and control the voice, not respiratory movements.

Nerve IX.—Concerned in articulation, mastication, and the commencing act of deglutition. Each function may be separately lost.

BRAIN DISEASE, SIGNS OF.

Head-pain ; Vertigo ; Cerebral Vomiting ; Convulsion ; Paralysis ; Hemiplegia ; palsy of Cranial Nerves ; strabismus ; palsy of Special Senses. Mental or intellectual disturbance ; Coma ; Aphasia. Changes in Optic Nerve. Pulse, intermittence of. Pupils. See general condition of the Nervous System ; Sensation ; Hysteria.

Examination.—Look for history of neuroses ; previous signs of **Brain Disease**. Indications of acute diseases, *e.g.*, take temperature and look for other signs of **Fever**. Examine **vascular system** and **urine**.

OPHTHALMOSCOPIC APPEARANCES.

Test sight and examine **Pupils** previous to using atropine. Some of the principal conditions of the fundus that may be observed are—**Optic Neuritis ; Optic Atrophy**, (1) primary, (2) secondary to neuritis or consecutive atrophy ; over-fulness of veins ; emptiness of arteries ; **Hæmorrhages ; Choroiditis ; Tubercle of Choroid ; retinitis albuminurica**.

OPTIC NEURITIS.—Disc blurred, outline indistinct ; vessels on disc in parts covered with effusion ; veins large. Vision may be perfect. Neuritis is very indicative of coarse intra-cranial disease, *e.g.*, **Tumour**. This condition may subside, leaving but little change noticeable, or it may leave consecutive atrophy. See signs of **Brain Disease**.

OPTIC ATROPHY.—May be sequent to neuritis. It differs in appearance from primary atrophy in having more disturbance of the choroidal pigment around the disc, a less sharply-defined margin, and sometimes thickening of the sheaths of the vessels remains ; it looks dull. Primary optic atrophy gives a more clearly-defined margin ; it is clean cut, and its general appearance brighter. Vessels atrophied or obliterated.

BRAIN DISEASE, SIGNS OF.

a condition of the brain may be judged of by observation of the optic discs and retinæ, as expansions of nerve matter in connection with the circulation of the brain. Also by the condition of parts supplied by nerves having their centres in the brain. Special signs are found in conditions of the muscles, paralysis, spasm, convulsion, want of co-ordination, etc. See **Motor Power**.

Examination. — Onset of acute febrile disease may cause cerebral symptoms. Cerebral symptoms with pyrexia contra-indicate a purely functional disturbance.

OPHTHALMOSCOPIC APPEARANCES.

HEMORRHAGES in the fundus are usually situated in the retinæ. They are common in **Pernicious Anæmia**; in **Retinitis Albuminurica**—here they are accompanied by white shining spots. They may be seen in *ague* and *leucocythæmia*. Hæmorrhages, even if considerable, may be quickly absorbed, and may recur.

OROIDITIS.—Dull yellowish patches over fundus; there may be subsequent atrophy, the shining sclerotic showing through. Around the patches the choroidal pigment is much disturbed, forming black rings or patches. It may be disseminated or marginal. It is often **syphilitic**.

BERCLE OF CHOROID.—Small circular spots, more or less circumscribed, reddish or greyish-white in colour. They may be elevated above the level of the choroid with retinal vessels passing over them; adjacent choroid may be normal. Their growth in size may be watched. See **General Tuberculosis**.

BERCLE IN CHOROID is rather a sign of general tuberculosis than a sign of meningitis; but coincident optic neuritis indicates probable tubercular meningitis. Tubercle in choroid in a case of continued fever suggests tuberculosis as its cause.

PUPILS.*

Let a full light fall upon the face. Keep one eye covered and test the other; letting light suddenly fall upon it, observe its reaction. Partially screening one eye, let light fall suddenly upon the other, and observe the reflex effect upon the first eye. This reaction involves the optic nerve on side exposed to light, corpora quadrigemina, and Nerve III. on the side shaded. Note contraction of pupil on near accommodation.

Observe.—1. Its shape, regularity, and outline; adhesions may cause irregularity; shape when dilated.

2. Size; may be measured by reference to the holes of catheter gauge.
3. Activity to light and on near accommodation.
4. Any differences between the two pupils.
5. Colour of iris, distinctness of muscular bundles.

Mydriasis = great dilatation of pupil. 1. Artificial, by atropine.

2. Paralytic, from palsy of Nerve III. 3. Spasmodic.

Myosis = contraction of pupil.

* See Mr. Hutchinson's article on "States of the Pupil." "Brain," Vol. I. II.

PUPILS.

large in **Anæmia** and debility; dilated during rigor and **Convulsion**. May be exceedingly mobile in debility. Sluggish pupils indicate defect of vaso-motor nerve, and then the pupil is rather small. A pupil sluggish to the direct action of light may respond immediately when the other eye is acted on by light, thus—(1) Irido-motor apparatus is sound; (2) Peripheral structures of the second eye are sound; (3) There is a defect in the percipient structures of the first eye. Pupils may remain active with optic atrophy. The movements upon accommodation (Nerve III.) may be good though reaction to light (vaso-motor) be lost, *e.g.*, in **Ataxy**. Precise symmetry in size of the eyes is not common.

iridoplegia = palsy of pupil to light, but not to drugs.

ycloplegia = absolute loss of accommodation.

phthalamoplegia interna = both the radiating and circular fibres of iris and the ciliary muscle are paralysed. Pupil is motionless and accommodation lost.

iritis may be a sign of previous **Syphilis**.

SPINAL CORD DISEASE, SIGNS OF.

Paraplegia, partial or complete ; **Spasms ; Tremors**. Dysæsthesia, principally confined to the lower extremities ; **Paralysis of Sphincters** ; sacral bed-sore ; atrophy of optic nerve.

Motor power.—See power of co-ordination of the limbs ; their state of nutrition. Enquire as to the state of sphincters. Test **reflex action** of extremities and patellar tendon reflex.* If there be paralysis, state what groups of muscles are involved, and which escaped ; gait in walking.

Sensation.—Objective sensibility ; examine the muscular sense. See **Muscular Anæsthesia**. Subjective sensibility ; dysæsthesia of lower extremities.

Look for Ophthalmoscopic appearances ; condition of spine. See **Pupils**.

Causation.—Exposure to cold ; over-exertion ; functional paraplegia in hysteria ; heredity ; reflex paraplegia, from urethral stricture, sequent to confinement ; spinal meningitis ; spinal hæmorrhage ; injury to back ; **Syphilis ; Alcoholism**.

* Dr. Gowers : " Med.-Chir. Trans." 1879.

SPINAL CORD DISEASE, SIGNS OF.

Muscles supplied by spinal nerves are alone paralysed. See if signs of **Brain Disease** and **Palsy of Cranial Nerves** are absent. Paraplegia may be purely functional.

Motor power.—If there is paralysis of a special group of muscles, see **Minor Paralyses**. Specially note the power of **Co-ordination of the Limbs**.

Sensation.—Sensation of girthing round abdomen, frequent in spinal cord disease.

"Lightning pains," darting, burning, or pricking; common prodromata of **Ataxy**, often mistaken for rheumatism.

Look for sacral bed-sore, very apt to form in myelitis, probably as the direct effect of the nervous lesion. No bed-sore in **Hysteria**. Test reflexes.

Paralysis.—Reflex paraplegia seldom complete, less widely-spread, and less defined than paraplegia from myelitis. See **Paralysis, Functional or Organic**.

MINOR PARALYSES.

Paralysis of isolated muscles, or groups of muscles. **Spinal (Infantile) Paralysis.** Onset sudden; most common in infancy; frequent in healthy children; occurs but once; large muscles principally affected, *e.g.*, deltoid rather than muscles of fingers.

PROGRESSIVE MUSCULAR ATROPHY.—A chronic disease causing atrophy of certain muscles, with corresponding loss of power, attacking shoulder and ball of thumb by preference, gradually involving more muscles; no pain.

PSEUDO-HYPERTROPHIC PARALYSIS.—Enlargement of muscles paralysed; usually attacks calves, thighs, buttocks, erector spinal muscles; mostly seen in children—male children; several children in same family may be affected.

PARALYSIS OF EXTENSORS OF FOREARM.—Usually due to plumbism.

CROSS PARALYSIS.—Palsy of face on one side, and hemiplegia of the opposite side.

LABIO-GLOSSO-LARYNGEAL PARALYSIS (Bulbar paralysis).—Paralysis of muscles of tongue, palate, pharynx, orbicularis oris; death by asphyxia.

PARALYSIS OF THE FACE.—See *Bell's Paralysis*. Paralysis of muscles of deglutition frequently due to *Diphtheria*.

NEURALGIA.

Symptoms.—Onset, whether sudden or gradual, whether preceded by general or local disturbance; the paroxysms, whether severe, their frequency, the character of the pain. The effect of heat and cold upon the pain. Look for tender points in the course of the nerve affected, and its branches. Examine cutaneous sensibility at the seat of pain.

Causation.—Age, sex, heredity, injury to nerve, frequent movement of the limb, or pressure upon a nerve. *Malaria, Syphilis, Gout, Rheumatism, Alcoholism, Anæmia, Hysteria*, cold, mental anxiety, carious teeth. Reflex causes, *e.g.*, from pregnancy, pain in eyeball from caries of a tooth.

Conditions characterised by neuralgia.—*Locomotor Ataxy*, lower extremities; *Herpes Zoster*, a long area of skin supplied by nerve affected; *Herpes Labialis*.

MINOR PARALYSES.

Paralysis of isolated muscles, or groups of muscles. See **Infantile Paralysis.**

PROGRESSIVE MUSCULAR ATROPHY.—Enquire for injury to nerves ; lead poisoning ; the nature of the employment, as to its using one particular set of muscles. **Electric tests.** Irritability of muscles when struck. Cutaneous sensibility,

PSEUDO-HYPERTROPHIC PARALYSIS.—Test reflex action, and electric tests. See motor power.

PARALYSIS OF EXTENSORS OF FOREARM.—Supinator longus and extensor carpi rad. longior usually escape.

BOSS PARALYSIS.—May be due to disease of pons.

ABIO-GLOSSO-LARYNGEAL PARALYSIS.—Often accompanies hemiplegia and chronic brain disease.

PARALYSIS OF FACE.—May be due to lesion of brain, or **Bell's Paralysis.**

NEURALGIA.

Symptoms.—Pain localized, almost invariably unilateral ; in recent cases paroxysmal or distinctly intermittent. Gradual formation of tender points, where nerve-branches become superficial, passing through bone or fascia, the points of Valleix.* Absence of local causes of pain, such as inflammation, periostitis, new growth. Absence of fever or local heat.

Prevalence.—Most common in females at puberty ; when developing at forty years or older, is very intractable. Malarial neuralgia, usually in supra-orbital nerve. Injury to a nerve may cause neuralgia of branches communicating with it.

Conditions characterized by neuralgia.—The subjects of hysteria and epilepsy are very liable to neuralgia.

* See Anstie on "Neuralgia."

NEURALGIA.

TRIGEMINAL.—Tender points. 1. Supra-orbital. 2. Palpebral, in upper eyelid. 3. Nasal, at junction of nasal bone and cartilage. 4. Ocular, a point in the eyeball. 5. Trochlear, at inner angle of orbit

Superior maxillary division.—1. Intra-orbital. 2. Malar. 3. A point in the line of the upper jaw.

Inferior division.—1. Temporal, a little in front of the ear.
2. Inferior dental (mental), towards front of lower jaw.
3. Lingual, at side of tongue.

SCIATICA.—Is a neuralgia of the sensory fibres of the sciatic plexus. Note gait in walking; the muscular power of the limb; the state of its nutrition. Look for tender points—along the course of the nerve and its branches, *e.g.*, superficial cutaneous branches in gluteal region; down back of thigh, calcanean and malleolar branches; also behind trochanter.

INTERCOSTAL NEURALGIA.—There is pain and tenderness in the course and distribution of the nerve or nerves affected. It is most common in the left infra-mammary nerve. Pain is constant, at times shooting. Painful points.
1. Vertebral. 2. Lateral, along outer margin of trapezius.
3. Sternal.

NEURALGIA.

TRIGEMINAL.—*Causation* : Any cause of neuralgia, especially malaria ; dental or maxillary disease ; cerebral tumour. It mostly occurs in conditions of low nervous depression. Some severe cases are associated with hereditary insanity. With disease of trigeminal nerve there may be profound disturbance in the eyeball, as in cases of herpes in this region. Ulceration of Cornea, iritis, suppuration, and disorganization.

■

SCIATICA.—*Causation* : Rare under twenty years. Cold ; peripheral irritation, *e.g.*, tight boots. May arise from pressure on the sacral plexus, *e.g.*, pelvic tumours, ovarian, hard fæces. Examine hip-joint.

Pain is more constant and less paroxysmal than in other neuralgia ; motor as well as sensory fibres often affected, diminishing muscular strength ; the limb may emaciate and become somewhat anæsthetic. In walking, the foot on side affected is planted carefully, so as to avoid any jar which would increase the pain.

HEMIPLEGIA.

State side affected. Give history of the onset, whether sudden, gradual, with convulsion or loss of consciousness; whether preceded by abnormal sensations; whether first attack.

P.C.—General condition of **Nervous System**. Look for palsy of **Cranial Nerves**. Examine limbs affected as to **Motor Power**, coarse movements, *e.g.*, power to raise limb from the bed, to move large joints, pronate and supinate; to lift weights. As to finer movements, *e.g.*, use of fingers, to pick up a pin, button shirt, point with index and little fingers, etc., to write. Note power of tongue and face. Palsied limbs, their temperature, atrophy, or rigidity, condition of **Sensation**. Look for signs of **Brain Disease**. **Special Senses**. Condition of cranial nerves. **Sight**, examine for limitation of the field of vision. **Facial Palsy** from cerebral disease. Examine **Optic Discs**. Look for bed-sore.

Causation.—Examine heart, and look for signs of **Vascular Degeneration**. Look for signs of **Bright's Disease**. Look for signs of **Syphilis**. **Hysteria**.

Superficial **reflexes** diminished on side of palsy. Rigidity in part paralysed later on. If involuntary movements of *parts* palsied.

HEMIPLEGIA,*

Right hemiplegia commonly associated with **Aphasia**. Hemiplegia from **Embolism** most commonly right-sided. Onset sudden in embolism, and in cases of extensive hæmorrhage. Sometimes premonitory warnings are experienced in the head or limbs.

P.C.—Nerve VII., when affected, is usually partially paralysed, muscles about mouth being most weakened. There may be the following phenomena :—

1. Head turned to side of lesion.
2. Conjugate deviation of the eyes, both being turned to the side of lesion.
3. Muscles of chest and belly weakened on side opposite to lesion.
4. Paralysis of muscles passing from the trunk to the limbs paralysed.
5. The face paralysed on the side of hemiplegia.
6. The tongue protruded to side of hemiplegia.
7. Arm and leg paralysed on the side opposite to the lesion.

Nos. 1 and 2 are very temporary. Those parts suffer most and longest which have the most voluntary uses. Sensibility is usually restored before motor power.

Causation.—**Valvular Disease of the Heart** may lead to embolism, **Atheroma** to cerebral hæmorrhage or thrombus. Bright's disease, being associated often with disease of vessels and hypertrophy of heart, frequently leads to cerebral hæmorrhage. Syphilitic disease of arteries.

* Dr. Hughlings-Jackson : Reynolds' "System of Medicine."

CHOREA.

If there have been previous attacks, say whether one-sided, and state side affected. The manner of commencement. Previous history as to the general condition of the **Nervous System**. History of school-life. **Headaches**.

F. H. of Neuroses, headaches, hysteria, chorea, fits in infancy epilepsy, **Rheumatism**.

P. C.—Note state of nutrition; general condition of the nervous system. Look for signs of **Brain Disease**. Specially note condition of **Intelligence, Speech, Sleep**.

Motor Power.—Whether muscles supplied by cranial and spinal nerves are alike affected. Examine face, tongue, soft palate, movements of eyes, movements of head, respiratory movements, movements of trunk and head.

Examine the extremities in detail, e.g., right upper extremity. Is the shoulder much moved? in which direction principally? by the action of what muscles? The elbow, is it more or less moved than the shoulder; what are the principal movements—flexor, extensor, pronator, or supinator? The hand; movements of wrist, fingers, thumb. Fingers may twitch with extensor-flexor or adductor-abductor movements; some digits may move more than others. Postures of hands when held out, also of trunk and spine.

Complications.—Onset of **Rheumatism, Pericarditis, Endocarditis**. Mental symptoms. Look for **Rheumatic Nodules**.

Examine heart, its sounds, regularity. Look for signs of **Anæmia**. Examine urine for urea and uro-hæmatin.

Causation.—The most distinctly demonstrated lines of causation are in connection with **Rheumatism, Heart Disease**, and sudden mental impressions. Reflex causes, *e.g.*, intestinal worms, pregnancy. Enquire for arthritis with enlargement, attended with feverishness or not; over-work, or complaint of school lessons.

CHOREA.

character of the muscular movements.—Are the movements due to mere clonic jerks of certain muscles, repeated in a meaningless manner (muscular tic), or are they of the character of gesticulations, wriggling, twisting movements, turning the limbs about? Do the movements greatly place the limbs, or after the movements do the limbs always fall back into their previous position? Are the movements independent of voluntary efforts? are they increased by voluntary efforts? are they equal on the two sides? Accompanying muscular weakness. Urine often high sp. gr. and loaded with urea.

SCLEROSIS.*

CHOREA.

Chemical oscillations. In the arm, the main direction of the movement is in spite of the obstacles by the jerks of the arm, and it reaches its goal. The main direction of motion is disturbed from the outset by contradicting movements which cause the goal to be missed. Movements sudden, and unexpected when the limbs are at rest, and apart from the action of the will.

Complications.—In pregnant women miscarriage is frequently attended with danger.

Urine.—Mitral bruits, very common. Urine often scanty and very dense, being loaded with urea; uro-hæmatin often in large amount.

Course.—The connection with rheumatism is shown by its occurrence before, after, or with the chorea. The frequency of cardiac bruits has suggested that the disease is due to rheumatism. If pregnancy excites chorea, there has usually been chorea in childhood. Chorea most common in females, and in childhood near puberty. Exciting causes—light, falls, etc., over-work at school, imitation. Enquire of the symptoms before occurrence of acute movements; whether fidgety, frequently dropping things, clumsy, nervous.

* Charcot: New Syd. Soc. Trans.

HYSTERIA.

Describe briefly patient's complaints. State if able to perform ordinary work ; if not, say why. Enquire if any "attacks, fits, or **Convulsions** occur ;" if they do, note time and circumstance. Note general condition of **Nervous System** ; signs of **Brain Disease**.

Motor Power.—General character of movements, whether active or sluggish. Test reflex excitability.

Sensation.—Should be examined carefully. Globus (sensation of a ball rising in the throat and choking). **Headaches**. **Neuralgia**, specially **Infra-mammary Neuralgia**, and of Nerve V. Look for **Muscular Anæsthesia**. Note mental condition and **Intelligence**.

Causation.—Almost exclusively in female sex ; common in early life ; may be very persistent.

EPILEPSY.

A condition of disease characterized by convulsive paroxysms with loss of consciousness. Look to the general condition of the **Nervous System**, and signs of **Brain Disease**. See **Convulsions**. Note history of onset, frequency of paroxysms, their periodicity and characters, condition in intervals of the paroxysms.

Paroxysms.—Note state of consciousness, whether persistent, partially or wholly lost. Note carefully the degree, kind, and range of **Spasm**, whether **Tonic** or **Clonic**. The amount of fixation of respiratory muscles and signs of cyanosis. Whether head is drawn to one side, face distorted, or signs of opisthotonos. Position of eyes and state of pupils. Look for spasms in muscles supplied by cranial nerves, and one-sided, local, or repeated movements. Condition of sphincters. Temperature, pulse, heart. Next passed urine.

Starting points.—(1) Hand, usually index finger, thumb, or both ; (2) face, usually near mouth, or tongue, or both ; (3) foot, usually great toe. Note range of spasm.

HYSTERIA.

The will is defective ; all voluntary movements are usually sluggish and wanting in energy, but movements excited by emotion may be in excess. The condition is most common in young females, and is frequently associated with disordered menstruation. A special character is the liability to attacks of convulsive nature. Disturbance of *Sensation* is very common, sometimes assuming the form of hemi-anæsthesia, one half the body having lost sensibility, or hyperæsthesia. *Functional Paralysis* is common in this condition ; it may be paraplegic, hemiplegic, or of a single extremity—functional aphonia. *Spasm of Muscle*, more or less continued, is not uncommon, thus causing contraction of a joint, talipes, or a phantom tumour in the rectus abdominis. Among signs of disturbance of organic nerves are *Vomiting* and *Angina Pectoris*.

causation.—Inherited tendency to neuroses. Disordered menstruation. Depressing mental circumstances.

EPILEPSY.*

Symptoms of the Attack.—Stage I. Sudden loss of consciousness ; tonic rigidity of muscles ; arrested respiration, often with a cry due to forcing air through closed glottis. Pallor or duskiness. Pupils dilated. Stage II.—Unconsciousness continues ; clonic convulsion ;* laboured breathing and foaming ; profuse sweating. Stage III.—Partial return of consciousness and voluntary power.

Classes of Paroxysms.—I. Loss of consciousness without evident spasm.

II.—Loss of consciousness with local spasm.

III.—Loss of consciousness with general tonic and clonic convulsion.

IV.—Without complete loss of consciousness, convulsion being general or partial (abortive epilepsy).

e petit mal = classes I. and II.

* Dr. Reynolds' "System of Medicine."

EPILEPSY—*continued.*

Premonitory symptoms.—Mental condition, excitability, dullness, vertigo, dysæsthesia. Aura epileptica strictly implies a sensation of wind blowing upon a limb. An aura may commence in a limb, or the epigastrium, or in the pharynx, in each case passing upwards towards the brain. An aura may commence in an organ of special sense, *e.g.*, the vision of a shape or colour, a "nasty taste," a sound, a smell, a mental sensation. The aura is immediately followed by loss of consciousness.

Sequelæ.—Permanent impairment of intelligence and mental capacity. **Vertigo.**

Complications.—Post-epileptic mania may succeed the paroxysm; in this state acts of violence or homicide may be unconsciously performed. In a condition after the paroxysms termed "reduction" the patient may perform unconscious acts, *e.g.*, place things in strange places.

Causation.—Age, sex, psychical causes, and heredity. The commonest antecedents are reflex causes, teething, intestinal worms; physical causes, *e.g.*, blows on head, exposure to great heat.

Commonest in female sex and from thirteen to sixteen years of age; may be secondary to other organic changes; heart disease is common.

CONVULSIONS.

EPILEPTIC

or

HYSTERICAL.

<i>Onset.</i> —Sudden, often with an aura. Loss of consciousness usually complete.	Less sudden, with emotional disturbance. Loss of consciousness, more protracted, or very apparent.
<i>Prodroma.</i> —Aura epileptica.	Globus hystericus.
<i>Asphyxia.</i> —Often very complete.	Flushed; not asphyxiated.
<i>Face.</i> —Features distorted.	Not distorted.
<i>Coma.</i> —Usually profound, with stertorous breathing. Con-junctiva insensible.	Insensibility complete. Re-flex movements of eye usually continue on touching it.
<i>Subsequent state.</i> —Coma ; stupor ; drowsiness. Subjunc-tival hæmorrhages.	Exhaustion.
<i>Pyrexia.</i> —May arise if much tonic spasm is present.	Temperature normal.
<i>Sleep.</i> —Common during sleep and when falling asleep.	Usual during day-time when others are about.
<i>Tongue.</i> —Often bitten.	Not bitten.
<i>General condition.</i> —Signs of epilepsy.	Signs of emotional dis-turbance.
<i>Urine.</i> —Occasionally contains albumen or sugar.	Copious, limpid, light-coloured, sp. gr. low.

CEREBRAL TUMOUR.

Special symptoms.—**Vomiting, Head-pain, Paralysis of Cranial Nerves, palsy of Special Senses, Optic Nerve changes, Convulsions, Hemiplegia, or other form of Paralysis.** Temperature sometimes very high, without any inflammation.

Look for Syphilis, Scrofula, Phthisis, Cancer, or new growth in other parts. See **Motor Power**, and gait in walking. Examine urine for sugar and albumen.

Causation.—**Syphilis.** Scrofulous diathesis leading to tubercular mass. Tubercular tendency. **Cancer.**

Tumour may be caseous mass, gumma, glioma, cancer, growth of pituitary body, cyst, hydatid, aneurism, blood-cyst in membranes, exostosis.

CEREBRAL MENINGITIS.

Note symptoms, with date and manner of commencement.

Special symptoms.—See general condition of **Nervous System**, signs of **Brain Disease**, vomiting, paralysis of **Cranial Nerves**, intermittent pulse. Look for signs of **General Tuberculosis, Phthisis, strumous disease.** Ophthalmoscopic examination may show tubercles in the choroid. Note eyes, their movements, strabismus, state of **Pupils**, photophobia; general state of nutrition. Take temperature. Examine lungs as to phthisis and recent pneumonia or pleurisy. Examine urine, and note whether it be retained.

Look for Head-pain, Vomiting, ear disease, Syphilis. Take temperature.

CEREBRAL TUMOUR.

sin may be localized, and permanent, or intermittent with exacerbations. **Vertigo** is common. Hearing is not commonly palsied. Urine may be saccharine.

Seizures, partial, clonic, or tonic, not uncommon; they may resemble epilepsy, but usually differ from such attacks as follows:—1. Irregular in development, with less loss of consciousness and no asphyxia or subsequent coma. 2. Not specially a disease of female sex or early period of life. 3. Less tendency to mental disturbance. 4. No special hereditary inheritance of neurosis. 5. Characteristic symptoms of tumour develop.

Course of the disease is generally slow. **Hemiplegia**, if present, usually develops slowly; if on the right side may be accompanied by aphasia. Preceding death the temperature often rises high.

Prognosis.—Cerebral tumour may cause ventricular effusion, resembling **Hydrocephalus**.

CEREBRAL MENINGITIS.

often insidious; poorliness and loss of appetite, with head-pain and vomiting. Temperature is a very uncertain sign; vomiting, though important when present, is frequently absent throughout. Intermittence of the pulse and paralysis of a cranial nerve are very important signs. Abscesses may occur in the choroid, independently of meningitis.

Prognosis.—**Miliary Tuberculosis**. Disease of ear. **Syphilis**. Injury to head. **Cerebral Tumour**.

CHRONIC HYDROCEPHALUS.

History of family ; of the pregnancies and labours of the mother. State of the head at birth, or date at which symptoms were first observed. Enquire for **Convulsions**. Note general **Motor Power**. **Sensation**. Nutrition ; power to hold head up. Eyesight. Hearing. **Intelligence**, whether child notices sounds and colours, and plays with toys, or is backward for age.

Head.—Is it held well up, well shaped ; its circumference, measurement from ear to ear, over the vertex, and from the nose to the occiput. State of sutures and fontanelles, whether patent or ossified. Take tracings of skull with cyrtometer.

Eyes, whether of normal direction ; condition of optic nerve. Dentition. Look for signs of **Rickets**.

Look for signs of Defective Development.

CHRONIC HYDROCEPHALUS.

be mistaken for the large head of rickets. Congenital hydrocephalus usually causes difficult labour. Head may be normal at birth, subsequently enlarging. Sometimes accompanied by spina bifida. Tendency to enlargement of the head is progressive.

HYDROCEPHALUS.

is of **Rickets**, but
Brain Disease.

is a tendency to globe; eyes depressed; strabismus; optic nerves thin. Cranial bones thin. Head held up. Parosmia, or a contracted

RICKETS.

Signs and symptoms of
Rickets.

Head large, tending to broadness and squareness; there may be irregular thickening of bones. No paralysis, head held up, child playful.

try to increase of size of head to body, by measurements. No enlargement; parosmia continuing. Inebecile. Optic atro-

As signs of **Rickets** pass away the relative size of the head less noticeable. May have good power.

ALCOHOLISM.

See **Nervous System**. See **Motor Power**. Tendency to tremor; tongue tremulous, coated, glazed. Muscular weakness and want of muscular co-ordination. See co-ordination of the limbs. Muscular inquietude; muscular fidgetiness. Look to **Muscular Sense** (usually diminished). **Paralysis**, **Paraplegia**. See signs of disease of **Spinal Cord** and **General Paralysis**. Flushing and congestion of the face and eyes. **Vomiting**, especially in the morning. Conditions of **Sleep**. Neuralgic pains. **Anæmia**.

Mental disturbance.—Deterioration of mental power, restlessness, loss of memory, hallucinations, delusions. Mental alteration, *e.g.*, inaptitude for business, avoidance of friends.

Sensation.—Cutaneous sensibility, dysæsthesia, *muscæ volitantes*, buzzing in ears, vertigo. Note state of nutrition.

Complications.—Look for signs of disease of liver, kidneys, vascular system, emphysema. Acne rosacea of nose. Bronchitis. Pneumonia. **Delirium** and symptoms of delirium tremens.

ACUTE ALCOHOLISM.

Excessive dose may produce **Coma**; breathing stertorous; breath smelling of alcohol. Appearance of face. Examine urine for albumen and alcohol. Look for signs of general condition of **Nervous System**. **Vomiting**. **Paralysis**. See causes and examination of cases of coma. Examine heart and condition of blood-vessels.

Look for—Injury to head. **Uræmia**. Simple exhaustion. **Meningitis**. There may be **Albuminuria** from acute renal congestion. Complications of chronic alcoholism.

ALCOHOLISM.

principally produces nervous symptoms ; affects next the digestive system. Nutrition may become much impaired. In chronic cases, kidneys and liver often become cirrhotic. Vascular system degenerates. **Emphysema.**

Chronic Cases.—In advanced stages, the lower extremities may become unsteady, hands and fingers tremulous, so also the tongue. At first the tremors may be restrained by voluntary effort. *Acne rosacea.*

Diagnosis from—

Commencing **General Paralysis of Insane**, mind depressed.

Paralysis Agitans.

Plumbism, with tremor and delirium.

Locomotor Ataxy.

Paraplegia, from **Disease of Cord.**

Senile Degeneration.

Sclerosis.

Hysteria.

Nervous malaise, from simple dyspepsia.

Complications.—**Cirrhosis of Liver and Ascites.** **Chronic Bright's Disease.** Atheroma or **Degeneration of Arteries** and small vessels. Chronic gastritis. Fatty degeneration of heart and liver.

ACUTE ALCOHOLISM.

acute symptoms may be due to **Delirium Tremens**, or to an excessive dose causing toxic effects, *e.g.*, coma, etc. When drunk—**Coma**, face livid, breath smelling of alcohol, tendency to vomit. Vomits or washings of stomach contain spirit. In very deep coma there may be strabismus. There may be great excitement in place of coma. Cerebral hæmorrhage may occur during intoxication. Alcohol in urine.

DELIRIUM TREMENS.

Delirium, delusions, illusions of **Sight** and hearing. **Vomiting**, inability to take food. Intense restlessness. Look for the degenerative changes of chronic alcoholism. Specially examine lungs, urine, heart, and pulse. Note muscular condition, general strength, and power of movement. **Tremor**, subsultus tendinum. Sleep; degree of consciousness.

Complications.—**Typhoid State**. Subsultus. **Coma**. Heart failure and pulmonary congestion. Syncope. Albuminuria. **Pneumonia**. Rapid development of phthisis.

INSANITY.

1. Mania. 2. Monomania. 3. Melancholia. 4. Puerperal mania. 5. Moral insanity. 6. Dementia. 7. Idiocy, including imbecility. 8. **General Paralysis** or **Paresis**.

Causation.—Heredity of primary importance; enquire back to the third generation in the families of each parent. See also as to collateral relations.

Alcoholism. Habits and mode of life. Mental anxiety. Injuries to head.

Signs of Insanity.—Talking to self, fantastic dress, refusing food, squandering property, kleptomania, self-injury, violence, delusions, melancholy, incapacity for business, avoiding friends, delirium. See signs of **Brain Disease**.

Illusions of the senses. Sight; they may be coloured, moving forms. Hearing, smell, taste, touch. The perception of the sense is mistaken, and the impression made is false.

Complications.—**Phthisis**, fragile bones, heart disease, **Epilepsy**. Attacks of partial coma.

Examination of Patients.—Test **Motor Power**, **Pupils**, **Muscular Sense**, **Nervous System**, **Sensation**, heart and lungs.

DELIRIUM TREMENS.

Usually the effect of long-continued drinking, with dyspepsia and deprivation of food. Commences with disturbance of general condition of the **Nervous System**; diminished motor power. Insomnia, night-wandering, and horrors, with delusions, passing on to delirium with violence and suicidal tendency. Delirium may be busy, low muttering, or talkative.

Complications.—Sudden syncope during violent struggling in the delirium may lead to sudden death.

GENERAL PARALYSIS OF THE INSANE.

Characterised by progressive diminution of mental power, followed by paralysis, involving the whole of the muscular system. Pupils show want of symmetry of size, and want of mobility. Mental condition characterized by an exaggerated feeling of power, extravagant exalted ideas, loss of memory, attacks of excitement and violence. Hallucination; delusion.

Motor Power.—Failure first seen in tongue; inaccurate articulation, fibrillar trembling of the tongue. Pupils unequal. Automatic and reflex actions lessened; electric contractility of muscles retained. Teeth-grinding. Late in disease, sphincters lose their control, and there is tendency to choking. Bones may be very brittle.

Sensation.—Cutaneous sensibility usually diminished, and later lost. **Muscular Sense** lost. Attacks of excitement and violence; epileptiform convulsions. Face becomes expressionless.

Diagnosis from Alcoholism.—Ideas of exaltation; pupils unequal: effect of removing alcohol; paralysis of sphincters.

Causation.—*Inheritance*, intemperance; most common in men.

PARALYSIS AGITANS.

State principal sites of tremor ; hemiplegic type, paraplegic, or confined to one extremity. Examine **Motor Power** ; whether movements of head, face, tongue ; if speech be affected. Expression of face. Power to walk ; gait in walking. Note any tendency to involuntary forward or backward movement, or dragging of limbs, etc. Ability to perform certain acts, walk, hold out limbs, pick up a pin, or write ; keep specimen of writing. Let him hold a glass of water, and carry it to his mouth. Note effect of emotion on tremors. Describe the **Tremor**.

SCLEROSIS OF CORD.

Examine condition of the **Motor Power** and reflex excitability. **Tremors** ; see whether they cease during repose and are increased by voluntary acts. Let patient raise a glass of water to his mouth, and describe the result. Let him stand and walk ; then close his eyes and again perform the same acts. Note whether tremors are fine or coarse. Note the extent of parts affected by tremor ; whether head, trunk, and all the extremities are affected. Examine for **Brain Disease**. Ankle clonus.

PARALYSIS AGITANS.

Characterized by muscular tremor, constant even in repose; muscular power diminished. Head not tremulous, but may be shaken by movements of the body. Tremor consists of jerks, more regular and rapid than in **Disseminated Sclerosis**. No real difficulty of speech, but the utterance is slow and with jerk-like effects. Respiratory movements not affected. In advanced cases muscular rigidity may lead to deformity; this is specially seen in the hand. There may be a subjective sensation of heat. No **Nystagmus**. Face stiff, expression still.

SCLEROSIS OF CORD.

Characterized by muscular tremor, increased in direct proportion to the extent of any movement executed. It is only manifested by voluntary movements of some extent, and ceases when the muscles are in complete repose. The oscillations are larger than in **Paralysis Agitans**, and more resemble the gesticulations of **Chorea**. Voluntary acts may be performed despite the tremors. Closing the eyes does not affect the tremors, as in ataxy. Movements are not seen, independent of voluntary efforts, as in chorea. The head is usually affected with tremor; **Nystagmus** is common. Patellar tendon reflex is exaggerated in sclerosis, obliterated in **Ataxy**.

TETANUS.

General condition.—T. = ; sweating. Occurs in the robust rather than in the weak.

Special condition of muscular system.—Tonic spasm. Trismus (lock-jaw). Opisthotonos, *i.e.*, body drawn backwards, or emprostotonos, body drawn forward. Eyes may be retracted from spasm of recti muscles. Face set hard with sardonic grin. Tetanic convulsions frequently repeated.

Modes of onset.—Commencing about six days or sooner after injury, may be two to four weeks after injury. Injury may have been overlooked, and the wound healed. The earlier the commencement, the more rapid the case and the more certainly fatal the result. There is a gradual progress of the symptoms. Tonic contraction commences with trismus, extending to throat, back of neck, abdomen.

Causation.—Injury, blows, burn ; forcing bodies under skin or under nail. Excited by exposure to cold ; war ; operation. Injury to nerve. Males rather than females. Idiopathic cases due to exposure to wet and cold : this not common. May occur spontaneously in infants within eight days of birth.

Diagnosis.—From hysterical opisthotonos. Strychnia poisoning. Hydrophobia.

TETANUS.

riod of onset ; not con-
with food.

iffness first perceived
; it then progressively
downwards, attacking
y and limbs, the hands
ig commonly affected
last. Progressive in-
with somewhat gradual

iration of the case
ess than twenty-four
Seldom fatal in idio-
ases.

scovery of nux vomica,
ia, brucin, or other
in food, vomits, or
s of stomach.

iscular rigidity almost
intermission.

STRYCHNIA POISONING.*

1. Onset soon after food.

2. Sudden and violent onset
of symptoms: commences with
shivering, gasping for breath,
trembling ; the body and limbs
are then simultaneously af-
fected, hands clenched, feet
curved ; at a later date the
jaw becomes fixed during a
paroxysm.

3. Rarely survive two hours
after a fatal dose.

4. Absence of wound, ulcer,
or traumatism to account for
tetanus ; exposure to cold, or
special nervous susceptibility.

5. Intervals or remissions of
rigidity of muscles.

* See Taylor on the Poisons.

LOCOMOTOR ATAXY.

Examine condition of the **Motor Power**; especially the gait in walking, and co-ordination of the limbs. Let him walk with his eyes open, then shut; let him walk with slight assistance or using a stick. Also test power to keep knee flexed or extended. Test upper extremities, *e.g.*, precision with which he can touch an object, his eye or nose, or execute definite movements. **Reflexes**, and patellar tendon reflex.

Electric Tests.

Sensation.—Tactile sensibility; sense of heat and cold. Subjective sensibility; consciousness of ground in walking; perverted sensations in lower extremities. **Sight**, reaction of Pupils, **Ophthalmoscopic Appearances**.

Examine the joints and skin. Look for signs of **Disease of Cord**. Temporary defects of third nerve common. Bowels and action of bladder sluggish. This disease is often associated with **Syphilis**. Occasionally gastric crises or attacks of vomiting.

LOCOMOTOR ATAXY.

characterized by difficulty in walking, especially with the eyes shut; there being no motor paralysis and no loss of nutrition of the lower extremities—patient still having voluntary power to keep the limb flexed or extended with good force. Commonest in males, and at ages thirty-five to fifty years. In walking there is exaggeration of the movements; the feet are lifted too high and the heel brought suddenly down. The lower extremities are the most affected, but there may be want of co-ordination of the upper extremities also. Electric irritability not impaired. **Patellar tendon reflex** obliterated more or less completely. In early stages “lightning-pains” in legs and back are usual; they may last for years and cause much distress.

pupils usually small, inactive to light (vaso-motor palsy); contraction for near accommodation intact, *i.e.*, ciliary muscle sound—it is supplied by Nerve III.; ptosis sometimes. The optic nerves sometimes become white from atrophy. Large joints may be the seat of effusion and chronic absorption of the cartilages. It is distinguished from disseminated sclerosis by the marked increase of symptoms produced by closing the eyes; this does not so modify the rhythmic jerks of sclerosis.

forming ulcer of foot; deafness from atrophy of nerve—sometimes found.

ine may be retained.

INFANTILE PARALYSIS.

History.—General condition of health. Look specially for **Rickets**. Test general strength of motor power. State which extremity is affected; note its state of nutrition. Examine the separate muscles. Test reflex action and sensation. **Electric Tests.** Observe temperature of the paralysed limb, and the condition of the skin.

Upper extremity.—Can he move the fingers separately? point with index and little fingers, etc.? Movement of wrist; power of pronation and supination; hold out the limb from the shoulder; put his hand to back of his head. Measure length and circumference, and compare with opposite side.

Lower extremity.—Can child walk, stand, move toes, flex ankle and knee, or hold out the limb? When sitting down can he get up?

Causation.—Age six months to six years. Equally in both sexes. Sequel to exposure to cold or an exanthematous fever. Possibly due to dentition.

Rigid contraction frequently causes talipes in leg; such deformities much less common in upper extremity. Occasionally the bones do not grow in length, and a shortened limb results in after life.

INFANTILE PARALYSIS

urs during ages from six months to seven years, attacking a certain muscle or group of muscles. It is unattended by pain or signs of brain disease. It often occurs in children apparently perfectly healthy. The attack of paralysis is never repeated.

on.—There may be premonitory symptoms two or three days, or more; then the limb may be found paralysed. Such premonitory symptoms may be wholly absent. Onset not usually attended with much disturbance of the general condition of the nervous system. Paralysis may be noted without any premonitory symptoms.

se of disease.—Usually the general health remains good. Most of the muscles first paralysed usually regain power in two or three weeks, leaving some muscles, or a single muscle, *e.g.*, deltoid, permanently weakened. In regaining power the order of recovery is the reverse of that seen in paralysis from brain disease; the finer movements are first regained, *e.g.*, movements of fingers and toes before the wrist and ankles. The muscles permanently paralysed atrophy. The growth of the limb may be checked, especially in the lower extremity. Permanent paralysis may be in one leg only. The palsied limb becomes cold and bluish.

ation not affected. **Reflex excitability** impaired or abolished, and electric excitability lost.

GRAVES' DISEASE (Exophthalmic Goître).

Exophthalmos.—Frequently eyelids cannot close over eyeballs, the eyes remaining open even during sleep. The degree of prominence of either eye is usually equal, but may be more marked on one side. Eyelids tremble on endeavouring to cover eyeball. Eyes appear staring, bright, and glistening. Test sight and optical refraction, and movements of eyes. Ophthalmoscopic appearances. Examine pupils.

Goître.—This sign may be absent. Enlargement usually moderate, with a thrill felt and hæmic murmur on auscultation. It is very rarely cystic. Thyroid enlargement usually first seen on the right side.

Vascular system.—Throbbing in arteries of neck, and in thyroid. Hæmic bruits over goître and vessels in neck. Violent and frequent action of heart even without exertion. Left ventricle may be dilated. Valvular lesion not very common.

Complications.—Dilatation of heart. Asphyxiating attacks. Diarrhœa. Vomiting. Bronchitis. **Paraplegia**. Headache.

PLUMBISM.

General condition.—**Anæmia**; emaciation; gout.

Digestive system.—Attacks of colic and constipation, may be with vomiting, nausea, loss of appetite. Blue line on margin of gums, especially opposite the teeth. Abdomen retracted. Breath fœtid.

Nervous system.—Paralysis usually of extensors of forearm, attended with atrophy and loss of electrical reaction. Usually paralysis is preceded by attacks of colic. Look to the general condition of the **Nervous System**. **Motor Power**. **Sensation**. See **Optic Discs**.

GRAVES' DISEASE (Exophthalmic Goître).

mainly characterized by prominence of the eyeballs. Pulsating goitre. **Palpitation.** Usually there is **Anæmia** and disordered menstruation. Emaciation. Mental irritability and want of sleep. See **Motor Power.** There is a tendency to intercurrent attacks of diarrhoea; appetite capricious. Occasionally enlargement of liver, spleen, and mammae. Frequently there is increase of the symptoms at the menstrual periods.

apils.—No alteration from the normal; natural size and activity; accommodation normal.

rusation.—Usually develops in females above age of puberty; rare in men. May date from a mental shock or period of over-work. It is connected with anæmia and disordered menstruation.

PLUMBISM.

characterized by colic, anæmia, blue lines on gums, **Paralysis** of extensors of forearm, and brain disturbance.

ervous system.—There may be profound disturbance of the brain. Optic neuritis. **Delirium.** Epileptiform convulsions. **General Tremors.** Palsy of the extensors of the forearm, principally marked on the right side; the supinator longus and extensor longior carpi radialis escape palsy.

sensation may be at fault; numbness in limbs, neuralgia, headache.

omplications.—**Gout; Bright's Disease;** optic nerve changes; paralysis.

DIPHTHERITIC PARALYSIS.

History.—Previous attack of sore-throat. Possible source of infection; evidence as to the diphtheria.

General condition.—W. = ; nutrition; T. = .

Mouth.—Condition of mucous membrane as indicating previous inflammation. Movement of palate; fauces; tonsils; pharynx.

Nervous system.—Speech, whether nasal or twangy, understandable or voiceless. Pain, giddiness.

Motor power.—Ability to stand and walk; gait. Eyes, their movements; vision, accommodation, pupils.

Sensation.—Dysæsthesia, with numbness and formication, precede palsy in limbs.

Urine may be albuminous.

HERPES ZOSTER.

History of illness; date of onset of symptoms, and of appearance of rash. Enquire as to recent use of arsenic.

Look to general condition, debility, **Anæmia**, etc.

Look for signs of **Neuralgia**, condition of skin at seat of inflammation, sensibility, subjective pain, etc., tenderness, tenderness present along the course of the nerve supplying area affected.

Note any nutritional effects on parts affected, ulcerated areas, scars (with Nerve V. see Iritis); note subsequent state of

Sensation.

DIPHTHERITIC PARALYSIS.

.—Palsy follows the sore-throat in two to six weeks. Primary illness rarely attended with laryngitis. Complains usually as to motor power, sight, speech, deglutition.

Area of muscular weakness may be limited to fauces and accommodation of eyes. The limbs, if much weakened, atrophy proportionally. Respiratory muscles may be involved. Palsy is usually symmetrical; lower extremities more palsied than upper. Eye-muscles, and tongue and face may be palsied.

is.—Cases usually recover. Danger from heart failure, shock, paralysis of respiratory muscles.

HERPES ZOSTER.

usually occurs in young subjects; it has been noted as common in persons taking arsenic. The disease does not burn. Pain precedes the eruption; it may be severe and last for days. The rash is vesicular, vesicles appearing along the area of a cutaneous nerve; the vesicles contain clear watery fluid, and may have inflamed bases. The vesicles seldom cross the median line. Vesicles dry up in a scab; in debilitated subjects ulceration may follow.

DISEASES OF THE VASCULAR SYSTEM.

HEART—PHYSICAL EXAMINATION.

Inspection.—See front of chest. Look for and define apex-beat; it should be seen in fifth space an inch below, and internal to, left nipple. Look for other sites of pulsation. Pulsation of left auricle may be seen in third space.

Palpation.—Feel the general force of the cardiac impulse, indicating strong or weak action, **Hypertrophy** or **Dilatation**. Determine area of impulse and site of apex-beat. Search for a thrill, especially towards apex; feel first with tips of fingers, afterwards with ends of metacarpal bones. Look for friction fremitus. See pericarditis. See **Displacement of Heart**.

Auscultation.—Listen for 1st and 2nd sounds; each should be clear "lub-dub."

1st Sound.—Systolic, coinciding with the impulse. Loudest towards apex; to be traced upwards to the base, towards epigastrium and to axilla. Note character of sounds, sharp, clear, feeble, dull, prolonged, or short, or much resembling 2nd sound (tic-tac). Accompanying bruits are termed systolic.

2nd Sound.—Diastolic, coinciding with subsidence of cardiac impulse. Loudest at level of second costal cartilage; aortic valves to the right side (aortic cartilage), pulmonary valves to the left (pulmonary cartilage). Trace the sound to the apex. The whole 2nd sound may be accentuated, or either the aortic or pulmonary only. It may be reduplicated. Accompanying bruits are termed diastolic.

Cardiac Murmurs.—The fact of a cardiac murmur being decided, determine its periodicity—systolic, diastolic, or presystolic; the site of maximum intensity; and relative conductivity in various directions, towards base or apex, to axilla or along sternum, or along the vessels at the base. Observe if audible by spine or at angle of left scapula. Character of murmurs—plain bellows sound, musical, rasping.

HEART—PHYSICAL EXAMINATION.

Inspection.—May detect a diffused wave of impulse, *e.g.*, **Pericarditis**. Hypertrophied left auricle may be seen pulsating in mitral stenosis or contraction of left lung. Abnormal site of pulsation from **Aneurism**, usually in right third space. See bulging of precordium.

Palpation.—Pulsation may be detected in epigastrium in dilatation. Thrill systolic over aortic cartilage (second right) in aortic stenosis or aneurism; at apex in mitral regurgitation. A diastolic thrill in base at aortic regurgitation; at apex just before the systole in mitral stenosis. Strong heaving impulse with hypertrophy.

Auscultation.—Determine if heart's sounds are healthy and in due rhythm; if accompanied by, or replaced by, abnormal sounds (bruits or murmurs) which are generally due to pathological conditions of the valves. Note they may be due to **Anæmia** or **Aneurism**.

1st Sound.—Indicates the muscular condition of the heart, and how it is working; strong in **Hypertrophy**, weak in degeneration of the walls. It may be reduplicated; may be masked by **Emphysema**. Anæmic bruit at base common.

2nd Sound.—Due to closure of semilunar valves, aortic and pulmonary; each should be examined separately. Pulmonary 2nd sound not often accompanied by a bruit unless from anæmia; it is accentuated in recent pulmonary congestion, as from recent mitral regurgitation. Aortic 2nd sound accentuated in obstructed arterial (systemic) circulation, as in **Bright's Disease**.

Cardiac Murmurs.—If the normal heart's sounds are heard, and the other physical signs and the pulse are healthy, we may conclude that the heart is healthy. If a bruit be heard, look for all the **Signs of Heart Disease** and the presence or history of some cause likely to produce valvular defects. If there be no other proof of heart disease than the bruit, look for signs of **Anæmia** and anæmic bruits. The character of murmurs often changes.

HEART—PHYSICAL EXAMINATION.

Percussion.—Determine and mark out the area of relative and absolute precordial dulness. In health it extends from about the third left cartilage to the apex-beat, being limited below by the line of the liver, and not crossing the median line. Area of dulness may be diminished by atrophy of the heart, as in old age, or heart may be overlapped by **Emphysema** of the lungs. The area may be increased by pericardial effusion as a triangle, larger than the normal area, with its apex towards the top of the sternum.

PULSE.

Usually felt in radial artery; it may be examined in any superficial artery.

1. *Frequency.*—Frequent or infrequent refers to the number of pulsations per minute. $P. =$.
2. *Quick or slow.*—Refers to the time occupied by each beat, not including the interval between it and its successor.
3. *Rhythm.*—Regular or irregular implies the order of succession. Intermittent, the occasional dropping of a beat.
4. *Large or small.*—Refers to the degree of dilatation of the artery.
5. *Jerking or collapsing.*—Full, rising quickly and falling suddenly.
6. *Tension.*—Soft or hard. Felt by the fingers and measured by the force required to extinguish the pulse by pressure.
7. *Dicrotous.*—The wave is double-headed and the pulse soft.
8. *Locomotor.*—When the artery is seen to travel like a snake under the skin.

State of arteries.—Examine radial, brachial, femoral, dorsalis pedis, temporal, etc. The artery as a piece of tissue may be hard or soft, irregular on the surface, dilated, etc.

HEART—PHYSICAL EXAMINATION.

Percussion.—Hypertrophy of right ventricle increases the width of the area of dulness, so that it may reach to the right of the median line. Hypertrophy of the left ventricle extends the dulness outwards and downwards. Abnormal areas of dulness adjoining the heart may be due to **Consolidation of Lung**, mediastinal tumour, or **Aneurism**.

PULSE.

May indicate the condition of the cavities of the heart and valves, and the state of the nervous system.

. *Frequency.*—High in fever and in mental excitement; in disease of the valves and walls of the heart; in **Graves' Disease**.

. *Quickness.*—Chiefly affected by conditions of the nervous system.

. *Rhythm.*—Irregularity may depend upon valvular lesions, especially mitral disease, or on the state of the muscular walls of the heart; brain disease, *e.g.*, **Meningitis**; reflex causes, *e.g.*, dyspepsia.

. *Large or small.*—Depends upon strength of the left ventricle and condition of valves. It may be small in mitral disease or depressed innervation.

. *Jerking.*—In aortic regurgitation with hypertrophy of left ventricle. This character may be less marked if mitral disease coexist.

. *Tension.*—High in **Chronic Bright's Disease**, and in the cold stage of ague. Low in **Typhoid State** and conditions of adynamia.

. *Dicrotous.*—In fevers, especially in the typhoid state.

. *Locomotor.*—Indicates a hard, thickened, or **Atheromatous Artery**, or an hypertrophied left ventricle.

tate of Arteries.—Rigid, tortuous, and rough upon the surface in atheroma. See **Vessels, Disease of**.

PASSIVE (Cardiac) CONGESTION.

Starting from an obstructed circulation on the left side of the heart, *e.g.*, mitral obstruction.

Pulmonary veins overfull (open into left auricle) ; receive blood from pulmonary capillaries and some of the bronchial capillaries.

Bronchial capillaries overfull ; hence tendency to **Bronchitis**.

Pulmonary capillaries overfull ; hence **Pulmonary Œdema**, *i.e.*, effusion into air vesicles.

Pulmonary artery (leading from right ventricle) conveys blood to the overfull pulmonary capillaries ; hence tension rises in the pulmonary artery, and pulmonary 2nd sound may be accentuated.

Right ventricle (drives blood into the pulmonary artery, which is overfull). It becomes over-distended and dilated ; this may lead to **Tricuspid Regurgitation**.

Right auricle (drives blood into the right ventricle, which is overfull). It receives blood from superior and inferior venæ cavæ and bronchial veins.

Superior vena cava receives blood from *bronchial veins* ; these carry blood from bronchial capillaries (which also partly empty into pulmonary veins) ; hence **Bronchitis**. The bronchial veins also receive blood from the pleura ; hence **Hydrothorax**.

Jugular veins, and the veins of the head and upper extremities, send their blood to the superior cava ; hence **Cyanosis of the Face**, jugulars standing out in the neck, **Congestion of the Brain**, **Œdema** of the upper extremities. If there be tricuspid incompetence, jugulars may be seen and felt pulsating.

(Continued next page.

PASSIVE (Cardiac) CONGESTION.

Inferior vena cava receives blood from the *hepatic vein*; hence congestion of intra-lobular veins and hepatic capillaries in the lobules causes **Enlargement of the Liver** and **Jaundice**, also obstruction to the outflow from the *vena portæ*, and congestions of the vessels emptying into the *portal system*, viz., gastric, splenic, intestinal, hæmorrhoidal; hence **Spleen large, Ascites, Hæmatemesis, or Melæna.**

Renal veins (branches of the inferior cava) receive the veins which collect blood from the capillary plexus surrounding the uriniferous tubes; this plexus becomes primarily congested, and as it receives blood from the afferent vessels of the Malpighian tufts, these capillaries become secondarily congested, leading to **Scanty Secretion of Urine** and **Albuminuria.**

Iliac and femoral veins return blood from the lower extremities, and their over-fulness leads to capillary congestion and **Œdema of the Feet**, the pressure being the greatest in the most dependent set of capillaries.

IMPORTANT ANASTOMOSES.

In portal obstruction, anastomosis of inferior hæmorrhoidal veins of the internal iliac with branches of the inferior mesenteric of the portal system. In portal obstruction, blood flows from intestines through the rectum to the internal iliac veins. Piles result.

In obstruction of inferior (abdominal) vena cava, e.g., by pressure of a growth or tumour, anastomosis of epigastric veins of the iliacs with mammary branches of superior cava. Enlarged veins on abdominal walls common in ascites.

Radial and ulnar arteries.—When radial pulse is obliterated we may have a return current by deep palmar arch.

MITRAL REGURGITATION.

Inspection. — Apex-beat displaced outwards and downwards; impulse diffused. Right ventricle probably dilated.

Palpation. — Right ventricle usually hypertrophied or dilated. Apex-beat displaced outwards and downwards.

Pulse frequent, small, irregular. Systolic thrill at apex. Heart's action may be irregular.

Auscultation. — Systolic bruit at apex conducted well into axilla, also heard at angle of left scapula. Pulmonary 2nd sound accentuated.

Percussion. — **Dilatation** or hypertrophy of right ventricle.

MITRAL OBSTRUCTION

Pulsation of hypertrophied left auricle sometimes seen in third left interspace. Right ventricle probably dilated.

Right ventricle hypertrophied. Thrill at apex just preceding impulse.

Pulse small.

Presystolic bruit at apex. Pulmonary 2nd sound accentuated from increased tension in pulmonary artery. Aortic 2nd sound feeble at apex. Bruit almost localized to apex-beat.

Left auricle and right side of heart hypertrophied. Left ventricle not hypertrophied.

AORTIC REGURGITATION.

Inspection.—Left ventricle **Hypertrophied**; apex-beat displaced outwards and downwards; much precordial impulse seen. Pulse seen locomotor.

Palpation.—Thrill diastolic, distinct at base; great hypertrophy of left ventricle, precordium thrust forward at systole, etc.

Pulse full and **Collapsing**.

Auscultation.—Diastolic bruit at aortic cartilage and conducted down left side of sternum.

Co-existing **Mitral** disease common.

Percussion.—Area of dulness increased downwards and outwards, from hypertrophy of left ventricle.

AORTIC OBSTRUCTION.

Left ventricle hypertrophied, but less dilated than with regurgitation.

Thrill systolic over aortic valves. Signs of hypertrophy; impulse strong, heaving.

Pulse small or not abnormal.

Systolic bruit over aortic cartilage conducted to right sterno-clavicular joint.

Exclude **Anæmic Bruit**.

HYPERTROPHY OF HEART.

Inspection.—Heart's impulse may be seen over an extended precordial area, shaking and thrusting forward the chest-walls. Apex-beat displaced, usually outwards and downwards. In children the cardiac area may be bulged forward.

Palpation.—Shock of heart against chest-wall very distinct, raising the hand or stethoscope. Impulse felt in several interspaces. Epigastric pulsation if right ventricle is hypertrophied. Pulse full and strong in proportion to the hypertrophy.

Auscultation.—First sound prolonged, dull, strong. Aortic 2nd sound intensified. *Note* diagnosis from pericardial effusion by intensity of 1st sound coinciding with increased area of dulness.

Percussion.—Area of dulness. Left ventricle enlarged downwards and outwards, and may extend a little upwards. Right ventricle enlarged laterally, and may extend to right of sternum.

HYPERTROPHY AND DILATATION.

Causation—

Obstructions in the pulmonary circulation (right heart affected). — **Emphysema.** Chronic bronchitis. Chronic pleurisy. Adhesions of lung preventing expansion.

Obstructions in the aortic circulation (primarily affecting left side).—Arterial disease. **Chronic Bright's Disease,** with thickening of small arteries. General plethora. Repeated pregnancies.

Causes originating in or about the heart.—Primary dilatation, *e.g.*, after fevers, in **Anæmia.** Valvular disease and stenosis of the outlets. Adherent pericardium. Excessive exercise. **Displacements of the Heart.** Malformations. Emotional disturbance long continued. Fatty or other form of degeneration.

DILATATION OF HEART.

Inspection.—Impulse diffused, not bulging or shaking walls of chest. If right ventricle is dilated, epigastric pulsation may be seen, and **Tricuspid Regurgitation** may lead to pulsation in jugular veins. Cyanosis and dropsy common.

Palpation.—Impulse diffused ; it may be heaving if ventricles are hypertrophied, or feeble if walls are degenerate. Impulse may be masked by emphysematous lung overlapping the heart. Pulse weak, especially if walls of ventricle are degenerate.

Auscultation.—If walls of heart are degenerate, 1st sound feeble, short, and much resembling 2nd sound.

HYPERTROPHY AND DILATATION.

Dilatation of either ventricle may exist without much compensative hypertrophy ; perhaps this is most common on the right side. A dilated and hypertrophied heart may be capable of carrying on the circulation so perfectly as to compensate for a valvular lesion, but when degeneration of the heart-walls follows, then signs of **Cardiac Congestion** are apt to supervene. Very great hypertrophy of the left ventricle without any bruit or valvular lesion is common with **Granular Contracted Kidneys**. The cardiac impulse and area of dulness are often masked by emphysema of the lungs ; but still, with hypertrophy the pulse is strong. Examine heart, lungs, urine.

CARDIAC DISPLACEMENTS.

Pleuritic Effusion, pushing the heart to the opposite side; subsequent contraction of lung may draw it to the side affected. Cirrhosis of lung, or other form of contraction, drawing heart to side affected. Cancer of lung, if diffused, dragging it to side affected by contracting. Mediastinal tumour, cancer, **Aneurism**, glandular, pushing heart aside. **Abdominal Tumour**, hepatic, ovarian, etc., pressing up diaphragm may displace heart.

VALVULAR DISEASE.

Causation.—**Rheumatism**, atheroma, rupture of valves, scarlet fever, **Syphilis**, **Alcoholism**, muscular over-strain, congenital heart defect.

HEART DISEASE.

General symptoms.—Seldom attended with pain, unless **Angina**.

General condition.—**Anæmia**, mal-nutrition, **Œdema**, hæmorrhages, faintness, languor.

Digestion.—Dyspepsia, **Jaundice**, **Ascites**, **Liver** large.

Vascular system.—**Palpitation**. Irregularity of heart's action. Hæmorrhages. Cyanosis. See **Passive (Cardiac) Congestion**. Dropsy. **Embolism**. Irregular pulse, feeble, etc. Over-fulness of veins.

Nervous system.—Disturbance of general condition of **Nervous System**. Insomnia. **Vertigo**, **Headache**, **Chorea**, **Convulsions**, **Paralysis**, **Angina Pectoris**.

Respiratory system.—Orthopnœa. Dyspnœa, especially on exertion. Respirations frequent. Bronchitis, **Emphysema**, **Cough**.

Urine.—Scanty, with deposit of lithates. Sp. gr. high. May contain albumen or blood.

CARDIAC DISPLACEMENTS.

diagnosed by palpation and percussion principally. The heart may be raised by pericardial effusion. Apex may be displaced by cardiac **Hypertrophy** or **Dilatation**. Displacement may cause cardiac dyspnoea, etc., in cases of sudden pleuritic effusion.

VALVULAR DISEASE.

note.—Rheumatism usually attacks the mitral valve, and may spread to the aortic. **Atheroma** spreads from the aorta to the valves.

HEART DISEASE.

Look for signs and symptoms of heart disease, then make careful physical examination. Listen for the normal sounds, and bruits accompanying or replacing them. Look for signs of **Hypertrophy** or **Dilatation**. If a bruit heard, look for signs of anæmia. Note state of pulse and respiratory system. Examine arteries. Valvular lesions are often combined, *e.g.*, aortic regurgitation and mitral regurgitation coexisting, etc. Many symptoms result from passive congestion, *e.g.*, œdema, cyanosis, pulmonary œdema, bronchitis, hydrothorax, hæmoptysis, enlargement of spleen, enlargement of liver, jaundice, ascites, albuminuria, congestion of brain. These symptoms then vary with the heart's condition.

PALPITATION.

FUNCTIONAL.

Disturbance of the general condition of the **Nervous System**.

Excessive smoking, or use of tea and coffee.

Often relieved by exercise.

Frequent in recumbent posture.

Mostly in hysterical women.

Attacks intermittent, as causes producing them vary, *e.g.*, dyspepsia, menstruation.

Between attacks heart and pulse natural.

Often accompanied by neuralgic pains.

In attacks face may be flushed, throbbing in ears, tinnitus aurium.

Gout; masturbation; want of sleep; **Graves' Disease**.

Causation.—Heart disease; **Dilated Heart**; **Hysteria**; hypertrophied heart with degeneration. Reflex uterine; dyspepsia.

Examine heart, its sounds, impulse, regularity. Look for **Vascular Degeneration**.

ORGANIC.

Physical signs of disease of walls of heart or its valves.

Accompanying general **Signs of Heart Disease**.

Excited by exertion, relieved by rest.

Mostly while at work.

Mostly in men who labour.

Coincide with the amount of exercise, but may be excited by emotion.

In intervals signs of heart disease may be best detected.

Not often accompanied by distinct pain, but there may be attacks of **Angina Pectoris**.

In attack, pulse small and irregular.

Disease of walls of heart, vessels; **Aneurism**.

ANGINA PECTORIS.

the exciting causes of paroxysms, the times and circumstances under which they occur.

the paroxysm, note position and attitude of patient, facial expression, ability to speak or otherwise, state of skin ; **Pulse**, its frequency and characters ; action of heart during attack. Note state of respiration. Examine urine passed after attack. Examine heart and vessels in the intervals of the paroxysms.

s characterized by sudden paroxysms of intense suffering, with the sense of impending death, or a sense of want of air, burning pain in chest, or sense of constriction, pain radiating from the chest down the left arm. In paroxysms there may be profuse sweating, face pale, occasionally flushed, palpitation, subsequent exhaustion.

se may be small and weak, or strong and not frequent.

sation.—Organic disease of heart, walls, or valves. Disease of vessels. See **Signs of Heart Disease**, especially hypertrophy and dilatation. Atheroma of arteries, **Syphilis**, **Aneurism**, over-exertion, **Alcoholism**, **Gout**, **Hysteria**. Reflex exciting causes, *e.g.*, dyspepsia, uterine derangement, mental excitement.

PERICARDITIS.

Precordial pain and tenderness. Dyspnoea, especially in upright posture. Tendency to syncope. **Palpitation.** Pain on swallowing. **Fever.**

Inspection.—A diffused wavy impulse may be seen. In young subjects the precordial region may bulge.

Palpation.—Precordial fremitus may be felt, especially if patient sit or stand up. Apex-beat may be elevated and slightly displaced outwards. Tenderness on upward pressure from epigastrium. Pulse feeble, irregular, intermittent.

Auscultation.—Friction sound not suspended on holding the breath; it may be altered by pressure—"to-and-fro," or only systolic; a brush or hard grating sound. Describe the heart's sounds heard as well as these adventitious sounds.

Percussion.—Tenderness. Enlarged area of cardiac dulness extending as a triangle, apex upwards, to second rib, and even passing to right of sternum. Area of dulness may pass outside apex-beat.

Causation.—**Rheumatism. Bright's Disease. Scarlet Fever. Erysipelas** and other fevers. **Pyæmia.** Cold. Neighbouring abscess or cancer, etc.

PERICARDITIS.

exist with other inflammatory conditions, *e.g.*, **Pneumonia**, **Pleurisy**. If valvular disease coexist there may be orthopnea. The oesophagus is in close relation to the cardium.

m. — Extended wave of impulse may be due to **oedema** or **Retracted Lung** exposing the left auricle.

n. — Fremitus may be absent in recumbent posture, effusion be excessive or purulent. Diffused impulse may be mistaken for cardiac dilatation ; it is weaker than cardiac **Hypertrophy**.

o. — Friction may be inaudible with excessive serous effusion or with pus ; it ceases when adhesion occurs. Motion of a roughened pleura moved by heart may be mistaken for pericarditis. Endocardial murmurs may accompany those exocardial.

p. — Extended area of dulness may suggest hypertrophy, but in pericardial effusion the sounds are indistinct. Extended dulness, apparently cardiac, may be due to elevation of the left lung.

q. — From endocardial murmurs by the friction being increased and being heard as localized and not specially inducted in certain directions. Fremitus may be altered by pressure of the stethoscope on the chest, especially in young subjects. The friction is heard as superficial more grating than an endocardial murmur.

CONGENITAL DEFECTS OF THE HEART.

Among the signs of malformation or congenital defects of the heart and vessels there may be cyanosis, clubbed fingers and toes, low temperature, a general want of development, or some special deformity of the mouth, ears, fingers, etc. Enquire as to the history of the mother's pregnancy.

Make physical examination of the heart; there may be hypertrophy of one or both ventricles in various forms of malformation. **Tricuspid Constriction** may result from foetal endocarditis.

See **Developmental Defects**.

TRICUSPID REGURGITATION.

Systolic murmur at the lower part of the sternum to its right side or near the ensiform cartilage, not conducted to the aorta. Examine for other valvular lesions and **Emphysema**. As a primary disease it is rare, and usually congenital. Examine jugulars for venous pulse, and see if they refill from below when emptied.

CONGENITAL DEFECTS OF THE HEART.

Some conditions are incompatible with life. The conditions most commonly met with are communications between the ventricles through the septum ; these may be accompanied by a systolic bruit heard near the base of the heart, not conducted into the arteries. Cyanosis is not a necessary accompaniment ; the bruit may not be constant. Patent foramen ovale but rarely produces a murmur ; it is often accompanied by a contracted pulmonary orifice, which produces a systolic murmur at the base.

TRICUSPID REGURGITATION.

Usually secondary to mitral disease or emphysema ; the right ventricle is then hypertrophied. At systole regurgitation takes place into the vena cava, causing venous pulse in the neck, the pulsation being perceptible to sight and touch. It may be temporary from over-distension of right ventricle during an attack of **Bronchitis**.

THORACIC ANEURISM.

Physical signs.—Pulsation, when the aneurism points against the chest-wall, most usually about third right rib. The chest-walls may be bulged, the ribs absorbed, and the tumour become prominent. Impulse may be felt with a thrill at same point. The heart may be displaced, but is not usually hypertrophied. The sternum or chest-walls may be heaved up without any prominent tumour. A systolic or double bruit may be heard at seat of impulse, but not necessarily so. Heart-sounds may be heard as distinctly over the aneurism as over the heart itself. The aortic valves may be incompetent. Dulness over chest in an abnormal situation without signs of lung consolidation.

Pressure signs.—Pressure on one lung or bronchus causes dyspnoea on exertion, and loss of respiratory murmur over the lung compressed. **Hæmoptysis**, or ulceration of bronchus or trachea. Dysphagia from pressure on the œsophagus. Constant pain in back. Pressure on sub-clavian artery causing unequal pulses in radials. Pressure on veins causing œdema and enlargement of superficial veins. Irritation of sympathetic nerve in chest causing dilatation of the corresponding Pupil, or its contraction if nerve is paralysed. **Paralysis** of one recurrent laryngeal nerve causing paralysis of the corresponding vocal cord, cough, laryngeal stridor, and metallic-toned voice. (*Note*—the left nerve turns round the arch of aorta, the right round the innominate artery.) Pressure on trachea causes spasmodic cough, often with tracheal respiration, heard over sternum and vertebræ.

THORACIC ANEURISM.

its symptoms.—Pulsation on the surface of the chest, dulness at a point remote from cardiac impulse; hæmoptoe on exertion; stridulous laryngeal breathing from lysis of one vocal cord; pressure signs in thorax; in **the Pectoris; Hæmoptysis.**

1.—Atheroma of aorta. Syphilitic arteritis. Strains injuries. Most common in men and in middle and life. See **Vessels, Disease of.**

2.—from chronic laryngitis; cough loud and paroxysmal with a ringing sound, laryngoscope showing palsy of the vocal cord with no other disease. The pulmonary artery or left auricle may be uncovered by retraction of the left lung, and abnormal pulsation on the surface result, with an enlarged area of dulness. A cancerous aneurism may be pulsatile. Rarely, an empyema may co-exist.

3.—*disease.*—The tendency of an aneurism is to increase in size. If blood is pumped into the sac at a pressure of one ounce to the square inch, that amount of pressure is exerted on each square inch of the aneurism. Pressure may cause absorption of vertebræ, ribs, sternum. The sac may extend into the pleura, lungs, pericardium, œsophagus, externally, etc.

4.—*rupture.*—Sudden or rapidly increasing dyspnoea. Death.

VESSELS, DISEASE OF.

Look for **Anæmia**, **Heart Disease**, **Œdema**, **Bright's Disease**, cutaneous hæmorrhages, loss of elasticity of skin, **Syphilis**, **Gout**, **Alcoholism**, **Epistaxis**.

Arteries.—Examine all the superficial arteries, *e.g.*, radials, brachials, temporals, femorals, dorsales pedis, etc. Feel the condition of the vessels, whether soft or hard, rough upon the surface, rigid, calcareous, locomotor, tortuous, snake-like. **Embolism** may occlude any artery in the limbs: in spleen, causes its enlargement with tenderness; in kidney, temporary albuminuria or hæmaturia; in brain, hemiplegia. Retinal artery may be blocked.

Veins.—Most often diseased in lower extremities; may be enlarged, showing situation of valves. Varicose veins may become hard from occurrence of thrombosis, the clot organizing and becoming hard and cord-like; then ulcer of the skin may result.

Phlebitis may occur during fevers, *e.g.*, enteric, scarlet fever, erysipelas, the vein becoming tender, swollen, hard, cord-like. There may be œdema and subsequent abscess. Look for gout, pressure on the vein, cancer, phthisis, or other cause of great debility. **Anæmia**.

Capillaries.—Often seen dilated over malar bones in persons exposed to the weather; in **Cirrhosis of the Liver**, chronic **Bright's disease**, heart disease, alcoholism.

VESSELS, DISEASE OF.

Often coexists with general degeneration of the tissues of the body, and especially of the kidneys.

Arteries.—Disease may be senile degeneration or due to local injury or strain, atheroma, gout, rheumatism, syphilis, alcoholism. There may result **Aneurism**, aortic incompetency, thrombus, embolism, gangrene, cerebral hæmorrhage. **Arteritis**; if diseased, specially liable to give way when heart is hypertrophied, as with **Granular Kidney**. It is apt to lead to vertigo, hemiplegia. **Embolism** may start from a diseased valve or point of atheromatous artery. Onset of symptoms often sudden; it may obstruct any systemic artery; it often occurs in brain.

Veins.—Staining of the legs in course of veins may result from constantly sitting before the fire. Varicose veins may result from long standing or constipation. Phlegmasia dolens after confinement. Loss of fat from the legs removes their natural support.

Phlebitis, or inflammation of a vein. The clot may break down and lead to pyæmia. It may be detached and carried to the right side of the heart, and plug the pulmonary artery or a branch, causing dyspnoea, hæmoptysis from collateral hyperæmia, syncope from arrest of circulation in the right heart, and sudden death. It may occur deep in a limb.

Capillaries.—Chronic capillary congestion in a limb often seen when the nervous centres are diseased, *e.g.*, paralysis, *idiocy*, *etc.*; hands blue and cold; chilblains.

DISEASES OF THE RESPIRATORY SYSTEM.

CLINICAL REGIONS OF THE CHEST.*

Supra-clavicular.—From outer end of clavicle to trachea.

Clavicular.—Behind inner half of clavicle.

Infra-clavicular.—From clavicle to lower border of third rib and outwards to a vertical line from the acromial angle, which divides the anterior from the lateral regions.

Mammary.—Extends to lower border of sixth rib. The nipple is usually over the fourth rib.

Infra-mammary.—Extends to lower margin of the ribs.

Lateral regions: Axillary.—From apex of axilla to line continuous with lower border of mammary region, and bounded posteriorly by scapula.

Infra-axillary.—Extends down to the margin of the ribs.

Upper and lower scapular regions.—Above and below spine of scapula.

Inter-scapular region.—Between inner edge of scapula and spines of dorsal vertebræ.

Infra-scapular region.—From angle of scapula to margin of ribs.

Upper sternal.—Extends to lower border of third rib.

Lower sternal.—From third rib downwards.

* After Dr. Walshe: "Diseases of the Lungs."

CLINICAL REGIONS OF THE CHEST.

clavicular.—Contains apex of lung; this is usually highest on the right side; also portions of subclavian and axillary arteries, and large veins.

axillary.—Lungs, large arteries.

sternal.—Upper lobe of either lung. Right side, close to sternal border lie the superior vena cava and part of the arch of aorta. Left side, edge of pulmonary artery, the base of the heart being below.

apical.—Right side, middle lobe of lung. Left side, pre-axillary area, sloping outwards and downwards to a point about an inch below and internal to nipple.

hepatic.—Right side, liver dullness, the lung encroaches to a variable extent on full inspiration. Left side, subcostal, and inner portion of left lobe of the liver. Splenic region to sixth rib in lateral region.

apical regions: *Axillary*.—Contains upper lobes of the lungs.

subcostal.—Lower margins of the lungs sloping downwards and backwards. Right side, liver; left side, spleen and stomach.

axillary and lower scapular regions.—Contains lungs.

sternal region.—Lungs, main bronchi, and glands, descending aorta, oesophagus.

apical region.—Lungs down to eleventh rib; liver lies above this on right side. Left side may be partially occupied below by intestine. Aorta descends along the inner boundary.

sternal.—Contains large vessels; transverse portion of arch of aorta. Aortic valves at level of third right rib, pulmonary valves to the left. Bifurcation of aorta at level of second rib.

apical.—Main portion of right ventricle and a small portion of the left resting upon the diaphragm and liver; upper part a small portion of the left lung.

PHYSICAL EXAMINATION OF THE CHEST.

Inspection.—Observe general configuration ; form, especially local or on one side, *e.g.*, bulging or retraction ; observe spine, if straight. Chest movements—thoracic, abdominal. In health, expansive movements are forward and upward. The sternum moves forwards and upwards on inspiration. Specially observe expansive movements in the infra-clavicular regions. In calm breathing, abdominal movements are scarcely observable. Observe position of heart's apex-beat, and the condition of the intercostal spaces.

PATHOLOGICAL CONDITIONS.

Expansion, or bulging, may affect one or both sides ; it may be general over one side or only affect a particular area. Observe intercostal spaces, whether bulged or sunken. Look for position of heart's apex-beat. In all cases carefully compare the corresponding regions on the two sides.

Retraction, or depression, may be general over one or both sides of the chest. It may be localized in one side, as in flattening or retraction in the infra-clavicular region or in the axillary regions. Examine spine ; it may be bent to side contracted, with dropping of that shoulder. Contraction in infra-mammary regions common in infants from *Rickets* and collapse of lung.

Chest movements.—Deficient expansion may be bilateral and general, one-sided or local. There may be a permanent condition of expansion, *e.g.*, **Pleuritic Effusion**, or permanent **Contraction**. In women, respiratory movements are principally thoracic. Movements of diaphragm may be restricted by various conditions of the abdomen, *e.g.*, **Ascites**, ovarian tumour, **Abdominal Tumour**.

Rhythm of the respiratory act.—In health, if the total duration of one movement be taken at 10, inspiratory movement=5, expiratory 4, pause 1.

PHYSICAL EXAMINATION OF THE CHEST.

Inspection.—The general form should be symmetrical on the two sides, and slightly convex in the infra-clavicular regions. Shoulders should be on the same level, and the neck straight. Specially observe movements, and **Signs**

Retraction in the infra-clavicular regions. The two sides of chest should be symmetrical, but, in men, muscular development may cause greater fullness on the right side. There may be expansion, or bulging, or retraction, or altered chest movements. Chest may be deformed from **Rickets**.

PATHOLOGICAL CONDITIONS.

Enlargement, or bulging.—General enlargement of both sides may be due to **Emphysema**. If one-sided from **Pleuritic Effusion** or **pneumothorax**, the heart is then generally displaced. Local bulging may be due to **Aneurism**, mediastinal tumour, encysted empyema; in right infra-axillary region to enlargement or tumour of liver. In children, **Cardiac hypertrophy** may cause local bulging.

Contraction, or depression. implies contraction of the lung corresponding, as from consolidation or pleurisy. It may be general in atrophous emphysema. In infra-clavicular regions it is an important indication of **Phthisis**. Collapse lung may occur from **Laryngeal Disease**, and accompanies "pigeon-breast" in rickets.

Movements.—Movement may be restricted by the pain of pleuritic stitch or by **pleurodynia**; by ossification of the ribs, or by conditions of the lung and pleura. Deficient movement in the infra-clavicular spaces accompanies contraction of the apex. In **Emphysema** vertical movement of the sternum is usually unaccompanied by any forward expansive movement.

Rhythm of the respiratory act.—Duration of expiratory movement may be greater than the inspiratory, *e.g.*, in obstruction to entry of air, in emphysema. Inspiration may be short and abrupt.

PHYSICAL EXAMINATION OF THE CHEST.

Palpation.—Observe movements of the chest, both general and local. Compare the two sides. Determine the intensity of tactile vocal fremitus (T.V.F.) in various situations. A friction fremitus from pleurisy or pericarditis, or from a rhonchus in young subjects, may be detected.

Percussion.—Percuss each region of the chest, and determine the boundaries of the heart and liver, height of apices of lungs in neck. If the percussion note varies from the normal, determine the area of this abnormality, and compare with the same region on the other side.

Hyper-resonant or tympanitic.

Cracked-pot sound.—Jerky and with metallic character.

Amphoric.—Like the sound of flapping the cheeks tensely distended.

AUSCULTATION.*

Note separately inspiration and expiration, their character, relative duration, and whether accompanied by adventitious sounds. Auscultate each region of the chest.

Normal respiration.—Vesicular murmur ; breezy.

Puerile respiration.—Exaggerated in both sounds, increased in intensity, especially the expiratory.

ABNORMAL SOUNDS FROM ALTERED CONDUCTIVITY OF LUNG-TISSUE.

Harsh respiration.—Loss of natural softness and breeziness. Expiration increased in duration and in intensity.

Bronchial respiration.—A higher degree of harsh respiration. Both inspiration and expiration are altered.

Tubular respiration.—Air heard drawn in and puffed back with a metallic character.

Cavernous respiration.—Hollow metallic sound.

* These definitions are mostly quoted from Dr. Walshe, op. cit.

PHYSICAL EXAMINATION OF THE CHEST.

Inspection.—Of great value in detecting local contractions and impairment of movement. T.V.F. (tactile vocal fremitus) increased (usually) over **Solidified Lung** and diminished over a **Pleuritic Effusion**.

Percussion.—In health, the sound is resonant, and resistance is rattle over lung. Sound approaches dulness, and resistance increases with various degrees of consolidation of the lung, or pleuritic effusion. Dulness may be noted on superficial or deep percussion only.

Hyper-resonant or Tympanitic.—Over **Emphysema** or **Pneumothorax**.

Cracked-pot.—Over a vomica; sometimes in young children without disease.

Amphoric.—Vomica. **Pneumothorax**.

AUSCULTATION.

To determine the physical condition of the lungs, and the position of their margins. In health, duration of inspiratory sound to the expiratory is as 3 : 1 (inspiratory movement of chest to expiratory as 5 : 6). Note separately the respiratory murmur and any adventitious sounds.

Respiration.—Normal in children. In adults, frequently limited to a portion of lung doing extra work (supplemental respiration) on account of neighbouring lung-tissue consolidated or compressed.

ADDITIONAL SOUNDS FROM ALTERED CONDUCTIVITY OF LUNG-TISSUE.

Respiration.—In moderate degrees of consolidation and **Emphysema**.

Altered respiration.—Indicates slight condensation of lung substance.

Crackling respiration.—Perfectly developed over hepatised lung pneumonia.

Rhonchus respiration.—Indicates probable cavity from phthisis; dilated bronchus.

AUSCULTATION.

ADVENTITIOUS SOUNDS.

Rhonchi.—Whistling, cooing, bubbling, crackling sounds.

Sonorous rhonchus.—Inspiratory and expiratory usually; sometimes heard without contact with the chest. It is a snoring sound.

Sibilant rhonchus.—Dry sounding; high pitched, sometimes hissing in character; whistling.

Crepitations are crackling râles occurring in successive puffs, all resembling one another. They may occur with inspiration or expiration.

Fine crepitation resembles the sound produced by rubbing hair near the ear; it occurs on inspiration in the first stage of **Pneumonia**.

Pleural friction sound.—Heard only with respiratory movements, except that occasionally a lung, roughened at its margin, is moved by the heart. It may be heard on inspiration and expiration; jerky in character; grating; like a simple brush; or a creak like that of new leather.

COUGH.

Note character and frequency; paroxysmal, *e.g.*, **Whooping Cough**; whether occurring in prolonged attacks; accompanied by **Sputa**.

Causation.—**Bronchitis**; lung disease; **Phthisis**; bronchopneumonia; **Pleurisy**; **Heart Disease**; pressure on air tubes in chest, *e.g.*, **Aneurism**, mediastinal tumour, enlarged bronchial glands. See whooping cough. Reflex causes: examine **Mouth**, fauces, pharynx, **Larynx**.

AUSCULTATION.

ADVENTITIOUS SOUNDS.

may be greatly altered by a cough; they may appear and return, being much less constant than frictions, which they sometimes resemble. They are characteristic of **Bronchitis**, and are frequently so loud to mask all respiratory sounds. The fremitus produced by a rhonchus may commonly be felt in children palpation.

ions may be mistaken for pleuritic friction. Small bubbling crepitations are heard at bases in **Pulmonary oedema**. Scattered crepitations are commonly heard at apices in **Phthisis**. Crepitations are sometimes absent if the patient has coughed and cleared the bronchus leading to the seat of crepitus.

friction sound.—It is more lasting than a rhonchus, and cannot be coughed away. It indicates a roughened pleural surface, but may not be heard in **Pleurisy** on account of **Pleuritic Effusion**, or the hepatization of lung beneath which it is preventing its movement.

COUGH.

Cough is a necessary accompaniment of lung disease, and often is dependent upon lung disease. Prolonged attacks of coughing sometimes cause so much asphyxia that temporary loss of consciousness arises from passive congestion of the brain.

SPUTUM.

Its amount, consistence, whether aërated, colour, mixture of substances, blood, colourless, mixed with blood, streaked with blood, yellowish, white; frothy, mucilaginous-looking, watery, viscid, grumous; mucus, purulent, unmulated, in viscid masses.

HÆMOPTYSIS.

Causation—

Valvular disease of left side of heart (pulmonary).

Valvular disease of right side of heart (bronchial).

Embolism of pulmonary artery from peripheral veins (infarction).

Embolism of bronchial artery from left side of heart.

Blow on chest.

Bronchitis. Plastic bronchitis. Foreign body in trachea.

Blood entering the larynx and coughed up.

Aneurism bursting into bronchus.

Spasmodic **Asthma** (bronchial).

Emphysema. Asphyxia (bronchial).

Scurvy. Hæmorrhagic diathesis.

Renal disease (vessels diseased). **Uræmia** (blood changes).

Degeneration of tissues and vessels (alcoholic).

Phthisis. Cancer of lung.

Pneumonia. Abscess of lung.

Vicarious menstruation attended with amenorrhœa.

After an attack of hæmoptysis there may be signs of blood having run down to base of lungs (crepitations and dulness). It may occur accidentally without organic disease. Hæmorrhage from the throat may be mistaken for hæmoptysis.

SPUTUM.

a frothy water, colourless in early **Phthisis**; later purulent, copious, and (when vomicae have formed) nummulated. Viscid, sticky, golden coloured in **Pneumonia**, and prune-juice colour if mixed with blood. White, aerated, frothy in simple bronchitis. Stinking with gangrene of lung, and in some cases of dilated bronchial tubes.

Dagnosis of HÆMOPTYSIS from HÆMATEMESIS.

<i>Expectorated.</i> —Bright, frothy, may be mixed with mucus. Saline.	Dark, clotted, mixed with food. Acid.
--	--

<i>Character of expectoration.</i> —Coughed expelled without effort; weakness subsequent to expecto- ration. No food expelled.	Vomited mixed with food. Acid. Patient often faints before expectoration.
---	---

<i>Diagnostic symptoms.</i> —Cough, history of Phthisis , previous expectoration of blood with expecto- ration.	Signs of Ulcer or Cancer of Stomach , pain with food, epigastric tenderness, mala- ria. Cirrhosis of Liver .
---	--

<i>Accompanying symptoms.</i> —Subse- quent expectoration of mucus and blood.	Subsequent blood by stool, usually black, tar-like matter.
---	--

Hæmoptysis is mostly due to disease of the lungs or heart. It may also be due to blood changes, *e.g.*, uræmia. Carefully examine heart, lungs, urine. P. = ; T. = ; H. = ; W. = . Enquire as to history of lung disease in patient or his family, also for early deaths in family. General condition of nutrition, etc. Hæmoptysis may apparently be sometimes purely accidental in a lung previously healthy, and blood remaining in the lung may be expectorated at up *phthisical changes*.

DYSPNŒA.

General condition.—Position of the patient, orthopnoea, cyanosis, fulness of the veins, **Œdema**, **Anæmia**. P. = T. = ; R. = . Note any stridulous breathing or signs of **Laryngitis**. Respiratory movements, whether thoracic or abdominal; if accompanied by collapse of the base of the chest or recession of the epigastrium on inspiration. Ability to speak; voice. Character of the dyspnœa, constant or paroxysmal; causing much distress; attended with pain, cough, and expectoration. Increased by exertion or occurring on exertion only (probably cardiac). Examine the lungs, heart, urine. Note condition of circulation, **Pulse**, **Vessels**. Respiratory muscles, if in state of over-action, especially the sterno-mastoids. Act of alae nasi. Fixation of the arms to enable chest muscles to act at greater advantage. General condition of **Nervous System**.

PULMONARY ŒDEMA.

At base of lungs abundant small bubbling râles. T.V.F. may be increased or diminished. On percussion resonance diminished and resistance increased. Dyspnœa.

Note position of patient; signs of **Typhoid** state. Examine urine. Note general condition of patient, specially **Nervous System**.

DYSPNŒA.

Causation—

Structural changes.—**Emphysema**; **Phthisis**; **Pneumonia**; **Bronchitis**. **Œdema** of lungs. **Pleuritic Effusion**, pneumo-thorax, acute pleurisy. Upward pressure of diaphragm from ascites.

Conditions of pulmonary circulation.—Congestion. Heart disease. **Embolism** of pulmonary artery. Clot in heart. Heart failure as when fatty or dilated. Aneurism or mediastinal tumour pressing on trachea or bronchus.

Laryngeal obstruction.—Laryngitis; paralysis of cord; growth upon cord; **œdema** of larynx.

General condition.—**Anæmia**. Fever. **Uræmia**.

Nerve conditions.—**Asthma**. **Hysteria**. Paralysis of nervous centres. **Graves' Disease**. Spasm of respiratory muscles, *e.g.*, from tetanus.

ŒDEMA OF LUNGS.

In the course of pneumonia it may occur in lung tissue adjacent to that inflamed, or in the opposite lung. May attend bronchitis or any lung disease. With pleuritic effusion may attack the other lung. **Uræmia**; **Fevers**; **Passive (cardiac) Congestion** from valvular disease, or degeneration of heart's walls. Frequent in conditions of prostration with dorsal decubitus.

CONTRACTION OF LUNG.

Inspection.—Over portion of lung contracted, thorax contracted; expansion (inspiratory) diminished. Contraction of one side of chest suggests previous **Pleurisy**; of an apex, **Phthisis**. Contraction of left lung may uncover left auricle. Look specially at infra-clavicular regions in adults, and at bases in infants.

Palpation.—Note diminished expansion, general, one-sided, or local. Position of heart; it may be drawn over by a contracting lung. Pulsation of left auricle may be felt if left lung is contracted. T.V.F. may be increased.

Percussion.—Sound may be of impaired resonance from thickening of pleura with lung consolidation. The resistance felt may be increased. Frequently dullness exists from coincident consolidation.

Area of pulmonary resonance above clavicle diminished over a contracted apex.

Auscultation.—Respiratory sounds usually weak and may be abnormal from altered conditions of the lung.

Look for signs of **Consolidation**. **Phthisis**.

SOLIDIFICATION OF LUNG.

Inspection.—Very commonly coincident signs of contraction, especially if the consolidation is at the apex.

Palpation.—T.V.F.* increased. Diminished expansive movement may also be detected. Note area affected, and whether over one or both lungs.

Percussion.—Dulness or various degrees of impaired resonance may be observed over area of solidification ; line of dulness not level, and changing with position of patient as in pleuritic effusion. Note effect of light and deep percussion.

Auscultation.—V.R.† increased. Respiration harsh, bronchial, or tubular ; may be cavernous if there be excavation. Puerile in neighbourhood of consolidation.

Look for signs of **Contraction of Lung ; Phthisis ; Pneumonia.**

* T.V.F. = Tactile vocal fremitus.

† V.R. = Vocal resonance.

Diagnosis of PNEUMONIA from PLEURITIC EFFUSION.

Inspection. — Expansion diminished. No contraction of chest unless lung shrinks from chronic changes.

Palpation. — T.V.F. increased (sometimes diminished), occasionally a pleuritic fremitus felt.

Mensuration. — Rarely any bulging.

Auscultation. — First stage, fine inspiratory crepitant râles, often also pleuritic rub. Second stage, tubular respiration. Rhonchus or scattered râles. V.R. increased. Resolution: Redux loose crepitus, inspiratory and expiratory. Friction sound may return.

Percussion. — Dulness at base, usually following the line of lower lobe downwards and forwards. Increased resistance felt. No change with alteration of position.

Determination of the position of heart and liver. No displacement.

Bulging of side of chest affected, also of the intercostal spaces. As fluid is absorbed, contraction and bending of spine to side affected.

T.V.F. absent below line of dulness; may be increased above. Fremitus in first stage.

Bulging usual. Tracing by cyrtometer.

First stage, pleuritic friction, inspiratory, expiratory, or both. Second stage, effusion. Respiratory murmur absent in axilla, frequently blowing respiration near spine; puerile at apex. V.R. absent or ægophonic. Resolution: Return of respiratory sounds at base. Redux friction.

Line of dulness at base level coming round to the front. Dulness shifting with position of patient. May be tympanic above fluid.

Heart displaced, especially with effusion on left side; liver may be depressed.

Hypodermic syringe may be used to draw off the fluid.

PLEURISY.

iction heard during inspiration, or expiration, or during both periods; it is lost after effusion has occurred, and may return after absorption of fluid or reduction of a pneumonia.

iction fremitus often felt.

iction of pleural surface usually attended with pain, causing patient to hold his breath; he lies on side affected.

the pleurisy be secondary to lung disease, *e.g.*, phthisis, symptoms will be those of the lung disease. Pyrexia in pleurisy, lower than the inflammatory fever of pneumonia.

e signs of **Pleuritic Effusion**. P. = ; T. = ; R. = .

uation, see same in **Pneumonia**.

euritic effusion is always albuminous; occasionally it coagulates from presence of fibrin.

EMPHYEMA.

ten not distinguishable from serous effusions before tapping.

It is most common in young subjects, debilitated or very strumous. Also when effusion is very chronic. Temperature not necessarily high. More displacement of chest walls and viscera than with serous effusion.

mperature often elevated, but not necessarily so. It may point under the skin in front of chest, laterally, or behind. May occur in Septicæmia, **Pyæmia**, **Erysipelas**, Scarlet Fever, **Puerperal Fever**. It may discharge by bronchus.

HYDROTHORAX.

ssive dropsical effusion without pleurisy. May occur from **Passive (cardiac) Congestion**, **Bright's Disease**, etc. It is usually *double* and unaccompanied by fever.

PHTHISIS.

Physical signs.—**Signs of Consolidation and Contraction of Apex of lung.** Carefully inspect movement in infra-clavicular fossa on each side, examining for signs of contraction of the apex. Palpate, noting if T.V.F. is increased. In some cases the left auricle is uncovered from contraction of the left lung. Percussion gives a dull or wooden note ; note the sound of light or deep percussion. The resistance increased over consolidated lung. It may be amphoric over a vomica, but still the resistance is augmented. Auscultation shows V.R. increased, respiration harsh or bronchial, with adventitious sounds, scattered râles, crepitation.

Digestion.—Dyspepsia often troublesome. Diarrhoea may be due to tubercular **Ulceration of Intestines.**

Circulation.—Note force and strength of heart's action ; it often partakes in the general wasting.

Nervous system.—General condition. Sleep.

Urine.—Albuminuria may be present. **Diabetes** is a frequent cause of phthisis.

PHTHISIS.

ough, with expectoration and **Hæmoptysis**, debility and weakness, emaciation. Sweatings especially at night. Flushings; fever; dyspnœa on exertion. Anæmia, and in women amenorrhœa. Muscular irritability often marked. In pregnant women phthisis is often temporarily arrested, becoming active after parturition. In advanced cases there may be œdema of the legs.

Causes.—Inheritance; history of consumption or **Scrofulous** disease in family; give ages of any members of the family who died. Hygienic conditions, locality of residence with regard to climate and dampness, dusty trades. Exposure to cold. Sequent to acute lung diseases, or hæmoptysis. A common termination in diabetes mellitus.

Possibly it is communicated from the diseased to persons predisposed.

Complications.—**Laryngitis**, bronchitis, pneumonia, hæmoptysis, **Pleurisy**, empyema, pneumo-thorax.

Failure of heart's action; thrombosis; bed-sores. Diarrhœa or **Melæna** from tubercular ulceration of intestines; fistula; **Peritonitis**; **Liver large**, fatty or amyloid; **Albuminuria**; **General Miliary Tuberculosis**; **Edema** of legs.

Signs of a cavity (**vomica**). Percussion, giving a metallic cracked-pot sound on auscultation; respiration blowing, tubular, cavernous, with moist râles at apex. Pectoriloquy.

PNEUMONIA.

Physical signs.—Signs of **Consolidation** over hepatized lung.

Earliest sign, fine inspiratory crepitation resembling the rustling of hair; there may be also a pleuritic friction. In hepatization, dulness along outline of the lobe solidified; if at base, sloping downwards and forwards. T.V.F. usually increased. Respiration tubular and often accompanied by rhonchus and râles. Voice broncho-phonic. On resolution respiration becomes less tubular; crepitation, loose inspiratory and expiratory (=redux crepitation). A return of the friction rub may be heard.

Digestion.—Tongue furred; thirst; anorexia. There may be vomiting, diarrhoea, **Jaundice**.

Circulation.—Note force of impulse and first sound of heart. Characters of pulse.

Nervous system.—General condition of **Nervous System**; sleep, restlessness, **Delirium**.

Urine.—Scanty, with excess of lithates; chlorides deficient. May be albuminous.

Complications.—Pulmonary œdema; collateral congestion. Bronchitis; high fever; failure of heart, pulse becoming weak and soft. **Jaundice**; **Delirium**; **Albuminuria**; **Typhoid State**.

PNEUMONIA.

In acute cases onset sudden with rigor, fever, quick breathing. Pleuritic pain and dyspnoea usually subside with the pyrexia, and coincidently with the signs of hepatization. Cough; expectoration viscid, golden colour, occasionally streaked with blood; it may be accompanied by aerated, frothy bronchial sputum. Note date of disease; P. = ; T. = ; R. = . Pleuritic pain may return during resolution. Symptoms usually subside by crisis; dyspnoea, fever, distress passing off suddenly, leaving lung hepatized and patient prostrated.

Classes of Pneumonia.—Acute sthenic as above described: usually at base. Asthenic with adynamic symptoms: less sudden onset and no marked crisis; less distinctly marked signs of solidification; much tendency to bronchitis and pulmonary oedema, patient tending to the **Typhoid State**. It may end in **Gangrene of Lung**.

Pneumonia of the apex. Frequently accompanied by grave nervous disturbance, and long convalescence or subsequent phthisis.

Causation.—Exposure to cold. A complication of fevers. Secondary to chronic disease, *e.g.*, of lungs or kidneys; rheumatism; injury; adjacent inflammation or disease, *e.g.*, pneumonia, cancer, tubercle.

EMPHYSEMA.

Physical signs.—Chest may be large or small ; expansion is markedly diminished, and such movement as there may be is usually vertical without forward expansion. Heart's impulse more or less encroached upon, and marked by lung covering it, but it may be felt as somewhat diffused. General hyper-resonance on percussion. Absolute dullness over heart may be wanting with an extended area of relative dullness. On auscultation, expiratory sound much prolonged ; feeble and toneless, harsh, often accompanied by rhonchi and sibili. Liver may be depressed.

Circulation.—Pulse feeble ; right ventricle dilated ; heart may be hypertrophied. **Passive Venous Congestion.**

Urine.—May be scanty and albuminous. **Chronic Granular Kidney** not uncommonly accompanies emphysema.

Causation.—Vicarious dilatation, *e.g.*, adjacent to pulmonary collapse or consolidation, or cells obstructed by bronchitis, Paroxysmal cough ; laborious work ; **Hooping Cough** ; heart disease ; **Alcohol** ; **Gout** leading to ill-nourished condition of lungs. Senile changes.

EMPHYSEMA.

lose their elasticity, much aërating surface is lost, and pulmonary capillaries destroyed, thus obstructing the flow from the right ventricle. Passive venous congestion results. The difficulty of expanding lungs with diminished elasticity throws respiratory muscles into strong action, the sterno-mastoids are often hypertrophied. Patient may emaciate or grow fat; in neither case is nutrition good. Usually chronic winter cough and liability to acute bronchitis.

Signs and accompaniments.—Heart: right **Ventricle** enlarged and hypertrophied; veins large; cyanosis; **Tricuspid Regurgitation**. **Œdema** of feet. **Bronchitis** due to passive congestion of bronchial veins, which empty their contents into the right heart. Dyspnoea on exertion. **Albuminuria** may be from coexisting Bright's disease, and is usually constant; if albumen be due to renal congestion it may pass off with other signs of congestion, the symptoms lessening and the quantity of urine increasing. The capillaries of cheeks often enlarge.

BRONCHITIS.

Physical signs.—If bronchitis is secondary to, or complicates other disease of lungs, the signs will be partly those of that other diseased condition.

Auscultation.—Often gives negative results, especially in chronic winter bronchitis. Rhonchi; sibili; râles.

Palpation.—Rhonchi are sometimes felt by the hand especially in the elastic chests of infants. Palpate heart.

Percussion.—No change from the normal, or temporary tonelessness in parts.

Inspection.—Observe chest movements; collapse of chest at apices, or in hypochondriac regions. **Dyspnoea.**

Urine may be albuminous, a similar cause producing Bright's disease and bronchitis. See **Albuminuria.**

Inquire for—P. = ; T. = ; R. = ; W. = . Signs of **Consolidation of Lungs**; signs of contraction. **Cough**; **Expectoration**; **Hæmoptysis.**

ASTHMA.

Respiration.—Percussion unaltered during paroxysms; shrill whistling sibili. Examine lungs during paroxysms and during intervals. The paroxysms, note their frequency and duration, exciting and predisposing causes. **Cough**; **expectoration.**

Causation.—Hereditary tendency to neurosis. Reflex causes, uterine, constipation. Tubercular diathesis; **Emphysema**; **Heart Disease**; **Uræmia.** May occur in Bright's disease without other signs of anæmia.

BRONCHITIS.

This condition may be acute or chronic ; primary or secondary to other disease, *e.g.*, **Emphysema**, **Phthisis**, **Pneumonia**, etc.

It is characterized by cough, with expectoration usually frothy and watery, sometimes viscid or purulent ; dyspnœa. Fever usually slight, but high in children. Post-sternal pain and tenderness, increased on coughing ; skin over sternum sometimes sore.

Causation.—Exposure to cold ; fevers ; bronchitis secondary to chronic lung conditions ; phthisis ; emphysema secondary to acute conditions ; pneumonia ; pleurisy.

Secondary to heart disease ; **Rickets** ; mechanical irritants.

Course of disease if towards fatal termination.—Inability to expectorate. Rapid respiration. Pulse becoming weak, compressible, irregular ; heart distended on the right side ; veins prominent ; cyanosis. **Edema** of legs increasing. Râles all over lungs. Sleeplessness. Tendency to **Coma** and the **Typhoid State**.

Urine scanty and albuminous. Temperature falling. In children collapse of chest at bases with infalling of epigastrium.

ASTHMA.

An affection characterized by paroxysms of dyspnœa.

Paroxysms.—Orthopnœa ; respiratory muscles, ordinary and extraordinary, at work. Chest fully dilated and respiratory movement almost nil. Sense of want of air. Voice weak or lost. Onset of paroxysm sudden, subsidence rapid ; they frequently occur at night. They may be preceded by drowsiness and a sense of fatigue.

LARYNX, DISEASE OF.

Acute conditions.—**Diphtheria**, croup, catarrh, œdematous, stridor.

Chronic conditions.—Laryngitis: syphilitic, strumous, catarrhal. **Hysteria**. Palsy of vocal cords, or one cord.

General condition.—P. = ; T. = ; R. = ; W. = ; signs of nutrition ; signs of struma or **Syphilis** ; **Phthisis**.

Digestion.—Examine mouth and fauces, using laryngoscope.

Circulation.—Examine heart as to strength and dilatation ; right side ; venous fulness ; strength and regular pulse. Seek for signs of **Aneurism**.

Respiration.—Signs of laryngeal disease ; laryngoscopic examination ; movements of cords. Bronchitis, œdema of larynx, pneumonia, etc. Chest movements. Look for **Ph**.

Nervous system.—Signs of **Convulsions**, thumb turned inward ; chronic spasm of muscles ; hysteria. Palsy of one cord usually from pressure on recurrent nerve. See **Ane**.

Urine often albuminous in diphtheria ; there may be coincident acute Bright's disease.

LARYNX, DISEASE OF.

Signs of Laryngeal disease.—Voice husky or lost ; stridulous inspiration, aphonia, cough, dyspnoea, cyanosis. Dilation of right side of heart, and other signs of obstruction to the entrance of air, *e.g.*, infalling of supra-sternal notch, supra-clavicular spaces, and epigastrium, and in young children collapse of the hypochondriac regions. Tracheotomy may be required when this obstruction is extreme ; in such a case observe the condition of the heart, pulse, and circulation before and after operation.

Laryngismus Stridulus.—Mostly in children ; spasmodic crowing sound on inspiration, child being well in intervals. Frequent during dentition, in **Rickets**, and associated with general **Convulsions**.

Edema may occur during **Bright's Disease**, **Erysipelas**, etc., acute catarrh from cold, or with onset of **Measles**. In **Hysterical Aphonia** cords are seen healthy but motionless ; pharynx often very anæsthetic.

Functional Aphonia.

DISEASES OF THE DIGESTIVE SYSTEM.

SIGNS OF DIGESTIVE FUNCTIONS.

Appetite.—Good, bad, indifferent, altogether lost. Frequency of recurrence, capricious and fanciful; variable, excessive, voracious. Nausea. **Vomiting**.

Fulness or pain after food. Enquire how soon after food; its character and duration; whether pain is relieved by vomiting. Whether pain without food. Flatulence and eructations.

Eructations. Heartburn. Water-brash. Pyrosis.

State of Bowels.—Regular, constipated, relaxed, with or without pain. **Diarrhœa**; frequency of action. If disturbed see and describe the motions—solid, liquid, light, clay-coloured, dark, black; hard scybala, flattened or tape-like, well formed, with blood, pus, etc.

INTESTINAL WORMS.

Tænia mediocanellata—beef tapeworm.—The head is at the narrow portion of the worm; it has four sucking discs, but is unarmed.

Tænia solium.—Less common in England. Pork tapeworm; four suckers and an armed head. *T. Bothriocephalus latus*. They live in small and large intestines. Thread worms—*Oxyuridis*—live chiefly in rectum; common in children.

Lumbricus, round worm. *Ascaris lumbricoidis* lives in upper intestine, and may be vomited.

SIGNS OF DIGESTIVE FUNCTIONS.

etite increases with thirst in **Diabetes**. Anorexia (loss of appetite) and thirst in **Fever**. In children often variable, especially in nervous cases; they often drink much in health, when urine is scanty with high sp. gr. Appetite is often lost in functional disturbance of the nervous system, *e.g.*, over-work, loss of sleep. In **Hysteria** and insanity the appetite may be greatly perverted; so also during pregnancy.

of Bowels.—Constipation may result from **Plumbism**, senile atrophy of bowels, inactive habits of life, ill-arranged diet.

axation or looseness from **Dysentery**, **Ulceration of Bowels**, or other organic condition. In infants from ill-feeding or summer heat.

EXAMINATION OF THE MOUTH AND THROAT.

On obtaining a good view of all parts of the mouth, see—tongue ; hard and soft palate, with uvula ; pillars of the fauces, anterior and posterior ; tonsils ; pharynx ; the buccal cavity ; cheeks and lips, mucous membrane ; gums ; teeth.

Tongue.—Mucous membrane and condition of muscle. Indented at edges by the teeth ; flabby ; clean or coated with fur ; white, yellow, dirty, dry, or moist. Enlarged papilla at tip projecting through fur. How protruded ; straight or deviating to one side, kept well out and steady, or a jerked, tremulous, distinct muscular tremor.

Palate and Uvula.—High arched roof, cleft. Ulceration, destruction of soft palate, adhesions. Movements of soft palate and fauces.

Tonsils.—Enlarged, one or both. Smooth, pale or congested ; with large follicles. Ulcers superficial or deep, if symmetrical. Exudation on surface.

Pharynx.—Mucous membrane and movements. Look for ulcers or old scars and adhesions. Thrush in children ; exudation in diphtheria.

Teeth.—Look for tender teeth ; those subjects of caries ; see if wisdom teeth be cut. Note condition as to dentition in infants.

Gums.—Whether of normal substance or shrunken ; condition of mucous membrane.

EXAMINATION OF THE MOUTH AND THROAT.

It is necessary to obtain a good light in the pharynx ; hence it is often convenient to use a lamp and the frontal reflector of laryngoscope. There may be signs of local or general disturbance.

Tongue.—Flabby and coated in dyspepsia ; often red with **Gastric Ulcers** and **Cerebral Vomiting**. Protruded to one side in **Hemiplegia**. Tremulous in **Alcoholism**, excessive smoking, **General Paralysis of the Insane**. Jerked and twitching in **Chorea**. Ulceration from local irritation. Ulcer of frænum in hooping cough. **Syphilis**. Epithelioma.

Palate and Uvula.—Palate high, arched, flat, cleft. Ulceration, scars, adhesion from scrofulous disease or **Syphilis**. Uvula commonly elongated. Movements of palate and uvula affected in palsy of Nerve VII.

Tonsils.—Chronic enlargement in rickets, often with deafness. Ulcers symmetrical in secondary syphilis. See Quinsy, **Diphtheria**, **Syphilis**.

Pharynx.—Scars from syphilis or strumous ulceration. Paralysis from diphtheria. Epithelioma. Post-pharyngeal abscess from spinal caries.

Teeth.—Upper central incisors (of second dentition) may be dwarfed, with atrophy of the middle lobe in **Inherited Syphilis**. Much ground in gouty people and children who suffer from **Headaches**.

Gums.—Blue line in **Plumbism**. Spongy in mercurialism. Swollen and bleeding in scurvy. Covered with sordes in *fever*

DIARRHŒA.

Note mode of onset and duration ; if attended with painful griping ; **Melæna** ; tenesmus (frequent desire to evacuate the bowels, but without effect). Whether acute with paroxysmal griping, melæna, collapse, as in cholera.

Motions passed.—Relaxed, liquid, pea-soup-like ; containing bile or not ; scybala, shreds of mucous membrane, undigested food, worms.

VOMITING.

Note the frequency of vomiting ; whether it occurs only after food ; whether giving relief to symptoms ; if affected by position. State of tongue and bowels. Examine abdomen for tenderness ; signs of disease of stomach. See general condition of **Nervous System**. Signs of **Brain Disease**. Examine urine.

Matters vomited.—Food unchanged ; bile-stained fluid ; clear acid fluid ; yeast-like matter containing sarcinæ seen on microscopical examination ; blood (*hæmatemesia*) ; dark coffee-grounds-like matter, altered blood ; lumbrici. Faecal matters may be thrown up in obstruction of bowels low down.

DIARRHŒA.

be indicative of local disease or general disturbance.

causes.—Disease of the bowels; **Tubercular Ulceration**; **Amyloid Disease**; stricture of bowel, rectum; scybala; enteric fever; **Dysentery**; cholera; erysipelas, etc.; **Bright's Disease**; ill-feeding; **Alcoholism**; exposure to heat and cold; poisoning; **Rickets**; nervous disturbance; **Graves' Disease**. Previous constipation.

VOMITING.

may indicate local or general disturbance or brain disease. See **Cerebral Vomiting**.

causes.—

Stomach.—Gastritis; dilatation of stomach; catarrh secondary to **Cirrhosis of the Liver**; irritating food; **Alcoholism**; poisons; **Cancer**; **Gastric Ulcer**; constriction of pylorus or duodenum.

Reflex causes.—Pregnancy; ovarian disease; uterine disturbance; dysmenorrhœa; dentition; intestinal worms; **Gall-stones**; **Renal Calculus**; **Addison's Disease**; liver disease, cancer, abscess, etc.; disturbance of special senses, glaucoma, ear disease. Attendant on paroxysms of whooping cough.

Brain disease.—**Headache**; **Hysteria**. See **Cerebral Vomiting**.

Blood conditions.—Fever; malaria; **Bright's Disease**; **Obstruction of Bowels**; **Peritonitis**.

ACUTE ABDOMINAL PAIN.

Enquire as to digestive functions ; previous attacks of **Biliary Colic**, **Renal Colic**, gastric ulcer.

Examine mouth and tongue for indications of **Gastric Ulcer**, poisoning ; and gums for blue lead line.

Palpate and examine abdomen ; note if tender and tympanitic ; position of the patient, whether still and prostrate or moving about. In females look for signs of pregnancy, uterine action, or hæmorrhage.

Examine heart, pulse, skin, pupils, urine. T. = . Note if much collapsed ; whether able to speak ; whether vomiting.

DYSPHAGIA.

General Condition.—**Anæmia**. Signs of **Cancer**. General condition of **Nervous System**. Syphilis. Senile degeneration.

Digestion.—Examine mouth and throat for ulceration, scars, etc. Auscultate spine while patient drinks, looking for gurgling at one point. Pass œsophageal bougie.

Vascular system.—Signs of disease of vessels or aneurism.

HÆMATEMESIS.

Enquire as to the general signs of the **Digestive Functions**, previous vomiting, pain, tenderness, etc. See causes of **Vomiting**. Examine the matters vomited and the motions as to hæmorrhage, etc. Examine the abdomen generally. Look for disease of stomach and liver. Examine lungs and heart to determine absence of causes of hæmoptysis. See diagnosis of **Hæmoptysis** from **Hæmatemesis**. Urine. **Anæmia**. Amenorrhœa.

ACUTE ABDOMINAL PAIN.

Causation.—Rupture of hollow viscera, stomach, intestine, bladder. **Renal or Biliary Calculus.** Irritant poison. Over-feeding. Colic, simple or from gout. **Plumbism**, often relieved by pressure. Rupture of abdominal **Aneurism**, abscess, **Hydatid**, **Peritonitis**. Perihepatitis. Ulceration of bowels with peritonitis or perforation. **Tubercular Ulceration.** Acute disease, *e.g.*, cholera. In females during pregnancy, concealed accidental hæmorrhage. Pain and tenderness suggest enteritis or peritonitis, rather than colic, the latter being often relieved by pressure.

DYSPHAGIA.

Causation.—Tonsillitis; syphilitic ulcerations; disease of larynx; cancer of œsophagus or of cardiac end of the stomach; thoracic tumour; abscess, post-pharyngeal or mediastinal; **Aneurism**; traumatic injury or action of caustics; ulcer of stomach at cardiac end; **Diphtheritic Paralysis**; **Bulbar Paralysis**.

In **General Paralysis of the Insane** there is much tendency to choking. **Hysterical dysphagia.**

HÆMATEMESIS.

Causation.—**Gastric Ulcer.** Corrosive poisons. **Cancer of Stomach.** Continued vomiting (reflex). Acute gastric catarrh. **Lardaceous Disease.** Pyloric ulcer. **Bright's Disease**; uræmia. **Passive Congestion** of stomach. Scurvy. **Cirrhosis of Liver.** Vicarious menstruation. Blood swallowed and vomited. Patient often faints from hæmorrhage, previous to the discharge of the blood from the mouth.

MELÆNA.

General condition.—State of nutrition. **Anæmia** or cachexia

Digestion.—Examine motions; presence of abdominal pain signs of gastric disease, pain on defæcation, etc.

Enquire for signs of **Cancer**; history of malaria; previ diarrhœa; signs of stricture of bowels.

Examine abdomen; if necessary examine rectum; urine; luns as to signs of tubercular disease.

OBSTRUCTION OF THE BOWELS.

General condition.—Position of patient; pain; abdominal tenderness; signs of collapse. T. = . Note what bowels last acted.

Digestion.—Habitual condition of bowels, regular, costive, relaxed. Previous signs of disease, e.g., **Melæna**, **Vomiting**. State of tongue. See and describe the motions passed.

Examine abdomen, especially the abdominal rings, and femoral rings for hernia. Note fulness, tenderness, local swelling or tumour; an elongated tumour from **Intussusception**. Signs of **Peritonitis**. Track out colon, if distended by palpation and percussion. Note if any signs of contraction at any point.

Examine rectum with the finger, or give enema noting what quantity of fluid can be retained; pass the lubricated tube. Sometimes the whole hand is introduced into the rectum.

Examine per vaginam.—Signs of pregnancy. See general signs of **Cancer**.

MELÆNA.

May be caused by all the causes of **Hæmatemesis**, the blood passing from the stomach to the intestines. **Cirrhosis of Liver**, or other obstruction to portal circulation. **Ulceration of Bowels**, tubercular. **Gastric Ulcer**. Cancer of bowels. **Enteric Fever**. **Dysentery**. **Intussusception**. Pelvic hæmatocele or abscess. Piles may cause bleeding from the anus. Villous growth in rectum. **Bright's Disease**.

OBSTRUCTION OF THE BOWELS.

Causation.—I. *Compression*.—Cancer or inflammatory mass involving intestine; **Abdominal Tumour**; pregnant uterus; ovarian tumour; pelvic tumour; uterine, ovarian, cellulitis; retroverted uterus.

II. *Changes in wall of gut*.—Cicatrization of intestinal ulcers, dysenteric, tubercular; congenital deformity of rectum, etc.; **Cancer**; epithelioma and syphilitic disease of rectum.

III. *Strangulation*.—Generally in small intestine, hernia; constriction from mesentery of portion of intestine drawn into a hernial sac; or from bowels, due to peritonitis.

IV. *Plugging*.—Undigested substances, fruit stones and seeds, hardened fæces, masses of worms.

Intussusception; volvulus.

Symptoms vary according to the position of the obstruction, its degree, its cause, the complications. If in small intestines there may be no marked and characteristic symptoms.

The motions may be pipe-like or not formed. Formed motions may be produced by fæces passing the stricture and being moulded in rectum. Constipation; flatulence. If in rectum, pain and straining on defæcation.

GASTRIC ULCER.

Digestion.—Pain immediately after food, relieved only by vomiting. Water-brash vomiting. **Hæmatemesis**. **Melæna**. Inability to take solid food. Localized tenderness at epigastrium; no tumour felt. Bowels usually confined; examine the motions. Tongue usually red.

Note.—General condition; position of patient; state of nutrition; signs of **Anæmia**. General condition of abdomen. Signs of **Hysteria**. General condition of the **Nervous System**. W. = . Urine.

TYPHLOITIS.

Local examination of right iliac fossa. Vaginal or rectal examination to determine absence of pelvic cellulitis.

Causation.—Hardened fæces; undigested food; dysentery. Local concretion in appendix. Cherry stone, or fish bone, etc.

GASTRIC ULCER.

Pain may be less if the ulcer is on the lesser curvature of stomach. In long-lasting cases, some thickening of walls of stomach may be felt, or stricture of the pylorus may result. Usually there is emaciation, anæmia, or cachectic appearance. Menstruation absent or disturbed. Recovery may occur for a while with tendency to relapse of the symptoms, or perforation and **Peritonitis**, vomiting, hæmorrhage, exhaustion.

Causation.—Most common in females; specially accompanies disordered menstruation; may result from action of caustics.

Complications.—Fistulous communication with external surface or with other parts of intestines.

Pyæmia.

TYPHLITIS.

Abdominal pain; local signs of inflammation in the right iliac fossa, pain, tenderness, swelling. Local peritonitis with infiltration of the cellular tissue; it may suppurate. Constitutional disturbance with fever may be considerable if the bowel is involved; less acute if only around the bowel. Perforation of bowel may follow. Usually pain and difficulty in moving right leg.

ABDOMINAL CANCER.

General condition.—Note state of nutrition. W. = .
Emaciation. Pain in back, exhaustion, and cachexia,
with the general signs of **Cancer**.

Digestion.—Signs of digestive functions. Vomiting, hæmorrhage, acid secretions, stomach pain.

Examine abdomen.—Clear out bowels with purgatives or enemata ; empty bladder. Palpate and percuss to detect any abdominal tumour. Note any signs of **Obstructed Bowels ; Peritonitis ; Ascites**. Examine rectum and per vaginam if necessary.

Cancer of Stomach.—A mass may be felt in epigastrium, or an increased resistance, often most distinct along greater curve of stomach. A rounded and movable mass may be felt over the pylorus.

Cancer of Intestines.—A mass may be felt on palpation ; it may be movable. Clear out bowels. Inspect and describe the motions, whether full-sized or flattened and small. **Melæna**. If there is obstruction or arterial hæmorrhage, examine rectum with finger. Look for piles.

ABDOMINAL CANCER.

7 affect stomach, intestines, peritoneum, mesenteric glands, liver, kidneys, spleen, uterus. Secondary deposits in the liver are common. See **Abdominal Tumour**.

Cancer of Stomach.—General signs of cancer. Pain in region of stomach, a very varying symptom. Vomiting acid frothy matter, often with sarcinæ; there may be arterial **Hæmatemesis** or coffee-ground-like matter. Excessive acid secretion. Usually it is primary. Secondary deposits may occur in the liver; it may creep on to pylorus and involve gall-duct. **Jaundice**. A mass may thicken the pylorus causing a tumour that can be felt there, and stricture with vomiting late after food. Scirrhus of stomach may run its course through many years. With a mass that can be felt, patient may still gain weight.

Cancer of Intestines.—Usually primary; most common in the sigmoid flexure, cæcum, and rectum. Abdominal pain. Tendency to annular contraction, causing **Obstruction**. May be mistaken for fæcal accumulations.

2.—Ulceration of rectum may be from epithelioma or **Syphilis**.

ULCERATION OF BOWELS.

Typhoid. Cancer. Epithelioma at anus. Syphilis. Ulceration from gall-stones, scybala, intussusception, etc.

Dysentery.—Note state of nutrition; **Anæmia.** P. = ; T. = ; R. = ; W. = .

Digestion.—Appetite. Abdominal pain or tenderness. Evacuations: colour, consistence, smell, bile, mucus, or sloughs.

Liver.—Size, absence of tenderness, jaundice, etc.

Complications and Sequelæ.—Chronic dysentery. Hæmorrhage from bowel. Abscess of liver.

Tubercular Ulceration. — *General condition.* — Emaciation; excessive sweating.

Digestion.—Appetite. Abdominal condition; fulness, tenderness, pain, general or localized. Bowels relaxed; may be acting with pain and **Melæna.** Enquire for fistula in ano.

Respiration.—Examine lungs, and look for signs of **Phthisis.**

ULCERATION OF BOWELS.

ysentery.—A disease more common in the tropics than here.

Caused by malaria, scorbutus, bad water, salt food, etc. It may occur in an acute or chronic form. It is febrile, characterized by tenesmus with the passage of mucus without fæcal matter or bile, sloughs may be passed with blood. These symptoms depend upon inflammation of the colon with exudation; it may extend to the small intestine.

tubercular Ulceration.—Common in cases of phthisis and other strumous affections. Abdominal pain, diarrhoea, and melæna may result. The tubercular ulcers in the bowels are transverse; they may heal up, leading to scars, which may cause stricture of the bowels. Ulcers occur mostly in the lower part of the ileum and cæcum.

Complications.—Peritonitis. Ascites. Perforation of bowels.

Acute Miliary Tuberculosis.

ABDOMINAL TUMOURS.

General condition.—State of nutrition. W. = . Signs of **Cancer** or **Scrofulosis**. Abdominal pain, tenderness, vomiting, condition of bowels, signs of **Obstruction**. Digestive functions. Look for **Ascites**, **Peritonitis**, **Abdominal Cancer**, œdema. Urine.

Examination of abdomen.—Palpate and percuss abdomen ; thus endeavour to detect any tumour present. Define its position with regard to the anatomical regions ; determine its boundaries and connections ; particularly note if distinct from liver and pelvic organs. Map out liver and spleen, showing them of normal size. Note physical conditions of tumour, its size, if smooth, rounded, lobulated, hard, impressible, doughy, fluctuating. If movable or moving with respiratory movements. Measure the abdomen, girth at base of chest and at the umbilicus, vertical measurements from umbilicus to pelvis, and umbilicus to xiphoid cartilage. In the normal the umbilicus is about an inch nearer to the pubes than to the sternum. Note pain or tenderness. Empty bowels and bladder.

History.—Commencing on one side ; enlarging from below upwards ; enlarging of the abdomen uniformly ; with pain and fever or not. Did symptoms commence at a menstrual period ?

Percuss, palpate liver, define and mark on skin the vertical and other dimensions.

ABDOMINAL TUMOURS.

Ovarian.—Globular, movable, fluctuating; usually situated more to one flank than the other. Springing from the pelvis and may be felt there. Usually dulness in centre of abdomen with resonance in the flanks. If very large may be mistaken for ascites. See diagnosis of **Ovarian Tumour** from **Ascites**. It may be accompanied by ascites. Dulness over an ovarian tumour shows that no intestines are in front of it; so also with a pregnant uterus.

Kidney.—Colon usually passes in front of tumour however large it becomes; this may be indicated by partial and varying resonance over it. There may be mixed resonance and dulness, varying on different occasions. Tumour may be felt in the flank, usually between false ribs and ilium; a tumour in this region may be renal, peri-nephritic, faecal in colon. Abscess; cancer; hydro-nephrosis; blood-tumour. The outline is rounded or lobulated (cystic tumour), not easily defined. Absence of fluctuation.

Liver.—See **Large Livers**. A hepatic tumour descends on inspiration. Gall-bladder; **Hydatid**; **Cancer**.

Abdominal Pulsation common in hysterical women and dyspeptic subjects. Throbbing of abdominal aorta also common in emaciation.

FLUID IN PERITONEUM.

Physical signs.—Enlargement of abdomen. In dorsal position, dulness on percussion over the fluid, which gravitates into the flanks leaving central region clear; on the dependent side of abdomen, a distended colon may give a tympanitic note, but on palpation in this flank the weight of the fluid is felt, line of dulness shifting with position. Thrill transmitted on filling abdomen; fluctuation. When placed on hands and knees, fluid will gravitate to the umbilicus. Clear out bowels; empty bladder; examine per vaginam.

Symptoms.—Dyspnoea and thoracic breathing. Pressure on renal veins may cause scanty urine and **Albuminuria**. Pressure on iliac veins causing oedema of legs. Superficial abdominal veins enlarged.

Causation.—**Cirrhosis of Liver**. Cardiac disease. Disease of peritoneum, tubercle, cancer, **Peritonitis**. Exposure to cold. Ovarian or other abdominal tumour.

Conditions simulating Ascites.—Ovarian cyst. Hydatid cystic kidney. Pregnant uterus. Distended urinary bladder. Fluid in intestines. See **Abdominal Tumours**.

ABDOMINAL TUMOURS.

en.—Feel for the notch towards anterior margin. Usually firm, flat superficial under abdominal walls without intestine in front. Stretching from left hypochondrium. Surface may be lobulated ; it may be tender and movable.

creas.—Has been stated to be frequently the seat of cancer. Examine fæces for fat ; shake up with ether.

sation.—Hypertrophy ; chronic congestion from cardiac or liver disease ; **Ague** ; **Amyloid Disease** ; cancer. Examine blood for leucocythæmia. In children **Rickets**. **Syphilis**. Large sometimes in fevers, specially enteric. Frequent seat of **Embolism**.

ominal Aneurism.—A tumour pulsating and laterally expansile, with a thrill and systolic bruit often also heard over spine. Pain. No necessary dyspeptic symptoms. Pressure signs less common than in thorax ; it may press on vena cava, or cause erosion of vertebræ, producing great pain. It does not fall forward when patient is in knee-elbow position. A tumour lying on the aorta may receive a communicated impulse. The pulsating aorta without disease may often be felt in nervous or dyspeptic patients, especially in females if emaciated. See **Aneurism**.

Diagnosis of
OVARIAN TUMOUR from ASCITES.

Palpation.—Definite margins may be felt. Usually situate more in one side of abdomen than in the other. It may be traceable into the pelvis and felt there.

Fluctuation may be detected; thrill transmitted in any direction on filling the surface.

Percussion.—Dulness in central region, intestines giving a resonant note in the lumbar regions. But little shifting of dulness on alteration of position of patient.

Dulness in flanks; central region tympanitic as patient lies on her back, and shifting with alteration of position.

Mensuration.—Distance of umbilicus from sternum, equal to or less than that from the pelvis. Greatest girth below the umbilicus.

Distance between umbilicus and sternum maintains normal ratio. Greatest girth at umbilicus or above it.

Inspection.—General roundness of abdomen; tumour may be seen somewhat rounded and prominent.

Abdomen flattened, but prominent and broad.

ABDOMINAL TUMOURS.

Tumours arising from the pelvis.—Examine per vaginam.

Ovarian. Perimetritis. Pregnant uterus.

Inflammatory swellings.—Renal or perinephritic abscess. Pelvic cellulitis. Parametritis; towards groins and iliac fossæ.

Fæcal accumulations.—Usually in colon, in either iliac fossa. There may be coincident diarrhœa.

Tubercular mesenteric glands.—Masses may be felt. Belly large and tender, emaciation, diarrhœa. Signs of **Scrofulosis**. Usually coincident signs of **Tubercular Peritonitis**. Tympanites. **Ascites**. **Abdominal Cancer** may cause enlarged glands.

Phantom Tumour.—Arises from local contraction of rectus muscle, one or both. It may be dull on percussion, and visibly prominent; usually it occurs in the lower portion of the abdomen. It subsides under chloroform. Not uncommon in **Hysteria**.

Intussusception.—Cylindrical tumour produced by intussuscepted bowel, movable from day to day. Tenesmus; passage of blood and mucus. Signs of **Obstruction of Bowels**.

PERITONITIS.

General condition of patient.—Position, complaints of pain, state of skin, tongue, pulse. Look for emaciation or other signs of chronic disease. T. = ; R. = ; P. = ; W. = . See **Ascites**; **Abdominal Tumour**; **Abdominal Cancer**; **Acute Abdominal Pain**; **Hysteria**. Examine abdomen.

Causation.—Traumatic. Rupture of bladder or other viscera. Ulceration from a gall-stone, etc., action of poisons, pressure on gut from hernia, etc. Exposure to cold. **Pyæmia**. Puerperal fever. **Bright's Disease**. Enteric fever. Enteritis. Cancer. **Tubercular Ulceration of Bowels**. Pelvic inflammations. **Perityphlitis**. **Abdominal Tumour**.

Diagnosis.—From gastritis, enteritis, metritis, cystitis, and distension of bladder; colic; abdominal hysteria.

PERITONITIS.

acute and chronic. Acute cases characterized by abdominal pain and tenderness, with fever, nausea, vomiting, constipation, abdominal distension, cold sweats. Patient usually lies on his back with the legs drawn up on the abdomen; collapse, pulse small and wiry, skin moist, extremities cold. Abdomen distended and tympanitic. There may be effusion of fluid. Bowels constipated. Respiration shallow and thoracic. Tubercular peritonitis usually occurs in young scrofulous subjects. Masses of glands may be felt in abdomen.

DISEASES OF THE LIVER.

JAUNDICE.*

A.—*Mechanical Obstruction of Bile Duct.*

- I. Obstruction by foreign bodies within the duct.
- II. Obstruction by stricture or obliteration of the duct.
- III. Obstruction by **Abdominal Tumours** closing the orifice of the duct, or growing into its interior.

B.—*Jaundice Independent of Mechanical Obstruction of the Bile Duct.*

- I. Poisons in the blood interfering with chemical changes in bile.
- II. Mineral poisons.
- III. Liver diseases.
- IV. Nervous causes.
- V. Intestinal accumulation.

Jaundice.—Shade and depth of colour. It affects also urine, sebaceous matter and sweat, milk. Taste bitter. Heart's action slow. Cerebral depression common in cases dependent upon obstruction, and when there is no obstruction tendency to stupor, coma, typhoid state. Skin liable to urticaria, lichen, boils, vitiligoidea; itchiness of skin may precede the jaundice. Digestion disturbed, constipation, flatulence, emaciation. In chronic hepatic affections hæmorrhages are common.

* See Dr. Murchison's table: "Diseases of the Liver."

JAUNDICE.

A.—*Mechanical Obstruction of Bile Duct.*

Gall-stones, inspissated bile, foreign bodies from intestines.

(a) Catarrh of duodenum, extending from gastric catarrh.

(b) Congenital defect. (c) Cicatrix after gall-stones.

. Also pressure of glands of transverse fissure of liver, amyloid or cancerous. **Cancer of Liver.**

B.—*Jaundice Independent of Mechanical Obstruction of the Bile Duct.*

Relapsing fever, enteric, typhus, pyæmia.

Phosphorus. Metallic poisons.

. Acute yellow atrophy. Congestion of liver in heart disease.

. Sudden fright.

Chronic constipation.

Indice.—Colour pale sulphur, lemon, deep olive. As it passes off skin is the last to clear. Urine may contain jaundiced casts. There may be a bitter taste from bile acids.

Diagnosis from—1. Yellow eye due to subconjunctival fat.

2. **Addison's Disease.** Here discoloration of skin is patchy and urine is normal.

3. Urine blood-coloured ; may resemble jaundice, but is also albuminous.

4. Infants soon after birth may be red and subsequently yellow, suggesting icterus neonatorum.

5. Cachexia from **Anæmia** or malignant disease.

LARGE LIVERS.*

1. **Lardaceous.** Uniform enlargement. See **Amyloid Degeneration.**
2. **Fatty.** Uniform enlargement.
3. **Hydatid tumour.** Bulging or projecting from liver.
4. Tight lacing may cause downward bulging of liver.
5. Congestion, passive, *e.g.*, from heart disease. Enlargement uniform.
6. Catarrh of bile ducts. Enlargement uniform.
7. Obstruction of common duct, *e.g.*, sequent to **Gall-stones.**
8. Pyæmic abscess. If numerous, enlargement uniform.
9. Tropical abscess, causing a bulging tumour.
10. **Cancer**, if secondary, is usually diffused, *e.g.*, secondary to cancer of sigmoid flexure or stomach.

Note size of liver, whether enlargement be uniform or irregular; whether it be tender; if accompanied by **Jaundice**. T. = Percuss; palpate and map out the liver.

Normal Liver Dulness.—Commencing posteriorly about the tenth or eleventh dorsal vertebra, it ascends slightly towards the axilla and the nipple, then again descends gradually towards the median line in front. In median line in front usually corresponds with the base of the ensiform cartilage, and to the left of this blends with the cardiac dulness at level of fifth space. In right mammary line 4—5 inches.

Cancer of Liver.—*General condition*, see **Cancer**, signs of. *Note* if jaundiced. **Liver** large; its measurements, outline, condition of surface and margin; if smooth, rough, nodular with masses. *Note* pain or tenderness. Spleen rarely enlarged. Look for other signs of **Abdominal Cancer**. **Ascites**.

* See Dr. Murchison.

LARGE LIVERS.

1. Firm, smooth, easily felt and defined.
2. Less definable ; there may be general fatty growth in the body.
3. A prominent and fluctuating tumour may be felt.
4. Tissue of liver may be healthy, and symptoms may be absent.
5. Active congestion in fevers ; frequent in tropical climates.
6. Accompanied by signs of dyspepsia and jaundice.
7. External pressure on duct may obstruct it.
8. There may be large abscesses, and irregular enlargement.
9. Usually secondary to **Dysentery**.
10. Primary cancer usually forms a mass that can be felt.

Look for—**Anæmia**. Causes capable of producing **Passive Congestion**. History of **Alcoholism** or residence in tropical climates. **Malaria**.

Cancer of Liver.—In primary cases usually cancerous masses, or large nodules, that can be felt. It may be secondary to other abdominal cancer ; then usually diffused in liver, enlarging it uniformly. Such deposits occurring may cause vomiting.

Diagnosis from—Nodular contractions of rectus muscles ;
Amyloid or Cirrhosis of Liver ; multiple hydatid.

SMALL LIVERS.

1. Simple atrophy. 2. Acute yellow atrophy. 3. Chronic atrophy.

1. Simple atrophy occurs in senile degeneration and inanition.

2. **Acute Yellow Atrophy.**—General condition much disturbed.

History.—Habits, especially as to intemperance. **Syphilis.**
Pregnancies.

Digestion.—Anorexia ; vomiting ; tongue furred.

Liver.—Note size and subsequent diminution. Jaundice, with bile in fæces.

Nervous system.—Headache ; loss of muscular power ; muscular twitchings. General disturbance of **Nervous System** tending to **Coma**.

Urine.—Urea, uric acid, and salts diminished. Presence of leucine and tyrosine, products of metamorphosis intermediate between albumen and the less complex nitrogenous compound, urea.

Causation.—**Alcoholism. Syphilis. Malaria. Typhus.** Strong emotional disturbance. Frequent pregnancy.

SMALL LIVERS.

1. No disease and no change of structure of the tissue.
2. **Acute Yellow Atrophy.**—Liver rapidly decreasing in size ; jaundice without obstruction ; symptoms of blood-poisoning.

Premonitory symptoms.—Digestion disturbed ; general vague pains. Jaundice slight, bile still appearing in fæces.

Fully established disease.—Sets in with sudden onset of symptoms due to the blood-poisoning, depending upon the defective formation of urea and uric acid ; this affects the general condition of the patient. Loss of strength. Jaundice increases ; headache, restlessness, delirium, convulsions, vomiting, coma. **Typhoid State.** Tongue dry and brown. Hæmorrhages in skin and mucous membranes may occur. Liver dulness constantly and rapidly diminishes. Spleen may enlarge.

CIRRHOSIS OF LIVER.

General condition.—**Anæmia**; emaciation; sallowness; epistaxis;
Ascites.

Digestion.—Dyspepsia; flatulence; vomiting; piles.

Spleen.—Often large.

Liver.—Usually small, but it may be enlarged in early stage; edge and surface rough, hob-nailed, hard. Jaundice may be present; then it is slight. Subsequently liver contracts. If there be peri-hepatitis, liver is tender.

SYPHILITIC DISEASE OF LIVER.

Gummata may be felt on palpation. Liver may be tender from peri-hepatitis; lobulated from irregular contraction, producing a notched margin. See signs of **Syphilis**.

CIRRHOSIS OF LIVER.

A chronic disease, mostly caused by chronic **Alcoholism**.
Dyspeptic symptoms, subsequently **Ascites** or **Hæm-**
temesis. Often associated with **Emphysema** and **Granular**
Kidneys.

Spleen large from obstruction to the return of its venous
blood.

SYPHILITIC DISEASE OF LIVER.

May result from inherited or acquired disease. There may be
gummata, a general change throughout the liver, or
peri-hepatitis.

GALL-STONES.

Occasionally they may be felt on palpation, or heard on auscultation. There may be pain on jolting or any sudden movement. They are common with cancer of gall-bladder. A stone may cause obstruction of the common duct and **Jaundice**. There may be recurrent attacks of biliary colic. Ulceration of gall-bladder may result, and extend to neighbouring organs, causing perforation of any of the hollow viscera.

BILIARY COLIC.

Attacks of severe **Abdominal Pain**, due to passage of a gall-stone from the gall-bladder to the duodenum. The attacks usually set in suddenly after exertion, and may subside suddenly, and be followed by jaundice. Attacks are apt to recur if there be many stones present.

HYDATID OF LIVER.

a chronic tumour causing an irregular outline to the liver ; usually painless, unless it be inflamed. It may be of any size ; is usually rounded, firm, slightly fluctuating. If there be no adhesion it is depressed on deep inspiration ; not accompanied by jaundice unless there be some complication. Usually single, but there may be many in the liver.

Diagnosis from — Cancer ; gummata or syphilitic liver with irregular contractions ; abscess of the liver ; distended gall-bladder ; cystic tumour of kidney ; ascites. See **Abdominal Tumours**. The spleen is not enlarged, as in some other conditions.

Course of disease.—The hydatid may suppurate and burst into the abdomen, lungs, pleura, etc. ; it may form adhesions ; it may shrink up.

Fluid in cyst.—Often removed by aspiration. It is not albuminous if there has been no inflammation. Sp. gr., about 1005 ; chlorides abundant. Microscopically, small cysts, with secondary cysts inside, may be seen ; “heads,” separate hooklets. Highly refractive particles. Cholesterin may be found in fluid.

DISEASES OF THE URINARY SYSTEM.

BRIGHT'S DISEASE.

General debility. **Anæmia.** Dyspepsia. **Œdema** or anasarca. Necessity to urinate frequently. Skin dry; often unable to perspire. **Uræmia.**

Digestion. — Dyspepsia; **Vomiting**; diarrhœa; **Hæmatemesis**; melæna.

Vascular system. — **Hypertrophy of Heart**; high tension of pulse; arteries thickened and hard; capillaries dilated on cheeks. Liability to hæmorrhages, epistaxis, hæmoptysis, etc. Excited action of the heart in uræmia.

Nervous system. — Disturbance of the general condition of **Nervous System**; **Vomiting**, **Headache**, **Vertigo**, etc. Retinitis albuminurica. See **Uræmia**, **Convulsions**.

Urine. — Albuminuria almost always present in Bright's disease. Quantity altered, usually diminished. Sp. gr. low; the total of uræa excreted diminished. Casts: fatty, hyaline, larger, small, epithelial, granular. Apparent absence of casts not an absolute proof of absence of Bright's disease, but evidence in that direction.

BRIGHT'S DISEASE.

The name signifies disease of the kidneys accompanied by **Albuminuria**, and dependent upon structural changes. The disease is usually unattended with pain, or any subjective symptoms characteristic of the disease. Pallor of the face is often a marked sign, and in elderly people is often suggestive of albuminuria. Attention must always be given in taking the history, and in observing, to determine if the disease be **Acute** or **Chronic**.

Vascular system.—May be profoundly altered and disturbed, as indicated on the other page, the blood changes being shown by anæmia, tendency to hæmorrhages, secondary inflammations, etc.

Causation.—Exanthemata, specially scarlet fever; febrile conditions, *e.g.*, pneumonia, rheumatic fever, ague, erysipelas; **Alcoholism**; exposure to cold; wet and cold work; repeated pregnancies; dyspepsia; **Gout**.

It is of great importance to determine whether the disease is acute or chronic.

Complications.—Inflammatory conditions; **Pericarditis**; **Pleurisy**; **Pneumonia**. Cerebral hæmorrhage; hæmorrhages from mucous membranes. Epistaxis. **Uremia**.

URÆMIA.

General condition.—**Anasarca. Anæmia.** Skin harsh and dry. Look for signs of **Contracted Kidneys.**

Nervous system.—**Head Pain;** drowsiness; **Delirium; Coma;** temporary blindness; retinitis albuminurica; neuro-retinitis; **Typhoid State;** muscular twitchings; **Convulsions.**

Vascular system.—Liability to hæmorrhages from mucous membranes, *e.g.*, epistaxis, **Hæmoptysis;** **Hypertrophy of Heart,** pulse hard. Pulse often strong till death is at hand.

Digestive system.—**Dyspepsia; Diarrhœa; Vomiting.**

Respiratory system.—Breath smelling ammoniacal; paroxysmal dyspnœa, resembling asthma.

Urine.—Quantity; albuminous; sp. gr. low; deficient in urea and salts; **Hæmaturia.** Casts in deposit.

Causation.—Bright's disease, acute or chronic. Suppression of urine from obstruction of ureters. Obstruction to renal veins or arteries. Destruction of one or both kidneys by abscess, calculi, etc. Cystic kidneys; surgical kidneys, sequent to stricture and pyelitis.

URÆMIA.

Many of these signs may be met with without uræmia. Inflammatory complications, *e.g.*, **Pericarditis**, **Pleurisy**. Dropsical complications, **Hydrothorax**, hydropericardium, **Ascites**. Uræmia is a condition of blood-poisoning; the breath becomes ammoniacal, and the excretion of urea is much diminished. Diarrhœa or vomiting may lead to a favourable termination. The skin seldom acts spontaneously, but its action is favourable. Symptoms may set in gradually or suddenly, with convulsions. Progress may be towards recovery, especially in acute Bright's disease; it frequently ends in death. Relapses and the recurrence of symptoms are common. Pulse full, strong, hard; heart's impulse strong.

Urine.—Scanty or suppressed from Bright's disease, **Passive Congestion** of kidneys, or pressure upon renal vessels, etc.

Causation.—Ureters may be obstructed by calculi or pressed upon by pelvic tumour, *e.g.*, ovarian. Venous congestion may result from heart disease, **Emphysema**, etc. Renal arteries may be obstructed by embolism or pressure on arteries by an **Abdominal Tumour**.

ALBUMINURIA.

Causation.—**Bright's Disease.** **Passive Congestion** of the kidneys. Simple or latent albuminuria.* Albuminuria from fevers. **Calculus** disease, due to presence of pus or blood in urine. In females from leucorrhœal discharge, etc., or menstruation.

Passive (mechanical) congestion.—**Heart Disease** or **Emphysema**, etc., may produce over-fulness of vena cava, and congestion of the kidneys. Pressure on the renal veins may also prevent return of blood from kidneys, and be due to pressure of a pregnant uterus or **Abdominal Tumour**. **Ascites** pressing on renal veins.

Urine.—Sp. gr.; quantity. Albumen, its quantity and variability under circumstances. Deposit, casts, crystals, blood discs, epithelium. Reaction.

Look for—Signs of **Bright's Disease**. Heart disease and **Diseased Vessels**. Emphysema and other lung disease Causes of passive congestion.

* Dr. Geo. Johnson : "Brit. Med. Journ." Dec. 13, 1879.

ALBUMINURIA.

Passive congestion of the kidneys.—Then the amount of albumen tends to vary with the other signs of passive congestion, *e.g.*, ascites, jaundice, cedema of legs, etc. No history of Bright's disease previous to the cause of congestion. Albumen less abundant and casts but scanty if Bright's disease is absent. Urine of high sp. gr., scanty in quantity, a few granular casts.

Simple Albuminuria, *i.e.*, not dependent upon Bright's disease.

—Urine albuminous without any marked structural lesion. May be due to exposure to cold; excess of nitrogenous food. Often accompanied by oxalates. During fevers and febrile conditions, *e.g.*, typhus, enteric, cholera, diphtheria, pneumonia, rheumatic fever. But few casts, if any.

HÆMATURIA.

Causation.—Disease in renal tissue, pelvis of kidney, ureter, bladder, urethra. **Bright's Disease**, acute or chronic. Passive congestion of kidneys. See **Passive (Cardiac) Congestion**. Active congestion of kidneys from alcohol, turpentine, cantharides. Traumatic injury. Stone. See **Renal Calculus**. **Bladder, Disease of:** cystitis, stone, cancer, villous growth, etc. In females during menstrual period. **Paroxysmal Hæmaturia**.

Urine.—Albuminous, alkaline, smoky, blood-coloured, porter-like. Containing hæmatin, but no corpuscles. See if in subsequent course albumen occurs without blood. Note the colour in relation to the amount of albumen and sp. gr.

Deposit.—Lithates with high sp. gr. from renal congestion. Blood casts; renal casts; epithelial and hyaline casts in Bright's disease; granular and hyaline in renal congestion. Crystals.

Blood may be mixed with the urine; in clots; in clots moulded in ureter.

Note quantity of urine, and any difficulty in micturition.

HÆMATURIA.

If blood comes from the renal structure usually there are blood-casts and smoky urine ; if from the urinary passages no casts ; if from the bladder or urethra pure blood and clots may be passed, usually after micturition. Periodical attacks of discharge of porter-like urine, with granules and hyaline casts, and the deposit of brownish granular matter in place of corpuscles. See paroxysmal hæmaturia. Hæmaturia may appear in early inflammation, and in acute exacerbations. Occasionally late in cirrhosis. Rare in lardaceous disease.

Paroxysmal Hæmaturia.—At irregular intervals sudden attacks of rigors, the next urine passed being loaded with blood. Health may long continue good. The *paroxysms* are unattended with pain ; there may be a feeling of chilliness across the loins, weakness, nausea, vomiting, joint-pain. The patient becomes languid, weak, anæmic. See **Anæmia**. Examine heart and vascular system. Optic discs. See **Hæmaturia**, with description of urine ; **Renal Calculus**.

Causation.—It is independent of any known structural change in the kidneys. Supposed to be connected with ague, rheumatism, exposure to cold ; certainly such exposure may excite the paroxysms. It almost always occurs in males, usually adults. There is sometimes oxaluria.

BRIGHT'S DISEASE, ACUTE.

Signs and symptoms.—Anasarca. Suppression of urine, more or less complete. Skin harsh. Tendency to somnolence, head-pain, vomiting, coma. **Uræmia.** Usually after exposure to cold or scarlet fever. It may resolve or terminate in a large white kidney.

Urine.—Smoky; very albuminous; blood discs and large epithelial casts abundant. In quantity, scanty. Blood casts.

GRANULAR CONTRACTED KIDNEYS.

Signs and symptoms.—If any œdema it is slight and transient. Heart hypertrophied; pulse of high tension. Liability to epistaxis and hæmorrhages from mucous membranes. Albuminuric retinitis. Tendency to uræmia. Commonest in advanced life. Often there is coincident cirrhosis of the liver.

Urine.—Clear, with little deposit; quantity large; albumen, a trace. Small granular and hyaline casts. Sp. gr. low.

FATTY KIDNEYS.

Signs and symptoms.—Usually anasarca. Face pale and puffy. Has a fatal tendency. May result from acute Bright's disease. Not uncommon in phthisis.

Urine.—Fairly copious; albumen much. Fatty casts; fatty cells. Sp. gr. rather low.

AMYLOID KIDNEYS.

Signs and symptoms.—Anasarca moderate. Pasty anæmic look. Emaciation and signs of amyloid disease of other organs: spleen, liver, intestines.

Urine.—Urine copious. Sp. gr. various. Much albumen; a few hyaline casts.

LARGE WHITE KIDNEYS.

Signs and symptoms.—Anasarca. Anæmia. Results from acute Bright's disease. Liability to intercurrent acute attacks, with increase of the symptoms.

Urine.—Scanty; pale. Casts, hyaline or granular. During exacerbations blood in urine. Albumen.

BRIGHT'S DISEASE, ACUTE.

Kidney enlarged and congested, the whole structure of the organ being involved. Cortex much swollen ; pyramids very dark and congested ; glomeruli large and congested. Epithelium swollen and cloudy. Veins of the surface dilated.

GRANULAR CONTRACTED KIDNEYS.

Kidney small ; capsule adherent ; surface granular and reddish. Frequent cysts in cortex. Much wasting of cortex. Arteries thickened.

FATTY KIDNEYS.

Kidney large, yellow, pale, soft, easily broken down.

AMYLOID KIDNEYS.

Kidney large and pale ; surface smooth ; cortex thick ; glomeruli and vessels stain with iodine.

LARGE WHITE KIDNEYS.

Kidney large, smooth, white. Cortex much swollen from over-development of epithelium in convoluted tubes ; but little change in Malpighian tufts.

BRIGHT'S DISEASE.

ACUTE.

Causation.—Cold. Scarlet fever.

Anasarca.—Present.

Heart and pulse.—No hypertrophy. There may be palpitation in uræmia.

Ophthalmoscopic appearances.—Usually no changes.

Urine.—Smoky colour. Casts, with large granular epithelium and blood.

Duration of albuminuria.—Short period.

CHRONIC.

Alcoholism. Gout degeneration.

Present with fatty loid, and large white usually absent with kidney.

Hypertrophy; pul

May be hæmorrhagic retinitis albuminuric

Hyaline casts, large small; granular cast casts.

Many months.

BLADDER, DISEASES OF.

Disease of the bladder and genito-urinary excretory apparatus may be indicated by—

1. *Urine*.—Thick, with deposit of mucus, pus, phosphates, blood, etc.; reaction alkaline; smell offensive. Such urine is passed with cystitis.

2. *Micturition difficult*.—This may be from stricture of the urethra, a bladder paralysed with retention, or overflow, or complete incontinence. This may arise from disease of the **Spinal Cord** or **Brain Disease**, **Meningitis**, or **Hysteria**.

3. Hypogastric pain and tenderness with fever.

Cystitis may be acute or chronic. It may result from paralysis or atony of the bladder, calculus, cancer, villous growth. Much mucus renders the urine alkaline by causing the breaking up of the urea into ammonia salts; phosphates are then precipitated. Cystitis is a common and grave complication of **Disease of the Cord**; in such cases it is usually painless.

RENAL CALCULUS.

A chronic condition ; liability to acute attacks.

Chronic course.—Aching continuous pain in one lumbar region, shooting downwards. Occasional passage of blood-stained urine, pus, gravel, epithelial debris. **Hæmaturia**, especially after jolting exercise. There may be tenderness in the loin. Occasional attacks of renal colic. Bladder : there may be signs of stone in the bladder, or **Cystitis**.

Enquire for history of attacks of renal colic, **Gout**, **Uremia**, signs of disease of bladder.

Complications.—Stone in the bladder. Nephritic or perinephritic abscess. Suppression of urine from impaction of calculus in ureter on each side.

Urine.—Quantity.

RENAL CALCULUS.

Urine.—Varying on different occasions. It may be mixed with blood, usually not forming clots. **Albuminuria** usually proportioned to the amount of blood unless the kidneys are degenerated; then albuminuria may occur in degree over and above the albumen due to the blood. There may be crystals of oxalates or uric acid, etc. Usually no casts.

Renal Colic.—Attacks may come on without any previous symptoms of calculus. In the attack paroxysmal pain in one lumbar region, severe, causing collapse, vomiting, and sometimes suppression of urine. The attack may cease suddenly; then the next urine passed may be bloody, and may bring away the calculus per urethram. Such paroxysms especially occur after exertion; they may last days or weeks. There is often retraction of the testicle on the side of pain (irritation of the genito-crural nerve); the pain shoots down the inner side of the thigh, and is accompanied by frequent desire to micturate.

URINE, DESCRIPTION OF.

Quantity.—In healthy adult forty to sixty ounces per diem.

Colour.—Light or dark sherry ; colourless ; smoky ; bleached or coloured.

Reaction.—Acid (normal) ; neutral ; alkaline.

Sp. gr.—Normal, 1015—1025.

Urea.—Normal, 400—600 grains per diem ; 1·5 per cent 4·0 per cent.

Albumen.—Abnormal. See **Albuminuria**.

Sugar.—Abnormal. See **Diabetes**.

Deposit.—Bulk in proportion to urine ; colour ; light or heavy.

CHEMICAL EXAMINATION OF THE DEPOSIT.

Phosphates.—Soluble in nitric acid ; insoluble in liq. potash. Urine usually alkaline.

Lithates.—Soluble in liq. potassæ, or on warming deposit. Urine when warm as passed is clear.

Uric acid.—Soluble in liq. potassæ, and precipitated from solution by hydrochloric acid. See **Murexide Test**.

Mucus.—Coagulated by boiling with liq. potassæ.

MICROSCOPICAL EXAMINATION OF THE DEPOSIT.

1. *Casts*.—Large, small, hyaline, granular, epithelial, containing large swollen epithelium ; blood casts.
2. *Crystals*.—(a) Triple phosphate : Triangular prisms, small or large ; when very short they may be mistaken for hexahedral oxalates. (b) Uric acid : Usually coloured ; crystals regular, lozenge-shaped or square, elongated or acicular. (c) Oxalates : Octahedra with bright centres. Dumb-bells. (d) Cystine : Hexagonal plates.
3. *Epithelium*.—Glandular ; squamous from vagina or bladder.
4. *Pus*.—

URINE, DESCRIPTION OF.

Quantity.—Increased in **Diabetes**.

Colour.—May indicate **Jaundice**; **Hæmaturia**; greenish in diabetes.

Reaction.—In alkaline urine usually a deposit of phosphates.

Sp. gr.—Dense in **Diabetes**, or if much urea, etc. Low in **Granular Contracted Kidneys**.

Urea.—Usually a large percentage if sp. gr. is high without sugar.

Albumen.—May be a transient ingredient, therefore look for it repeatedly.

Sugar.—Occasionally present in **Brain Disease**.

Deposit.—Give general, chemical, and microscopical characters.

CLINICAL INDICATIONS OF THE DEPOSIT.

Phosphates.—Common in hot weather, when urine ferments readily. Abundant when there is much mucus or pus.

Lithates.—Copious deposit in febrile conditions and in **Passive Congestion** of kidneys; also usually in healthy scanty urine.

Uric acid.—Like grains of cayenne pepper. May indicate gouty tendency or calculus-formation. Deranged liver.

Mucus.—Copious in cystitis.

Casts.—Coming from uriniferous tubes indicate their condition. Numerous in acute **Bright's Disease**; abundant and varied in inflammation; few in lardaceous disease; few in cirrhosis; common in other cases of albuminuria.

Crystals.—(a) Triple phosphates are common in alkaline urine in cystitis, and in urine that has decomposed. (b) Uric acid: Deposited in the gouty diathesis. (c) Oxalates: Dyspepsia may produce oxaluria, so anæmia. (d) Cystine may form calculi.

Epithelium.—Common in Bright's disease.

Pus from a pelvic abscess may be discharged into the bladder or urinary tract. Copious in renal abscess, and in cystitis.

NORMAL CONSTITUENTS OF URINE.

Chlorides.—A few drops of nitric acid, then an excess of solution of *nitrate of silver*; white precipitate of chloride of silver thrown down. (N.B.—Nitric acid prevents phosphate being precipitated.) Wash precipitate and prove its solubility in ammonia.

Phosphoric Acid.—(a) Solution of *nitrate of silver* gives a white precipitate of phosphate of silver, soluble in nitric acid, but insoluble in ammonia. (b) To urine tested as below for sulphuric acid, and thus deprived of sulphates, add excess of ammonia; phosphate of baryta is thrown down.

Sulphuric Acid.—Add a few drops of nitric acid, then *chloride of barium*, which gives a white precipitate of the sulphate.

Urea.—If the sp. gr. of the urine be from 1023—1030 it usually crystallizes with an equal bulk of nitric acid, the solution being cooled. Beautiful crystals of *nitrate of urea* are formed. See quantitative examination.

Uric Acid.—Precipitated from urine by *hydrochloric acid*, and waiting twenty-four hours. Soluble in liq. potassæ. See **Murexide Test**.

(a) **Quantity.***—1. Diminished in early inflammatory conditions.

2. Normal in middle inflammatory stage, and in early stage of cirrhosis.

3. Increased in lardaceous kidney, and here may precede albuminuria. In cirrhosis, late stage. Sometimes in advanced inflammation and during absorption of dropsies.

4. Suppressed in acute and advanced inflammation, and in advanced cirrhosis.

(b) **Sp. Gr. and Solids.**—Depend upon water, urea, other solids.

* After Dr. Grainger Stewart.

ABNORMAL CONSTITUENTS OF URINE.

Albumen.—1. Heat urine, and when boiled add nitric acid ; a precipitate indicates albumen.

2. Float in test-tube on nitric acid ; a non-crystalline cloud at the junction of the two fluids indicates albumen.

Sugar: Moore's test.—Mix urine with half its volume of liq. potassæ and boil ; a brownish colour shows sugar.

Trommer's test. — Add to urine one or two drops of solution of sulphate of copper, then about half as much liq. potassæ as urine. If sugar be present, the precipitate at first produced dissolves, producing a blue solution. Now boil this solution ; sugar causes decomposition, and the brown oxide of copper is precipitated.

Fehling's test.—Cupric sulphate, 40 grammes ; potass. tartrat., 160 grammes ; liq. sodæ (sp. gr. 1·12), 750 grammes ; distilled water to 1,154·5 c.c. Boil some of this solution ; then add urine, a few drops at first, and if it be saccharine the red suboxide of copper precipitates at once.

Bile: Pettenkofer's test.—Dissolve a grain or two of white sugar in a drachm of urine ; then add, drop by drop, strong sulphuric acid. A characteristic violet-red colour will be produced if bile be present.

Urine.—A morbid product, crystallizes as small spheres which are composed of acicular crystals which radiate from a common centre.

Urosine.—Crystallizes in long white needles.

URINARY CALCULI.

Heat a specimen on platinum foil over spirit flame ; afterwards with blowpipe.

- I. It *burns away*, leaving only a minute trace of ash, probably either **Uric Acid**, **Urate of Ammonia**, or **Cystine**. Proceed to test calculus with (a) liq. potassæ ; soluble. See **Uric Acid**. (b) Soluble in hot water or with liq. potassæ, evolving ammonia = urate of ammonia. (c) Insoluble in hot water, but readily soluble in ammonia, the solution on evaporation giving hexagonal plates = **Cystine**.
- II. It proves *incombustible* before the blow-pipe. (a) Soluble in dilute hydrochloric acid = **Phosphate of Lime**. Ammonia added to such solution gives an amorphous precipitate. (b) *Fusible* before blow-pipe and soluble in hydrochloric acid = **Triple Phosphate**. The precipitate produced by ammonia from the solution is crystalline. (c) Before ignition soluble without effervescence in hydrochloric acid, this acid solution giving a white precipitate with ammonia. After ignition soluble with effervescence in hydrochloric acid, this solution giving no precipitate with ammonia = **Oxalate of Lime**.

URINARY CALCULI.

Uric acid and phosphatic calculi common.

Murexide test of Uric Acid.— Dissolve the substance to be tested in nitric acid, and gently warm; when cold touch residue with liq. potassæ; a beautiful purple solution indicates uric acid

SIGNS OF PREGNANCY.

General condition.—Chloasma.

Nervous system.—Head-ache. Altered mental condition, sometimes great sleepiness, at other times insomnia. Neuralgic pains of all kinds.

Vascular.—Heart beats become more frequent. Pulse of high tension.

Respiratory.—Dyspnoea on exertion. Cough, reflex.

Digestive.—Heartburn, salivation. Nausea may occur at once, but commonly not till second month. Sense of sinking in epigastrium ; cravings for food.

Digestive disturbance.—Appetite increased ; may be strangely altered or perverted. Vomiting ; morning sickness. Bowels disturbed ; piles from pressure ; often constipation.

Urine.—Kystein floats as a pellicle on urine after it has stood twenty-four to thirty-six hours, subsequently falling as a milky deposit. Not a sure sign. Albuminuria.

Diagnosis from—Ascites ; amenorrhœa from other causes. Ovarian dropsy. Phantom tumour ; abdominal tumour.

Duration of pregnancy.—Calculate the full time complete at forty weeks, dating from a fortnight after commencement of last menstruation.

PREGNANCY.—COINCIDENT SIGNS AND SYMPTOMS.

Special signs of pregnancy. — 1. Suppression of the menses, under the climacteric age, and without anæmia or known uterine disease. Balottement.

2. *Changes in the breasts* which early become somewhat enlarged, their sensitiveness increased, with a feeling of fulness, weight, and shooting pains. Veins in skin enlarged, the glands feeling hard and knotty, and being tender, and sometimes subcutaneous fat augmented. Areola darkened, with a secondary areola outside; moist with enlarged sebaceous follicles, milk in breast in last month. Nipples turgid and prominent.

3. *Changes in abdomen.* — No visible tumour till third month. At first umbilicus is sunken from growth of fat; later it is protruded by internal pressure. Uterine tumour elastic; very slightly fluctuating; at fifth month reaches half up to umbilicus.

4. *Auscultation of uterus.* — Fœtal heart. Placental souffle.

Complications. — Obstruction of bowels; hæmorrhage from uterus.

Pressure signs. — On bladder. Piles.

INDEX

-
- Abdomen, examination of, 166
 - Abdominal cancer, 162
 - pain, acute, 156
 - pulsation, 167
 - tumour, 166
 - Acute Bright's disease, 192
 - Acute yellow atrophy of liver, 179
 - Addison's disease, 28
 - Adynamia, *see* Typhoid state, 44
 - Ague, 12
 - Albumen, 201
 - Albuminuria, 188
 - latent, 189
 - Alcoholism, 86
 - acute, 88
 - Amnesia, 41
 - Amphoric respiration, 128
 - Amyloid degeneration, 24
 - kidneys, 192
 - Anæmia, 20
 - pernicious, 20
 - Anæsthesia, 57
 - muscular, 56
 - Analgesia, 57
 - Anasarca, 24
 - Aneurism, 120
 - abdominal, 169
 - Angina pectoris, 115
 - Ankle clonus, 48
 - Aortic obstruction, 109
 - Aortic regurgitation, 109
 - Aphasia, 41
 - Aphonia, 149
 - Appetite, 150
 - Arteries, 122
 - Arthritis, 34
 - Ascites, 165
 - Ascites or ovarian dropsy, 170
 - Asthma, 146
 - Ataxy, 94
 - Atheroma, 123
 - Athetosis, 52
 - Aura, 57
 - Auscultation, 128
 - Bell's paralysis, 61
 - Bile in urine, 201
 - Biliary colic, 182
 - Bilious attacks, 41
 - Bladder, disease of, 195
 - Blood in expectoration, 132, 133
 - motions, 158
 - urine, 190
 - vomit, 133, 156
 - Bone disease, 17, 22
 - Bowels, state of, 150
 - obstruction of, 158
 - ulceration of, 164
 - Brain disease, 64
 - Bright's disease, 184
 - acute, 192

- use, acute orchonic, 46
 skin, 29
 ble, 21
 ysis, 70
 heat, 127

 ry, 182
 96
 202

 al, 162
 ines, 163
 177
 ch, 163
 estion, 106
 n, 111
 nents, 112
 phy, 110
 y, 198
 ingitis, 82
 82
 3, 45
 nents of, 126
 of, 124
 se-taking, xiv
 urine, 200
 ani, 61

 35
 ht's disease, 192
 n, 54
 iver, 180
 , 50
 , 182
 7

 , 14
 effects of heart, 118
 philis, 14
 assive, 106
 of lung, 137
 of lung, 136
 50
 of limbs, 54

 Cough, 130
 Cracked pot sound, 128
 Cramp, 50
 Cranial nerves, 58
 Craniotabes, 14
 Crepitations, 130
 Cross paralysis, 70
 Crystals in urine, 199
 Cycloplegia, 67
 Cystine, 202
 Cystitis, 195

 Delirium, 44
 tremens, 87
 Developmental defects, 31
 Diabetes, 28
 Diarrhœa, 154
 Digestive functions, 150
 Dilatation of heart, 110
 Diphtheria, 8
 Diphtheritic paralysis, 100
 Diplopia, 58
 Displacements of heart, 112
 Dorsal decubitus, 42
 Drunkenness, 86
 Duchenne's disease, 70
 Dysæsthesia, 57
 Dysentery, 165
 Dysphagia, 156
 Dyspnoea, 134

 Electric tests, 46
 Emaciation, 22
 Embolism, 122
 Emphysema, 144
 Empyema, 139
 Enteric fever, 4
 diagnosis from tuberculosis,
 27
 Epilepsy, 78
 Epistaxis, 12
 Erysipelas, 8
 Examination of abdomen, 166
 Exophthalmos, 98
 Expectoration, 132
 Extensors, paralysis of, 70

- Face, paralysis of, 61, 75
 Facial nerve, 60
 Facial palsy, cerebral, 75
 Facial spasm, 50
 Fatty kidneys, 192
 Fehling's test, 201
 Fever, signs of, 2
 enteric, 4
 scarlet, 4
 typhus, 4
 Fevers, specific, 4
 Fine movements, 46
 Functional paralysis, 47

 Gait, 54
 Gall-stones, 182
 Gastric crises, 94
 Gastric ulcer, 160
 General paralysis of insane, 89
 Girding sensation, 57
 Glosso-labio-laryngeal palsy, 70
 Glosso-pharyngeal nerve, 62
 Goitre, exophthalmic, 98
 Gonorrhœal rheumatism, 38
 Gout, 38
 Granular contracted kidneys, 192
 Graves' disease, 98
 Gummata, 17
 Gums, 152

 Hæmatemesis, 156
 Hæmaturia, 190
 paroxysmal, 191
 Hæmoptysis, 132
 Hæmoptysis or Hæmatemesis, 133
 Hæmorrhage in retina, 65
 Headache, 41
 Head-pain, 40
 Hearing, 58
 Heart, congenital defects of, 118
 dilated and hypertrophied, 110
 disease, 112
 displacements of, 112

 Heart, hypertrophy, 110
 physical examination of, 102
 Hemianæsthesia, 57
 Hemipopia, 41
 Hemiplegia, 74
 Herpes zoster, 100
 Hooping-cough, 12
 Hydatids, 183
 Hydrocephalus, 84
 Hydrothorax, 139
 Hyperæsthesia, 57
 Hyperpyrexia, 5
 Hysteria, 78
 Hygienic conditions, 1

 Idiots, 19
 Illusions, 88
 Infantile paralysis, 96
 Insanity, 88
 Insomnia, 40
 Instructions for case-taking, xi
 additional for children, xiv
 Intelligence, 40
 Intercostal neuralgia, 72
 Intermittent hæmaturia, 191
 Intussusception, 171
 Iridoplegia, 67
 Iritis, 16

 Jaundice, causation, 174
 Joints, 34

 Knee-jerk, 46
 Kidney, tumour of, 167
 Kidneys, amyloid, 192
 fatty, 192
 granular contracted, 192
 large white, 192

 Labio-glosso-laryngeal palsy, 70
 Laryngeal obstruction, 8, 148
 symptoms, 8, 148
 Laryngitis, 149
 Laryngismus, 52, 149
 Larynx, disease of, 149

- Lead poisoning, 98
 Leucine, 201
 Lightning pains, 69, 95
 Liver, acute yellow atrophy,
 178
 cancer of, 176
 cirrhosis of, 180
 large, 176
 small, 178
 syphilitic disease of, 180
 Locomotor ataxy, 94
 pulse, 104
 Lung contraction, 136
 solidification, 137

 Malformation of heart, 118
 Mania, post epileptic, 80
 Marasmus, 23
 Measles, 6
 Melæna, 158
 Meningitis, 82
 Mercurial tremor, 53
 Michel's ganglion, 61
 Migraine, 41
 Miliaria, 37
 Minor paralyses, 70
 Nasal obstruction, 108
 regurgitation, 108
 Motor power, 54
 Mouth and throat, 152
 Movements of extremities, 54
 fine, 46
 of head and trunk, 54
 Mucous patches, 14
 Muscular anæsthesia, 56
 movements, hysterical, 78
 Murexide test, 203

 Nerve, recurrent laryngeal, 62
 superior laryngeal, 61
 Nervous system, 40
 Neuralgia, 70
 conditions characterized by,
 70
 trigeminal, 72
 Nodes, rheumatic, 37

 Normal respiration, 128
 Nutrition, 22
 Nystagmus, 58

 Obstruction of bowels, 158
 Oedema, 24
 pulmonary, 134
 Ophthalmoplegia interna, 67
 Ophthalmoscopic appearances, 64
 Optic discs, 64
 atrophy, 64
 neuritis, 99
 otorrhœa, 58
 Ovarian dropsy or ascites, 170
 Ovarian tumour, 167, 170
 Oxalates, 198

 Pain, abdominal, 156
 Palate, 152
 Palpitation, functional or or-
 ganic, 114
 Palsy of cranial nerves, 58
 Paralysis, 46
 agitans, 90
 diphtheritic, 100
 functional and organic, 47
 infantile, 96
 labio-glosso-laryngeal, 70
 of extensors of forearm,
 98
 of face, 61, 75
 of face, Bell's, 61
 Paraplegia, 68
 Paroxysmal hæmaturia, 191
 Passive cardiac congestion, 106
 Patellar tendon reflex, 48
 Pernicious anæmia, 20
 Pericarditis, 116
 Periostitis, 16
 Peritoneum, fluid in, 168
 Peritonitis, 172
 Phantom tumour, 171
 Pharynx, 152
 Phlebitis, 123
 Phosphates, 198
 Phosphoric acid, 200

- Phthisis, 140
 Physical examination of chest,
 126
 of heart, 104
 Pleural friction sound, 130
 Pleurisy, 139
 Pleuritic effusion, 138
 Plumbism, 98
 Pneumogastric nerve, 60
 Pneumonia, 138, 142
 or pleurisy, 138
 Portal circulation, 107
 Post-epileptic mania, 80
 Pressure signs in chest, 120
 Progressive anæmia, 20
 Progressive muscular atrophy, 70
 Pseudo-hypertrophic paralysis,
 70
 Ptosis, 58
 Puerile respiration, 129
 Puerperal fever, 10
 Pulmonary œdema, 134
 Pulsation, abdominal, 167
 Pulse, 104
 Pus in urine, 199
 Pupils, 66
 Pyæmia, 10

 Râles, 130
 Reflexes, 48
 Regions of chest, 124
 Renal calculus, 196
 Renal colic, 197
 Recurrent laryngeal nerve, 62
 Respiration, bronchial, 128
 cavernous, 128
 harsh, 128
 tubular, 128
 Respiratory movements, 126
 Retraction of chest, 126
 of lung, 126
 Rheumatism, 36
 gonorrhœal, 38
 Rheumatoid arthritis, 34
Rhonchi, 130
Rhythm of respiratory act, 126

Rhythmical muscular m
 ments, 52
 Rickets, 22
 Rigidity, 74, 96
 Rupia, 16

 Sanitation, 1
 Senile degeneration, 32
 Scarlet fever, 4
 Scars, 15
 Sciatica, 72
 Sclerosis of cord, 77, 90
 Scrofulosis, 25
 Sensation, 56
 Sensations of girthing, 57
 Sight, 58
 Sleep, 40
 Small-pox, 6
 Smell, 58
 Solidification of lung, 137
 Spasm, tonic, 50
 clonic, 50
 facial, 50
 Special senses, 58
 Specific fevers, 4
 Speech, 40
 Spinal cord, disease of, 68
 Spleen, large, 169
 Sputum, 132
 Stomach, ulcer of, 160
 Stiff neck, 36
 Strychnia poisoning, 93
 Subsultus tendinum, 44
 Sugar, tests for, 201
 Syphilis, acquired and con
 tional, 16
 Syphilitic disease of liver,

 Tabes dorsalis ataxy, 94
 Tables, ascites or ovarian tur
 170
 convulsions, epilepti
 hysterical, 81
 enteric fever or tul
 losis, 27
 hydrocephalus or ricke

- Cables, joints, 3
 palpitation, functional or organic, 114
 alsy, functional or organic, 47
 pneumonia or pleuritic effusion, 138
 sclerosis or chorea, 77
 specific fevers, 4
 tetanus or strychnia poisoning, 93
 valvular lesions, 108
 Cape worm, 150
 Taste, 58
 Teeth, 15, 152
 Tetanus, 92
 Thoracic aneurism, 120
 Throat, 152
 Thrombosis, 123
 Tinnitus, 59
 Tongue, 152
 Tonic spasm, 50
 Tonsils, 152
 Tonsillitis, rheumatic, 37
 Tophi, 39
 Tremor, 52
 muscular, 53
 Tricuspid regurgitation, 118
 Trigeminal neuralgia, 72
 Tubercle in choroid, 65
 Tubercular ulceration of bowels, 165
 Tuberculosis, miliary, 26
 or enteric fever, 27
 Tumour, abdominal, 166
 kidney, 167
 ovarian, 167
 Tumour, brain, 82
 Typhilitis, 160
 Typhoid fever, 4
 state, 44
 Typhus, 4
 Tyrosine, 201
 Ulcer, gastric, 160
 Ulceration of bowels, 164
 tubercular, 165
 Uræmia, 186
 Urea, 200
 Urinary calculi, 202
 casts, 198
 crystals, 198
 Uric acid, 198, 203
 Urine, description of, 198
 deposits, examination of, 198
 normal constitution of, 200
 Uvula, 152
 Valleix, points of, 71
 Valvular heart disease, 112
 Varicella, 6
 Varicose veins, 123
 Variola, 6
 Vascular system, 102
 Veins, disease of, 123
 Vertigo, 42
 Vessels, disease of, 123
 Vomica, 140
 Vomiting, 44, 154
 cerebral, 44
 Worms, intestinal, 150
 Writer's cramp, 50

1

.

1

2

.

.

J. & A. CHURCHILL'S
MEDICAL CLASS BOOKS.

ANATOMY.

BRAUNE.—An Atlas of Topographical Anatomy, after Plane Sections of Frozen Bodies. By WILHELM BRAUNE, Professor of Anatomy in the University of Leipzig. Translated by EDWARD BELLAMY, F.R.C.S., and Member of the Board of Examiners; Surgeon to Charing Cross Hospital, and Lecturer on Anatomy in its School. With 34 Photo-lithographic Plates and 46 Woodcuts. Large Imp. 8vo, 40s.

LOWER.—Diagrams of the Nerves of the Human Body, exhibiting their Origin, Divisions, and Connexions, with their Distribution to the various Regions of the Cutaneous Surface, and to all the Muscles. By WILLIAM H. FLOWER, F.R.C.S., F.R.S. Third Edition, containing 6 Plates. Royal 4to, 12s.

GODLEE.—An Atlas of Human Anatomy: illustrating most of the ordinary Dissections and many not usually practised by the Student. By RICKMAN J. GODLEE, M.S., F.R.C.S., Assistant-Surgeon to University College Hospital, and Senior Demonstrator of Anatomy in University College. With 48 Imp. 4to Coloured Plates, containing 112 Figures, and a Volume of Explanatory Text, with many Engravings. 8vo, £4 14s. 6d.

HEATH.—Practical Anatomy: a Manual of Dissections. By CHRISTOPHER HEATH, F.R.C.S., Holme Professor of Clinical Surgery in University College and Surgeon to the Hospital. Fifth Edition. With 24 Coloured Plates and 269 Engravings. Crown 8vo, 15s.

11, *NEW BURLINGTON STREET.*

J. & A. Churchill's Medical Class Books.

ANATOMY—*continued.*

HOLDEN.—**A Manual of the Dissection of the Human Body.** By LUTHER HOLDEN, F.R.C.S., Consulting-Surgeon St. Bartholomew's Hospital. Fifth Edition, by JOHN LANG, F.R.C.S., Surgeon to, and Lecturer on Anatomy at, St. Bartholomew's Hospital. With 208 Engravings. 8vo, 20s.

By the same Author.

Human Osteology : comprising a Description of the Bones, with Delineations of the Attachments of Muscles, the General and Microscopical Structure of Bone and its Development. Sixth Edition, revised by the Author JAMES SHUTER, F.R.C.S., late Assistant-Surgeon to St. Bartholomew's Hospital. With 61 Lithographic Plates and 89 Engravings. Royal 8vo, 16s.

ALSO,

Landmarks, Medical and Surgical. Third Edition. 8vo, 3s. 6d.

MORRIS.—**The Anatomy of the Joints of Man.** By HENRY MORRIS, M.A., F.R.C.S., Surgeon to, and Lecturer on Anatomy and Practical Surgery at, the Middlesex Hospital. With 19 Plates (19 Coloured) and Engravings. 8vo, 16s.

The Anatomical Remembrancer; or, Complete Pocket Anatomist. Eighth Edition. 32mo, 3s. 6d.

WAGSTAFFE.—**The Student's Guide to Human Osteology.** By WM. WARWICK WAGSTAFFE, F.R.C.S., late Assistant Surgeon to, and Lecturer on Anatomy at, St. Thomas's Hospital. With 23 Plates and 66 Engravings. Fcap. 8vo, 10s. 6d.

WILSON — BUCHANAN — CLARK. — **Wilson's Anatomist's Vade-Mecum : a System of Human Anatomy.** Third Edition, by GEORGE BUCHANAN, Professor of Clinical Surgery in University of Glasgow, and HENRY E. CLARK, M.R.C.S., Lecturer on Anatomy in the Glasgow Royal Infirmary School of Medicine. With 450 Engravings, including 26 Coloured Plates. Crown 8vo, 18s.

11, NEW BURLINGTON STREET.

J. & A. Churchill's Medical Class Books.

BOTANY.

BENTLEY.—A Manual of Botany. By Robert

BENTLEY, F.L.S., M.R.C.S., Professor of Botany in King's College and to the Pharmaceutical Society. With 1185 Engravings. Fourth Edition. Crown 8vo, 15s.

By the same Author.

**The Student's Guide to Structural,
Morphological, and Physiological Botany. With 660 Engravings.**
Fcap. 8vo, 7s. 6d.

ALSO,

**The Student's Guide to Systematic
Botany, including the Classification of Plants and Descriptive
Botany. With 357 Engravings. Fcap. 8vo, 3s. 6d.**

BENTLEY AND TRIMEN.—Medicinal Plants:

being descriptions, with original Figures, of the Principal Plants employed in Medicine, and an account of their Properties and Uses. By ROBERT BENTLEY, F.L.S., and HENRY TRIMEN, M.B., F.L.S. In 4 Vols., large 8vo, with 306 Coloured Plates, bound in half morocco. gilt edges, £11 11s.

CHEMISTRY.

BERNAYS.—Notes for Students in Chemistry;

being a Syllabus of Chemistry compiled mainly from the Manuals of Fownes-Watts, Miller, Wurz, and Schorlemmer. By ALBERT J. BERNAYS, Ph.D., Professor of Chemistry at St. Thomas's Hospital. Sixth Edition. Fcap. 8vo, 3s. 6d.

By the same Author.

Skeleton Notes on Analytical Chemistry,
for Students in Medicine. Fcap. 8vo, 2s. 6d.

BLOXAM.—Chemistry, Inorganic and Organic;

with Experiments. By CHARLES L. BLOXAM, Professor of Chemistry in King's College. Fifth Edition. With 292 Engravings. 8vo, 16s.

By the same Author.

**Laboratory Teaching; or, Progressive
Exercises in Practical Chemistry. Fourth Edition. With 83
Engravings. Crown 8vo, 5s. 6d.**

11, NEW BURLINGTON STREET.

CHEMISTRY—continued.

BOWMAN AND BLOXAM.—**Practical Chemistry**, including Analysis. By JOHN E. BOWMAN, formerly Professor of Practical Chemistry in King's College, and CHARLES L. BLOXAM, Professor of Chemistry in King's College. With 98 Engravings. Seventh Edition. Fcap. 8vo, 6s. 6d.

BROWN.—**Practical Chemistry: Analytical Tables and Exercises for Students.** By J. CAMPBELL BROWN, D.Sc. Lond., Professor of Chemistry in University College, Liverpool. Second Edition. 8vo, 2s. 6d.

CLOWES.—**Practical Chemistry and Qualitative Inorganic Analysis.** An Elementary Treatise, specially adapted for use in the Laboratories of Schools and Colleges, and by Beginners. By FRANK CLOWES, D.Sc., Professor of Chemistry in University College, Nottingham. Third Edition. With 47 Engravings. Post 8vo, 7s. 6d.

FOWNES.—**Manual of Chemistry.**—*See WATTS.*

FRANKLAND AND JAPP.—**Inorganic Chemistry.** By EDWARD FRANKLAND, Ph.D., D.C.L., F.R.S., and F. R. JAPP, M.A., Ph.D., F.I.C. With 2 Lithographic Plates and numerous Wood Engravings. 8vo, 24s.

LUFF.—**An Introduction to the Study of Chemistry.** Specially designed for Medical and Pharmaceutical Students. By A. P. LUFF, F.I.C., F.C.S., Lecturer on Chemistry in the Central School of Chemistry and Pharmacy. Crown 8vo, 2s. 6d.

TIDY.—**A Handbook of Modern Chemistry**, Inorganic and Organic. By C. MEYMOTT TIDY, M.B., Professor of Chemistry and Medical Jurisprudence at the London Hospital, 8vo, 16s.

VACHER.—**A Primer of Chemistry**, including Analysis. By ARTHUR VACHER. 18mo, 1s.

VALENTIN.—**Chemical Tables for the Lecture-room and Laboratory.** By WILLIAM G. VALENTIN, F.C.S. In Five large Sheets, 5s. 6d.

CHEMISTRY—*continued.*

VALENTIN AND HODGKINSON.—A Course of

Qualitative Chemical Analysis. By W. G. VALENTIN, F.C.S. Sixth Edition by W. R. HODGKINSON, Ph.D. (Wurzburg), Senior Demonstrator of Practical Chemistry in the Science Schools, South Kensington, and H. M. CHAPMAN, Assistant Demonstrator. With Engravings. 8vo, 8s. 6d.

WATTS.—Physical and Inorganic Chemistry.

By HENRY WATTS, B.A., F.R.S. (being Vol. I. of the Thirteenth Edition of Fownes' Manual of Chemistry). With 150 Wood Engravings, and Coloured Plate of Spectra. Crown 8vo, 9s.

By the same Author.

Chemistry of Carbon - Compounds, or

Organic Chemistry (being Vol. II. of the Twelfth Edition of Fownes' Manual of Chemistry). With Engravings. Crown 8vo, 10s.

CHILDREN, DISEASES OF.

DAY.—A Treatise on the Diseases of Children.

For Practitioners and Students. By WILLIAM H. DAY, M.D., Physician to the Samaritan Hospital for Women and Children. Crown 8vo, 12s. 6d.

ELLIS.—A Practical Manual of the Diseases

of Children. By EDWARD ELLIS, M.D., late Senior Physician to the Victoria Hospital for Sick Children. With a Formulary. Fourth Edition. Crown 8vo, 10s.

SMITH.—On the Wasting Diseases of Infants

and Children. By EUSTACE SMITH, M.D., F.R.C.P., Physician to H.M. the King of the Belgians, and to the East London Hospital for Children. Fourth Edition. Post 8vo, 8s. 6d.

By the same Author.

A Practical Treatise on Disease in Chil-

dren. 8vo, 22s.

STEINER.—Compendium of Children's Dis-

eases; a Handbook for Practitioners and Students. By JOHANN STEINER, M.D. Translated by LAWSON TAIT, F.R.C.S., Surgeon to the Birmingham Hospital for Women, &c. 8vo, 12s. 6d.

DENTISTRY.

GORGAS.—**Dental Medicine: a Manual of**
Dental Materia Medica and Therapeutics, for Practitioners and
Students. By FERDINAND J. S. GORGAS, A.M., M.D., D.D.S., Professor
of Dentistry in the University of Maryland; Editor of "Harris's
Principles and Practice of Dentistry," &c. Royal 8vo, 14s.

SEWILL.—**The Student's Guide to Dental**
Anatomy and Surgery. By HENRY E. SEWILL, M.R.C.S., L.D.S., late
Dental Surgeon to the West London Hospital. Second Edition.
With 78 Engravings. Fcap. 8vo, 5s. 6d.

STOCKEN.—**Elements of Dental Materia Medica**
and Therapeutics, with Pharmacopœia. By JAMES STOCKEN, L.D.S.R.C.S.,
late Lecturer on Dental Materia Medica and Therapeutics and Dental
Surgeon to the National Dental Hospital; assisted by THOMAS GADDES,
L.D.S. Eng. and Edin. Third Edition. Fcap. 8vo, 7s. 6d.

TAFT.—**A Practical Treatise on Operative**
Dentistry. By JONATHAN TAFT, D.D.S., Professor of Operative Surgery
in the Ohio College of Dental Surgery. Third Edition. With 134
Engravings. 8vo, 18s.

TOMES (C. S.).—**Manual of Dental Anatomy,**
Human and Comparative. By CHARLES S. TOMES, M.A., F.R.S.
Second Edition. With 191 Engravings. Crown 8vo, 12s. 6d.

TOMES (J. and C. S.).—**A Manual of Dental**
Surgery. By JOHN TOMES, M.R.C.S., F.R.S., and CHARLES S. TOMES,
M.A., M.R.C.S., F.R.S.; Lecturer on Anatomy and Physiology at the
Dental Hospital of London. Third Edition. With many Engravings,
Crown 8vo. [In the press.]

EAR, DISEASES OF.

BURNETT.—**The Ear: its Anatomy, Physio-**
logy, and Diseases. A Practical Treatise for the Use of Medical
Students and Practitioners. By CHARLES H. BURNETT, M.D., Aural
Surgeon to the Presbyterian Hospital, Philadelphia. With 87 Engra-
vings. 8vo, 18s.

DALBY.—**On Diseases and Injuries of the Ear.**
By WILLIAM B. DALBY, F.R.C.S., Aural Surgeon to, and Lecturer on
Aural Surgery at, St. George's Hospital. Second Edition. With
Engravings. Fcap. 8vo, 6s. 6d.

11, NEW BURLINGTON STREET.

EAR, DISEASES OF—*continued.*

JONES.—**A Practical Treatise on Aural Surgery.** By H. MACNAUGHTON JONES, M.D., Professor of the Queen's University in Ireland, late Surgeon to the Cork Ophthalmic and Aural Hospital. Second Edition. With 63 Engravings. Crown 8vo, 8s. 6d.

By the same Author.

Atlas of the Diseases of the Membrana Tympani. In Coloured Plates, containing 59 Figures. With Explanatory Text. Crown 4to, 21s.

FORENSIC MEDICINE.

OGSTON.—**Lectures on Medical Jurisprudence.**

By FRANCIS OGSTON, M.D., late Professor of Medical Jurisprudence and Medical Logic in the University of Aberdeen. Edited by FRANCIS OGSTON, Jun., M.D., late Lecturer on Practical Toxicology in the University of Aberdeen. With 12 Plates. 8vo, 18s.

TAYLOR.—**The Principles and Practice of**

Medical Jurisprudence. By ALFRED S. TAYLOR, M.D., F.R.S. Third Edition, revised by THOMAS STEVENSON, M.D., F.R.C.P., Lecturer on Chemistry and Medical Jurisprudence at Guy's Hospital; Examiner in Chemistry at the Royal College of Physicians; Official Analyst to the Home Office. With 188 Engravings. 2 Vols. 8vo, 31s. 6d.

By the same Author.

A Manual of Medical Jurisprudence.

Tenth Edition. With 55 Engravings. Crown 8vo, 14s.

ALSO,

On Poisons, in relation to Medical Jurisprudence and Medicine. Third Edition. With 104 Engravings. Crown 8vo, 16s.

TIDY AND WOODMAN.—**A Handy-Book of**

Forensic Medicine and Toxicology. By C. MEYMOTT TIDY, M.B.; and W. BATHURST WOODMAN, M.D., F.R.C.P. With 8 Lithographic Plates and 116 Wood Engravings. 8vo, 31s. 6d.

HYGIENE.

PARKES.—A Manual of Practical Hygiene.

By EDMUND A. PARKES, M.D., F.R.S. Sixth Edition by F. DECHAUMONT, M.D., F.R.S., Professor of Military Hygiene in the Army Medical School. With 9 Plates and 103 Engravings. 8vo, 18s.

WILSON.—A Handbook of Hygiene and Sanitary Science.

By GEORGE WILSON, M.A., M.D., F.R.S.E., Medical Officer of Health for Mid Warwickshire. Fifth Edition. With Engravings. Crown 8vo, 10s. 6d.

MATERIA MEDICA AND THERAPEUTICS.

BINZ AND SPARKS.—The Elements of Therapeutics; a Clinical Guide to the Action of Medicines.

By C. BINZ, M.D., Professor of Pharmacology in the University of Bonn. Translated and Edited with Additions, in conformity with the British and American Pharmacopœias, by EDWARD I. SPARKS, M.A., M.B., F.R.C.P. Lond. Crown 8vo, 8s. 6d.

OWEN.—A Manual of Materia Medica; incorporating the Author's "Tables of Materia Medica."

By ISAMBARD OWEN, M.D., Lecturer on Materia Medica and Therapeutics to St. George's Hospital. Crown 8vo, 6s.

ROYLE AND HARLEY.—A Manual of Materia Medica and Therapeutics.

By J. FORBES ROYLE, M.D., F.R.S., and JOHN HARLEY, M.D., F.R.C.P., Physician to, and Joint Lecturer on Clinical Medicine at, St. Thomas's Hospital. Sixth Edition. With 139 Engravings. Crown 8vo, 15s.

THOROWGOOD.—The Student's Guide to Materia Medica and Therapeutics.

By JOHN C. THOROWGOOD, M.D., F.R.C.P., Lecturer on Materia Medica at the Middlesex Hospital. Second Edition. With Engravings. Fcap. 8vo, 7s.

WARING.—A Manual of Practical Therapeutics.

By EDWARD J. WARING, C.I.E., M.D., F.R.C.P. Third Edition. Fcap. 8vo, 12s. 6d.

MEDICINE.

ARCLAY.—A Manual of Medical Diagnosis.

By A. WHYTE BARCLAY, M.D., F.R.C.P., late Physician to, and Lecturer on Medicine at, St. George's Hospital. Third Edition. Fcap. 8vo, 10s. 6d.

HARTERIS.—The Student's Guide to the

Practice of Medicine. By MATTHEW CHARTERIS, M.D., Professor of Materia Medica, University of Glasgow; Physician to the Royal Infirmary. With Engravings on Copper and Wood. Third Edition. Fcap. 8vo, 7s.

ENWICK.—The Student's Guide to Medical

Diagnosis. By SAMUEL FENWICK, M.D., F.R.C.P., Physician to the London Hospital. Fifth Edition. With 111 Engravings. Fcap. 8vo, 7s.

By the same Author.

The Student's Outlines of Medical Treat-

ment. Second Edition. Fcap. 8vo, 7s.

LINT.—Clinical Medicine: a Systematic Treat-

tise on the Diagnosis and Treatment of Disease. By AUSTIN FLINT, M.D., Professor of the Principles and Practice of Medicine, &c., in Bellevue Hospital Medical College. 8vo, 20s.

HALL.—Synopsis of the Diseases of the Larynx,

Lungs, and Heart: comprising Dr. Edwards' Tables on the Examination of the Chest. With Alterations and Additions. By F. DE HAVILLAND HALL, M.D., F.R.C.P., Assistant-Physician to the Westminster Hospital. Royal 8vo, 2s. 6d.

SANSOM.—Manual of the Physical Diagnosis

of Diseases of the Heart, including the use of the Sphygmograph and Cardiograph. By A. E. SANSOM, M.D., F.R.C.P., Assistant-Physician to the London Hospital. Third Edition. With 47 Woodcuts. Fcap. 8vo, 7s. 6d.

WARNER.—Student's Guide to Clinical Medi-

cine and Case-Taking. By FRANCIS WARNER, M.D., F.R.C.P., Assistant-Physician to the London Hospital. Second Edition. Fcap. 8vo, 5s.

WEST.—How to Examine the Chest: being a

Practical Guide for the Use of Students. By SAMUEL WEST, M.D., M.R.C.P., Physician to the City of London Hospital for Diseases of the Chest, &c. With 42 Engravings. Fcap. 8vo, 5s.

J. & A. Churchill's Medical Class Books.

MEDICINE—continued.

WHITTAKER.—Student's Primer on the Urine.

By J. TRAVIS WHITTAKER, M.D., Clinical Demonstrator at the Royal Infirmary, Glasgow. With Illustrations, and 16 Plates etched on Copper. Post 8vo, 4s. 6d.

MIDWIFERY.

BARNES.—Lectures on Obstetric Operations,

including the Treatment of Hæmorrhage, and forming a Guide to the Management of Difficult Labour. By ROBERT BARNES, M.D., F.R.C.P., Obstetric Physician to, and Lecturer on Diseases of Women, &c., at, St. George's Hospital. Third Edition. With 124 Engravings. 8vo, 18s.

CLAY.—The Complete Handbook of Obstetric

Surgery; or, Short Rules of Practice in every Emergency, from the Simplest to the most formidable Operations connected with the Science of Obstetrics. By CHARLES CLAY, M.D., late Senior Surgeon to, and Lecturer on Midwifery at, St. Mary's Hospital, Manchester. Third Edition. With 91 Engravings. Fcap. 8vo, 6s. 6d.

RAMSBOTHAM.—The Principles and Practice

of Obstetric Medicine and Surgery. By FRANCIS H. RAMSBOTHAM, M.D., formerly Obstetric Physician to the London Hospital. Fifth Edition. With 120 Plates, forming one thick handsome volume. 8vo, 22s.

REYNOLDS.—Notes on Midwifery: specially

designed to assist the Student in preparing for Examination. By J. J. REYNOLDS, L.R.C.P., M.R.C.S. Fcap. 8vo, 4s.

ROBERTS.—The Student's Guide to the Practice

of Midwifery. By D. LLOYD ROBERTS, M.D., F.R.C.P., Physician to St. Mary's Hospital, Manchester. Third Edition. With 2 Coloured Plates and 127 Engravings. Fcap. 8vo, 7s. 6d.

SCHROEDER.—A Manual of Midwifery; includ-

ing the Pathology of Pregnancy and the Puerperal State. By KARL SCHROEDER, M.D., Professor of Midwifery in the University of Erlangen. Translated by CHARLES H. CARTER, M.D. With Engravings. 8vo, 12s. 6d.

SWAYNE.—Obstetric Aphorisms for the Use of

Students commencing Midwifery Practice. By JOSEPH G. SWAYNE, M.D., Lecturer on Midwifery at the Bristol School of Medicine. Eighth Edition. With Engravings. Fcap. 8vo, 3s. 6d.

11, NEW BURLINGTON STREET.

MICROSCOPY.

CARPENTER.—The Microscope and its Revela-

tions. By WILLIAM B. CARPENTER, C.B., M.D., F.R.S. Sixth Edition. With 26 Plates, a Coloured Frontispiece, and more than 500 Engravings. Crown 8vo, 16s.

MARSH. — Microscopical Section-Cutting : a

Practical Guide to the Preparation and Mounting of Sections for the Microscope, special prominence being given to the subject of Animal Sections. By Dr. SYLVESTER MARSH. Second Edition. With 17 Engravings. Fcap. 8vo, 3s. 6d.

MARTIN.—A Manual of Microscopic Mounting.

By JOHN H. MARTIN, Member of the Society of Public Analysis, &c. Second Edition. With several Plates and 144 Engravings. 8vo, 7s. 6d.

OPHTHALMOLOGY.

HIGGENS.—Hints on Ophthalmic Out-Patient

Practice. By CHARLES HIGGENS, F.R.C.S., Ophthalmic Surgeon to, and Lecturer on Ophthalmology at, Guy's Hospital. Second Edition. Fcap. 8vo, 3s.

ONES.—A Manual of the Principles and

Practice of Ophthalmic Medicine and Surgery. By T. WHARTON JONES, F.R.C.S., F.R.S., late Ophthalmic Surgeon and Professor of Ophthalmology to University College Hospital. Third Edition. With 9 Coloured Plates and 173 Engravings. Fcap. 8vo, 12s. 6d.

NETTLESHIP.—The Student's Guide to Diseases

of the Eye. By EDWARD NETTLESHIP, F.R.C.S., Ophthalmic Surgeon to, and Lecturer on Ophthalmic Surgery at, St. Thomas's Hospital. Third Edition. With 157 Engravings, and a Set of Coloured Papers illustrating Colour-blindness. Fcap. 8vo, 7s. 6d.

TOSSWILL.—Diseases and Injuries of the Eye

and Eyelids. By LOUIS H. TOSSWILL, B.A., M.B. Cantab., M.R.C.S., Surgeon to the West of England Eye Infirmary, Exeter. Fcap. 8vo, 2s. 6d.

WOLFE.—On Diseases and Injuries of the Eye :

a Course of Systematic and Clinical Lectures to Students and Medical Practitioners. By J. R. WOLFE, M.D., F.R.C.S.E., Senior Surgeon to the Glasgow Ophthalmic Institution, Lecturer on Ophthalmic Medicine and Surgery in Anderson's College. With 10 Coloured Plates, and 120 Wood Engravings, 8vo, 21s.

PATHOLOGY.

JONES AND SIEVEKING.—*A Manual of Pathological Anatomy.* By C. HANDFIELD JONES, M.B., F.R.S., and EDWARD H. SIEVEKING, M.D., F.R.C.P. Second Edition. Edited, with considerable enlargement, by J. F. PAYNE, M.B., Assistant-Physician and lecturer on General Pathology at St. Thomas's Hospital. With 195 Engravings. Crown 8vo, 16s.

LANCEREAUX.—*Atlas of Pathological Anatomy.* By Dr. LANCEREAUX. Translated by W. S. GREENFIELD, M.D., Professor of Pathology in the University of Edinburgh. With 70 Coloured Plates. Imperial 8vo, £5 5s.

VIRCHOW.—*Post-Mortem Examinations: a Description and Explanation of the Method of Performing them, with especial reference to Medico-Legal Practice.* By Professor RUDOLPH VIRCHOW, Berlin Charité Hospital. Translated by Dr. T. B. SMITH. Second Edition, with 4 Plates. Fcap. 8vo, 3s. 6d.

WILKS AND MOXON.—*Lectures on Pathological Anatomy.* By SAMUEL WILKS, M.D., F.R.S., Physician to, and late Lecturer on Medicine at, Guy's Hospital; and WALTER MOXON, M.D., F.R.C.P., Physician to, and Lecturer on the Practice of Medicine at, Guy's Hospital. Second Edition. With 7 Steel Plates. 8vo, 18s.

PSYCHOLOGY.

BUCKNILL AND TUKE.—*A Manual of Psychological Medicine: containing the Lunacy Laws, Nosology, Ætiology, Statistics, Description, Diagnosis, Pathology, and Treatment of Insanity, with an Appendix of Cases.* By JOHN C. BUCKNILL, M.D., F.R.S., and D. HACK TUKE, M.D., F.R.C.P. Fourth Edition with 12 Plates (30 Figures). 8vo, 25s.

CLOUSTON.—*Clinical Lectures on Mental Diseases.* By THOMAS S. CLOUSTON, M.D., and F.R.C.P. Edin.; Lecturer on Mental Diseases in the University of Edinburgh. With 8 Plates (6 Coloured). Crown 8vo, 12s. 6d.

MANN.—*A Manual of Psychological Medicine and Allied Nervous Disorders.* By EDWARD C. MANN, M.D., Member of the New York Medico-Legal Society. With Plates. 8vo, 24s.

11, NEW BURLINGTON STREET.

PHYSIOLOGY.

ARPENTER.—**Principles of Human Physiology.** By WILLIAM B. CARPENTER, C.B., M.D., F.R.S. Ninth Edition. Edited by Henry Power, M.B., F.R.C.S. With 3 Steel Plates and 377 Wood Engravings. 8vo, 31s. 6d.

ALTON.—**A Treatise on Human Physiology :** designed for the use of Students and Practitioners of Medicine. By JOHN C. DALTON, M.D., Professor of Physiology and Hygiene in the College of Physicians and Surgeons, New York. Seventh Edition. With 252 Engravings. Royal 8vo, 20s.

REY.—**The Histology and Histo-Chemistry of Man.** A Treatise on the Elements of Composition and Structure of the Human Body. By HEINRICH FREY, Professor of Medicine in Zurich. Translated by ARTHUR E. BARKER, Assistant-Surgeon to the University College Hospital. With 608 Engravings. 8vo, 21s.

ANDERSON.—**Handbook for the Physiological Laboratory :** containing an Exposition of the fundamental facts of the Science, with explicit Directions for their demonstration. By J. BURDON SANDERSON, M.D., F.R.S.; E. KLEIN, M.D., F.R.S.; MICHAEL FOSTER, M.D., F.R.S.; and T. LAUDER BRUNTON, M.D., F.R.S. 2 Vols., with 123 Plates. 8vo, 24s.

YEO.—**A Manual of Physiology for the Use of Junior Students of Medicine.** By GERALD F. YEO, M.D., F.R.C.S., Professor of Physiology in King's College, London. With 301 Engravings. Crown 8vo, 14s.

SURGERY.

BELLAMY.—**The Student's Guide to Surgical Anatomy ;** a Description of the more important Surgical Regions of the Human Body, and an Introduction to Operative Surgery. By EDWARD BELLAMY, F.R.C.S., and Member of the Board of Examiners ; Surgeon to, and Lecturer on Anatomy at, Charing Cross Hospital. Second Edition. With 76 Engravings. Fcap. 8vo, 7s.

BRYANT.—**A Manual for the Practice of Surgery.** By THOMAS BRYANT, F.R.C.S., Surgeon to, and Lecturer on Surgery at, Guy's Hospital. Fourth Edition. With 750 Illustrations (many being coloured), and including 6 Chromo-Lithographic Plates. 2 Vols. Crown 8vo, 32s.

SURGERY—continued.

CLARK AND WAGSTAFFE. — **Outlines of Surgery and Surgical Pathology.** By F. LE GROS CLARK, F.R.C.S., F.R.S., Consulting Surgeon to St. Thomas's Hospital. Second Edition. Revised and expanded by the Author, assisted by W. W. WAGSTAFFE, F.R.C.S., Assistant Surgeon to St. Thomas's Hospital. 8vo, 10s. 6d.

DRUITT. — **The Surgeon's Vade-Mecum; a Manual of Modern Surgery.** By ROBERT DRUITT, F.R.C.S. Eleventh Edition. With 369 Engravings. Fcap. 8vo, 14s.

FERGUSON. — **A System of Practical Surgery.** By Sir WILLIAM FERGUSON, Bart., F.R.C.S., F.R.S., late Surgeon and Professor of Clinical Surgery to King's College Hospital. With 463 Engravings. Fifth Edition. 8vo, 21s.

HEATH. — **A Manual of Minor Surgery and Bandaging, for the use of House-Surgeons, Dressers, and Junior Practitioners.** By CHRISTOPHER HEATH, F.R.C.S., Holme Professor of Clinical Surgery in University College and Surgeon to the Hospital. Seventh Edition. With 129 Engravings. Fcap. 8vo, 6s.

By the same Author.

A Course of Operative Surgery: with Twenty Plates (containing many figures) drawn from Nature by M. LÉVEILLÉ, and Coloured. Second Edition. Large 8vo, 30s.

ALSO,

The Student's Guide to Surgical Diagnosis. Second Edition. Fcap. 8vo, 6s. 6d.

MAUNDER. — **Operative Surgery.** By Charles F. MAUNDER, F.R.C.S., late Surgeon to, and Lecturer on Surgery at, the London Hospital. Second Edition. With 164 Engravings. Post 8vo, 6s.

SOUTHAM. — **Regional Surgery: including Surgical Diagnosis.** A Manual for the use of Students. By FREDERICK A. SOUTHAM, M.A., M.B. Oxon, F.R.C.S., Assistant-Surgeon to the Royal Infirmary, and Assistant-Lecturer on Surgery in the Owen's College School of Medicine, Manchester.

Part I. The Head and Neck. Crown 8vo, 6s. 6d.

„ II. The Upper Extremity and Thorax. Crown 8vo, 7s. 6d.

11, NEW BURLINGTON STREET.

TERMINOLOGY.

DUNGLISON.—**Medical Lexicon : a Dictionary** of Medical Science, containing a concise Explanation of its various Subjects and Terms, with Accentuation, Etymology, Synonyms, &c. By ROBERT DUNGLISON, M.D. New Edition, thoroughly revised by RICHARD J. DUNGLISON, M.D. Royal 8vo, 28s.

MAYNE.—**A Medical Vocabulary : being an** Explanation of all Terms and Phrases used in the various Departments of Medical Science and Practice, giving their Derivation, Meaning, Application, and Pronunciation. By ROBERT G. MAYNE, M.D., LL.D., and JOHN MAYNE, M.D., L.R.C.S.E. Fifth Edition. Crown 8vo, 10s. 6d.

WOMEN, DISEASES OF.

BARNES.—**A Clinical History of the Medical** and Surgical Diseases of Women. By ROBERT BARNES, M.D., F.R.C.P., Obstetric Physician to, and Lecturer on Diseases of Women, &c., at, St. George's Hospital. Second Edition. With 181 Engravings. 8vo, 28s.

COURTY.—**Practical Treatise on Diseases of** the Uterus, Ovaries, and Fallopian Tubes. By Professor COURTY, Montpellier. Translated from the Third Edition by his Pupil, AGNES M'LAREN, M.D., M.K.Q.C.P. With Preface by Dr. MATTHEWS DUNCAN. With 424 Engravings. 8vo, 24s.

DUNCAN.—**Clinical Lectures on the Diseases** of Women. By J. MATTHEWS DUNCAN, M.D., F.R.C.P., F.R.S.E., Obstetric Physician to St. Bartholomew's Hospital. Second Edition, with Appendices. 8vo, 14s.

EMMET.—**The Principles and Practice of** Gynæcology. By THOMAS ADDIS EMMET, M.D., Surgeon to the Woman's Hospital of the State of New York. With 130 Engravings. Royal 8vo, 24s.

GALABIN.—**The Student's Guide to the Dis-**eases of Women. By ALFRED L. GALABIN, M.D., F.R.C.P., Obstetric Physician to, and Lecturer on Obstetric Medicine at, Guy's Hospital. Third Edition. With 78 Engravings. Fcap. 8vo, 7s. 6d.

WOMEN, DISEASES OF—*continued.*

REYNOLDS.—Notes on Diseases of Women.

Specially designed to assist the Student in preparing for Examination. By J. J. REYNOLDS, L.R.C.P., M.R.C.S. Second Edition. Fcap. 8vo 2s. 6d.

SAVAGE.—The Surgery of the Female Pelvic

Organs. By HENRY SAVAGE, M.D., Lond., F.R.C.S., one of the Consulting Medical Officers of the Samaritan Hospital for Women. Fifth Edition, with 17 Lithographic Plates (15 Coloured), and 52 Woodcuts. Royal 4to, 35s.

SMITH.—Practical Gynæcology: a Handbook

of the Diseases of Women. By HEYWOOD SMITH, M.D., Physician to the Hospital for Women and to the British Lying-in Hospital. With Engravings. Second Edition. Crown 8vo. [In preparation.]

WEST AND DUNCAN.—Lectures on the Dis-

eases of Women. By CHARLES WEST, M.D., F.R.C.P. Fourth Edition. Revised and in part re-written by the Author, with numerous additions by J. MATTHEWS DUNCAN, M.D., F.R.C.P., F.R.S.E., Obstetric Physician to St. Bartholomew's Hospital. 8vo, 16s.

ZOOLOGY.

CHAUVEAU AND FLEMING.—The Compara-

tive Anatomy of the Domesticated Animals. By A. CHAUVEAU, Professor at the Lyons Veterinary School; and GEORGE FLEMING, Veterinary Surgeon, Royal Engineers. With 450 Engravings. 8vo, 31s. 6d.

HUXLEY.—Manual of the Anatomy of Inverte-

brated Animals. By THOMAS H. HUXLEY, LL.D., F.R.S. With 156 Engravings. Post 8vo, 16s.

By the same Author.

Manual of the Anatomy of Vertebrated

Animals. With 110 Engravings. Post 8vo, 12s.

WILSON.—The Student's Guide to Zoology:

a Manual of the Principles of Zoological Science. By ANDREW WILSON, Lecturer on Natural History, Edinburgh. With Engravings. Fcap. 8vo, 6s. 6d.

11, NEW BURLINGTON STREET.



